

On May 14, 1946, S. G. Smith captured a living individual of this species as it came aboard the cruiser U. S. S. Pasadena off the California coast. At the time the sparrow was discovered, the cruiser was fifty miles south of the Golden Gate and thirty miles off-shore. According to Smith, the bird came aboard in an exhausted condition and died just prior to the time the cruiser put into Los Angeles harbor. Smith gave the specimen to Mrs. Mary V. Hood of Los Angeles who in turn brought it to the Los Angeles Museum.

The specimen, an adult male (L. A. Museum Coll. 20332), was compared with a series of skins of *ochracea* from the collections of the Los Angeles Museum and the Museum of Vertebrate Zoology and is definitely referable to this race.—KENNETH E. STAGER, *Los Angeles Museum, Los Angeles, California, September 1, 1946.*

Reactions of Cliff Swallows to a Buteonid Hawk.—On May 18, 1946, the author and four other bird students were observing a colony of Cliff Swallows (*Petrochelidon albifrons*) on the north-facing lava cliffs of Saw Mill Ravine at Cherokee, Butte County, California. The colony included several hundred birds, mostly busy at nest building. A large buteonid hawk flying by within about 100 feet of the cliffs caused considerable excitement among the birds. Their flight became more rapid and erratic and some swallows were seen to dive at the hawk, although none of them was actually seen to come in contact. The hawk alighted in a tree about 300 yards from the colony and the swallows followed to a position above the tree, where they milled around excitedly in a well defined spherical flock. They gradually quieted down and in about fifteen minutes were back to nest building. Half an hour later, a Turkey Vulture (*Cathartes aura*) was seen flying past within fifty or sixty feet of the cliffs, but none of the swallows exhibited concern. But an hour or so later, most of the swallows formed a milling flock above one of the observers who approached more closely to the nests than had others of the party.

The concern exhibited by the swallows over the presence of the hawk is at variance with the statement in Bent's "Life Histories of North American Flycatchers, Larks, Swallows and their Allies" (1942:480), that "the appearance of a hawk in the vicinity of a colony of Cliff Swallows never creates any evidence of excitement, . . ." However, it does not necessarily indicate that they recognized the species as an enemy. It is well known that the feeding habits of hawks varies with the situation, and this individual may have been recognized because of having previously attacked swallows.—J. BRUCE KIMSEY, *Chico, California, August 21, 1946.*

A New Flycatcher of the Genus *Monarcha* from the Bismarck Archipelago.—The widespread and variable flycatcher, *Monarcha cinerascens*, occurs as a common resident on small off-shore islands and along the coasts of larger ones from Timor around the north coast of New Guinea to the northern Solomon Islands. The species has been recorded from all parts of the Bismarck Archipelago and several subspecies have been separated in that region. It is not surprising, therefore, to find a well marked race inhabiting the hitherto ornithologically unexplored islet of Tench in the St. Matthias Group. This new race may be known as

***Monarcha cinerascens tenchi*, new subspecies**

Type.—Adult male, no. 90235 Mus. Vert. Zool., from Tench Island, St. Matthias Group, Bismarck Archipelago, collected August 19, 1944, by Charles G. Sibley, orig. no. 2427.

Diagnosis.—Intermediate in tone of coloration between *M. c. perpallidus* and *M. c. impediens*, the abdomen of *tenchi* being Ochraceous Tawny (Ridgway, Color Standards and Color Nomenclature, 1912), that of *perpallidus* Cinnamon Buff, and that of *impediens* Sanford's Brown.

Range.—The island of Tench, located 30 miles east of Emirau Island in the St. Matthias Group, Bismarck Archipelago.

Specimens examined.—Two adults (male and female) from Tench have been compared with fourteen males and eleven females from Emirau and Mussau and with seven males and five females from Feni and Green islands. In addition, one female from New Hanover, one female from Watom, one male from Manus, and two males and one female from Ahu, Ninigo Group, have been examined.

The color of the abdomen of *tenchi* exactly matches that of the three specimens of *M. c. fulviventris* from Ahu, Ninigo Group. These two races are separable by differences in dimensions as shown in the table.

The single specimen from Manus, Admiralty Islands, agrees in coloration with those from Ahu although the bill is larger; it is tentatively referred to *fulviventris*. This is also true of the single specimen from Watom which resembles *fulviventris* in color but has an even larger bill than the specimen from Manus. The specimen from New Hanover has a smaller bill than any of the 25 adults from the St. Matthias Group, but it is possibly an immature bird. A summary of the characters of the forms examined is presented in the table. The color of the posterior underparts is designated by comparison with Ridgway (*op. cit.*) and also with Maerz and Paul (Dictionary of Color, McGraw-Hill Book Co.,