



Field Records

May 1986

by George W. Gove, Robert H. Stymeist, Lee E. Taylor

May 1986 was dry and cloudy with near normal temperatures. The average was 58.4°F; the first half of the month was cool, averaging about six degrees below normal, and birders twitched, waiting for warmer weather and the first "wave." The second half of the month was warmer than usual. The first 90° day came on May 19, twenty-two days earlier than average; the heat also tied the record for that day set in 1949. Rain totaled 1.21 inches, 2.31 inches less than the average for May. Sunshine totaled 51 percent of possible, a big improvement over May of 1985. Southwest winds were recorded on eight days - six of these days occurred after May 16, causing the rather late migration waves noted this year from many areas.

R.H.S.

LOONS THROUGH CORMORANTS

A Red-necked Grebe coming into breeding plumage was noted from Plum Island on May 13, whereas reports of Pied-billed Grebes in spring continued to be poor with only two individuals noted. In Nantucket Sound, the usual small numbers of tubenoses were observed with Sooty Shearwaters showing up in good numbers by the end of the month. Three Great Cormorants were found all month in North Scituate, and Double-crested Cormorants continued to show increased populations, especially in Boston Harbor. R.H.S.

| SPECIES/DATE | LOCATION | NUMBER | OBSERVERS | MAY 1986 |
|-----------------------------|-------------------------------------|------------------|---------------------------------|----------|
| Red-throated Loon: | | | | |
| 11,13 | P'town, Eastham (F.E.) | 8+, 8+ | B.Nikula# + v.o. | |
| 18,20 | Wellfleet, Duxbury | 2, 2 | D.Morimoto#, R.Forster | |
| 25,26 | Plum Island, Nahant | 2, 2 | L.Taylor#, R.Forster | |
| 29 | S.Monomoy | 3 | J.Russell | |
| Common Loon: | | | | |
| 10,10-11 | E.Middleboro, E.Boston | 1 mig., 2 | W.Petersen, S.Zendeh | |
| 10,14 | Acoaxet, Braintree | 3, 2 | G.d'Entremont# | |
| 17,18 | Clinton, Quincy-P'town | 1, 30 | M.Lynch#, D.Morimoto# | |
| 23-25,24 | M.V., P.I. | 3, 3 | BBC(J.Barton), BBC(F.Bouchard) | |
| Pied-billed Grebe: | | | | |
| 4,29 | Boston(Muddy R.), S.Monomoy | 1, 1 | P.Thayer, B.Nikula# | |
| Horned Grebe: | | | | |
| 24 | Plum Island | 3 | BBC(F.Bouchard) | |
| Red-necked Grebe (details): | | | | |
| 13 | Plum Island | 1 | T.Collins + A.Sanborn | |
| Northern Fulmar: | | | | |
| 15,17 | E of Chatham: 10 mi., 2 mi. | 6, 10 | R.Humphrey# | |
| 26 | Nantucket Sound | 1 | S.Perkins | |
| Sooty Shearwater: | | | | |
| 25 | Stellwagen Bank | 2 | C.Leahy | |
| 29,30;31 | Nant. Sound; N.Monomoy | 20, 100; 3 | R.Humphrey#, W.Petersen# | |
| Manx Shearwater: | | | | |
| 12 | Nantucket | 2 | S.Perkins | |
| Wilson's Storm-Petrel: | | | | |
| 15;25 | 10 mi. E. of Chatham; P.I., Stellw. | 100+; 6+, 125 | R.Humphrey#;R.Stymeist#,C.Leahy | |
| 26,28;29 | Nant.Sound, Nant.; P'town | 200, 75; 2 | S.Perkins; B.Nikula | |
| Northern Gannet: | | | | |
| 12,13 | Dennis, Eastham(F.E.) | 50, 300 | B.Nikula | |
| 17,18 | Nant., Wellfleet | 475, 1 | M.Litchfield, D.Morimoto# | |
| Great Cormorant: | | | | |
| thr.,29 | N.Scituate, S.Monomoy | 3+, 2 | G.d'Entremont+v.o.,B.Nikula# | |
| Double-crested Cormorant: | | | | |
| thr. | S.Dartmouth | max.35 (5/10) | LCES | |
| thr. | Saugus-Boston (train) | max.60 (5/27+29) | J.Berry | |

| SPECIES/DATE | LOCATION | NUMBER | OBSERVERS | MAY 1986 |
|--|------------------------------|------------------|--------------------------------|----------|
| Double-crested Cormorant (continued): | | | | |
| 10,11,25 | E.Boston (Belle Isle) | 66, 156, 15 | S.Zendeh | |
| 11,14 | Gloucester, Halifax | 30, 52 | BBC(J.Nove), K.Anderson | |
| 15,17 | Mt.A., Newburyport | 40, 60 | R.Stymeist#, BBC(B.Drummond) | |
| 25 | Nahant (Egg Rock) | 1000s | R.Stymeist + G.Gove | |
| <u>HERONS</u> | | | | |
| Sixteen American Bittern were reported this May, compared with only eight last year, but Least Bittern was reported from only one location, compared with five reports in May 1985. On Monomoy, 57 active Snowy Egret nests were found along with one active nest of Glossy Ibis, the first confirmed nesting for this species on Cape Cod. Black-crowned Night-Heron nests on South Monomoy totaled 176, a big jump from 70 nests in May 1982. Yellow-crowned Night-Herons were noted from five locations. R.H.S. | | | | |
| American Bittern: | | | | |
| 4,16 | Wayland | 1, 2 | R.Forster | |
| 6 | Peabody, Bolton Flats | 1, 1 | R.Heil, E.Salmela | |
| 8,10 | P.I., E.Boston(Belle Isle) | 1, 1 | J.Grujan, S.Zendeh | |
| 10,10 on | Newbypt, Lynnfield | 1, max.3(5/17) | R.Heil, v.o. | |
| 17 | Bolton Flats, Middleton | 2, 1 | BBC(M.Lynch), J.Berry | |
| 18,26 | Milton(F.M.), Newbypt Harbor | 1, 1 | G.Gove#, G.d'Entremont# | |
| Least Bittern: | | | | |
| 17 on | P.I. | 1+ | S.Surner#+v.o. | |
| Great Blue Heron: | | | | |
| 1,3 | Whitman, Lakeville | 2, 1 | K.Holmes | |
| 8 | Salem-Saugus(train) | 11 | J.Berry | |
| 13,17 | Watertown, Bolton | 6 mig., 5 | R.Stymeist, BBC(M.Lynch) | |
| 22,26 | E.Orleans, Westport | 4, 5 | A.Williams,BBC(T.Athearn) | |
| Great Egret: | | | | |
| 1 | Fall River, Quincy | 1, 1 | K.Holmes, K.Ryan | |
| 3,11,26 | Ipswich | 1, 3, 1 | J.Berry | |
| 7,21 | E.Orleans, Newbypt | 3, 1 | D.+E.Williams,BBC(A.Blaisdell) | |
| 25 | Essex(Island Rd.), Lakeville | 4, 1 | R.Stymeist#, W.Petersen# | |
| Snowy Egret: | | | | |
| thr. | Saugus-Revere(train) | max.11(5/16) | J.Berry | |
| 10,11 | W.Newbury, Squantum | 8, 9 | BBC(S.Bolton), SSBC(A.Leggett) | |
| 31 | S.Monomoy | 57 nests | R.Humphrey + v.o. | |
| Little Blue Heron: | | | | |
| 15,17 | Manchester, Monomoy | 1, 1 | BBC(G.Hotz), BBC(J.Barton) | |
| 25,29 | M.V., Chatham | 2, 1 ad. | BBC(J.Barton), B.Nikula# | |
| Tricolored Heron: | | | | |
| 6,10 | S.Dartmouth, N.Monomoy | 1, 1 | R.Marshall, B.Nikula# | |
| 12,12-17 | Gloucester, Hingham | 1, 1 | C.Leahy, D.Morimoto# + v.o. | |
| 17,24 | S.Monomoy, Essex | 1, 1 | B.Nikula#, R.Heil | |
| 25 on,31 | P.I., Chatham | 1, 1 | M.Lynch#+v.o., B.Nikula# | |
| Cattle Egret: | | | | |
| thr. | Ipswich(Appleton Farm) | max.10(5/12) | J.Berry + v.o. | |
| 8,10 | Concord, Essex | 1, 3 | R.Forster, R.Heil | |
| 25 | Wenham | 2 | M.Lynch, S.Carroll | |
| Green-backed Heron: | | | | |
| 2,7 | Swampscott, Brookline | 2, 1 | L.Sager, H.Wiggin | |
| 10 | Lakeville, P.I. | 1, 2 | K.Holmes, BBC(S.Bolton) | |
| 13 | Watertown, Lynnfield | 1, 1 | R.Stymeist, W.VanCorf# | |
| 17 | Bolton-Oxbow NWR | 5 | M.Lynch + S.Carroll | |
| 17,18 | P.I., Mt.A. | 6, 3 | S.Surner#, S.Carroll# | |
| Black-crowned Night-Heron: | | | | |
| thr. | Watertown(Charles R.) | max.40(5/20) | R.Stymeist# | |
| 31 | S.Monomoy | 176 active nests | R.Humphrey + v.o. | |
| Yellow-crowned Night-Heron: | | | | |
| 6,14 | Peabody, Marshfield | 1 ad., 1 | R.Heil, D.Clapp | |
| 16,20 | Eastham, Jamaica Plain | 1 ad., 1 ad. | M.O'Connor, C.Jackson# | |
| 25 | Martha's Vineyard | 1 ad. | J.Barton# | |
| Glossy Ibis: | | | | |
| thr. | Essex-Ipswich | max.40(5/4) | W.Petersen# + v.o. | |
| 1,6,8 | Revere | 3, 3, 4 | J.Berry | |
| 6 | Bolton Flats, Chatham | 2, 30 | E.Salmela, R.Humphrey | |
| 15,26 | Manchester, S.Harwich | 2, 4 | BBC(G.Hotz), B.Nikula | |
| 31 | S.Monomoy | 1 active nest | R.Humphrey + v.o. | |

WATERFOWL

Two Snow Geese were noted during May, one in Newburyport on May 10, and another in the Quabbin area on May 17; both reports are unusual for May and perhaps represent ailing birds. Brant lingered into the last week of the month in Newburyport Harbor, despite the increased numbers of wind surfers who were using the harbor on a regular basis. Oldsquaw also lingered throughout the month but in numbers much reduced from previous years, a result perhaps of the presence of the wind surfers. A Eurasian Wigeon was found on South Monomoy at the end of the month. Three King Eiders were noted during the month, and two first-year male Harlequin Ducks could be found from May 16 to the end of the month in Nahant. Nahant was also a good vantage point for observing all three scoters, White-winged being the most common. The only Ruddy Ducks reported during the month were found on South Monomoy.

R.H.S.

| SPECIES/DATE | LOCATION | NUMBER | OBSERVERS | MAY 1986 |
|-----------------------------|--|---|--|----------|
| Mute Swan: | | | | |
| thr. 25,31 | Manchester Plymouth, Wareham | 6 13, 19 | G.Hotz# + v.o. BBC(R.Timberlake), L.Robinson | |
| Snow Goose: | | | | |
| 10,17 | Newbypt, Quabbin(Gate 40) | 1, 1 | R.Heil, K.Ryan# | |
| Brant: | | | | |
| thr. 7,22 11,12 31 | Newbypt Harbor E.Orleans Squantum, Nahant N.Monomoy | max.525(5/17) 100, 40 118, 150 1 | H.Wiggins# + v.o. D.+E.Williams SSBC(A.Leggett), D.Lazzaro W.Petersen | |
| Wood Duck: | | | | |
| thr. 10 17,18 | P'town Milford, Topsfield Bolton-Oxbow, Marshfield | max.9 3, 3 ad.+8 yg. 15, 4 | B.Nikula + v.o. R.Hildreth, J.Berry M.Lynch#, BBC(R.Timberlake) | |
| Green-winged Teal: | | | | |
| 3,11 14,17,29 17,18 | Newbypt, Ipswich S.Monomoy Oxbow NWR, P'town | 50, 7 35, 6, 8 15, 1 | BBC(G.d'Entremont), J.Berry B.Nikula# BBC(M.Lynch), D.Morimoto | |
| American Black Duck: | | | | |
| 17,29 17 | S.Monomoy Bolton Flats | 20, 60 10 | B.Nikula# M.Lynch# | |
| Northern Pintail: | | | | |
| 14,17,29 | S.Monomoy | 10, 6, 6 | B.Nikula# | |
| Blue-winged Teal: | | | | |
| 14,17,29 25 | S.Monomoy P.I. | 6, 10, 6 20 | B.Nikula# M.Lynch, S.Carroll | |
| Northern Shoveler: | | | | |
| 14,17 | S.Monomoy | 5, 2 | B.Nikula# | |
| Gadwall: | | | | |
| 3,17,25 14,17,29 31 | P.I. S.Monomoy N.Monomoy | 10, 8, 25+ G.d'Entremont#, D.Morimoto#, M.Lynch# 25, 20, 20 1 | B.Nikula# W.Petersen# | |
| Eurasian Wigeon: | | | | |
| 29 | S.Monomoy | 1 m. | B.Nikula# | |
| American Wigeon: | | | | |
| 14,17,29 | S.Monomoy | 2, 4, 3 | B.Nikula# | |
| Ring-necked Duck: | | | | |
| 14,25 | Peabody, Lakeville | 1 m., 1 m. | R.Heil, W.Petersen# | |
| Greater Scaup: | | | | |
| 4,26 | Newbypt, Salem | 2, 1 | W.Petersen#, R.Forster | |
| Common Eider: | | | | |
| 11, 18 | Gloucester, Manomet-Chatham | 12, 15 | BBC(J.Nove), D.Morimoto# | |
| King Eider: | | | | |
| 17 28-31 | N.Scituate, S.Monomoy Nahant | 1 f., 1 imm. m. 1 m. | P.Donahue#, B.Nikula L.Pivacek | |
| Harlequin Duck: | | | | |
| 16-31 | Nahant | 2 imm. m. | L.Pivacek + v.o. | |
| Oldsquaw: | | | | |
| thr. 31 | Newbypt Harbor Monomoy | max.145 (5/17) 2 | v.o. W.Petersen# | |
| Black Scoter: | | | | |
| 11,17 17,25;17 | Gloucester, S.Monomoy Nahant; Swampscott | 10, 400 6, 4; 2 | BBC(J.Nove), B.Nikula# C.Floyd#; R.Forster | |
| Surf Scoter: | | | | |
| 17,25 18,25 | Nahant Wellfleet, Marblehead | 6, 1 1, 2 | C.Floyd# D.Morimoto#, R.Stymeist# | |
| White-winged Scoter: | | | | |
| 4,11 | Scituate, Gloucester | 100, 27 | G.d'Entremont, BBC(J.Nove) | |

| SPECIES/DATE | LOCATION | NUMBER | OBSERVERS | MAY 1986 |
|----------------------------------|---------------------------|--------------|--------------------------------|----------|
| White-winged Scoter (continued): | | | | |
| 17,29 | S.Monomoy | 2000, 400 | B.Nikula | |
| 17,25 | Nahant | 250, 75 | R.Stymeist# | |
| Common Goldeneye: | | | | |
| 17 | Newbypt, P.I. | 1, 4 | G.Gove#, S.Surner | |
| 28 | Boston Harbor | 1 m. | D.Morimoto | |
| Bufflehead: | | | | |
| 6,8 | Salem | 12, 30+ | J.Berry | |
| 14,17 | S.Monomoy | 25, 10 | B.Nikula# | |
| 15,22 | Manchester, E.Orleans | 7, 3 | BBC(G.Hotz), D., E.+A.Williams | |
| Red-breasted Merganser: | | | | |
| 6,13,20,27 | S.Dartmouth (Allens Pond) | 42, 50, 8, 9 | LCES(R.Marshall) | |
| 10,11 | S.Dartmouth, Gloucester | 56, 17 | LCES(R.Maker), BBC(J.Nove) | |
| 14 | Chatham (Nant. Sound) | 750 | B.Nikula | |
| 19,31 | N.Monomoy | 250, 70 | B.Nikula, W.Petersen | |
| Ruddy Duck: | | | | |
| 14,17,29 | S.Monomoy | 9, 6, 6 | B.Nikula# | |

RAPTORS

An American Swallow-tailed Kite was reported flying over Chatham on May 19, the twelfth spring record in the state. Also of special note from the cape was an adult Mississippi Kite flying over Truro (and photographed) on May 6. This is the eighth record for this species in Massachusetts, and a full report can be found elsewhere in this issue.

We had reports of three immature Bald Eagles, one or more of which were hanging around Chatham off and on this spring. Osprey reports were lacking from their strongholds - Westport and Martha's Vineyard. Other interesting bird-of-prey reports during the month including nesting Northern Goshawks and Red-shouldered Hawks, both of which were successful. A Broad-winged Hawk on Nantucket was unusual for the island. A light-phase Rough-legged Hawk was carefully studied in Saugus on May 1 and 6, and there were two reports of Golden Eagle from Plum Island. Merlin were noted from nine locations and Peregrines from ten. A Peregrine pair was seen off and on in Boston, although no hacking was being done downtown this year.

R.H.S.

Turkey Vulture:

| | | | |
|---------|-----------------------|---------|--------------------------------|
| 3,9 | S.Dartmouth | 2, 6 | LCES (R.Maker) |
| 3,5 | Newbypt, E.Middleboro | 1, 1 | BBC(G.d'Entremont), P.Anderson |
| 6,10;10 | Sharon; Essex | 2, 1; 3 | K.Ryan; R.Heil |
| 15,26 | Westport | 3, 3 | R.Forster, BBC(T.Athearn) |
| 24,26 | Boxford, E.Middleboro | 1, 1 | R.Heil, K.Anderson |
| 29,31 | P'town, Petersham | 3, 6 | B.Nikula#, M.Lynch# |

Osprey:

| | | | |
|-------|----------------------|-------------------------|---------------------------|
| thr. | S.Dartmouth (2 loc.) | max. 5 | LCES(R.Maker, R.Marshall) |
| 10,16 | E.Boston, Lakeville | 3, 4 | S.Zendeh, K.Holmes |
| 17 | Nantucket | 2 pr. | M.Litchfield |
| 31 | Wareham | pr. feeding yg. on nest | L.Robinson |

No reports from Westport and Martha's Vineyard

American Swallow-tailed Kite:

| | | | |
|----|---------|---|---------|
| 19 | Chatham | 1 | R.Hardy |
|----|---------|---|---------|

Mississippi Kite (full details):

| | | | |
|---|-------|-------------|-----------------------|
| 6 | Truro | 1 ad. (ph.) | B.Nikula + R.Humphrey |
|---|-------|-------------|-----------------------|

Bald Eagle:

| | | | |
|-------|----------------------|----------------|----------------------|
| 9,14 | Chatham, Marshfield | 1 imm., 1 imm. | R.Humphrey#, D.Clapp |
| 29,31 | S.Monomoy, N.Monomoy | 1 imm. | J.Russell# + v.o. |

Northern Harrier:

| | | | |
|------|----------------------|----------------------------|----------------------------------|
| thr. | S.Monomoy | 1 m., 2+ f.+ nest w/ 5 yg. | R.Humphrey# |
| 3 | Newbypt, Ipswich | 7, 3 | BBC(d'Entremont), J.Berry |
| 4,5 | Plymouth, Wayland | 2, 1 m. | R.Abrams, G.Gove |
| 17 | Newbypt, P.I., Essex | pr., 2, 2 | C.Floyd#, R.Stymeist#, R.Forster |

Other reports of single individuals from seven locations.

Sharp-shinned Hawk:

| | | | |
|-------|-----------------------|------------------|----------------------------|
| 2,10 | Truro, P.I. | 104, 18 | R.Humphrey#, BBC(S.Bolton) |
| 10,11 | E.Boston (Belle Isle) | 50, 7 | S.Zendeh |
| 24,29 | Milton, P'town | 1, 1 (both mig.) | R.Forster#, P.Trimble# |

Other reports early in the month of 1 or 2 individuals from many locations.

Cooper's Hawk:

| | | | |
|---|------|-------|---------|
| 2 | MNWS | 1 ad. | L.Sager |
|---|------|-------|---------|

Northern Goshawk:

| | | | |
|---------|--------------------|----------------------------|------------------------|
| thr.,18 | E.Middleboro | pr. at nest, 3 yg. at nest | K.Anderson |
| 4 | Holliston, Mt.A. | 1, 1 m. | R.Hildreth, F.Bouchard |
| 11,28 | Lakeville, Ipswich | 1, 1 imm. | K.Holmes, J.Berry |

| SPECIES/DATE | LOCATION | NUMBER | OBSERVERS | MAY 1986 |
|----------------------|-------------------------------|--------------------------|------------------------------|----------|
| Red-shouldered Hawk: | | | | |
| thr., 25 | E.Middleboro | 2-3 ad.+nest, | K.Anderson | |
| thr. | Boxford State Forest | 1 yg. at nest pr.nesting | v.o. | |
| 4,11 | Holliston, Quabbin (G-40) | 1, 2 | R.Hildreth, G.Gove# | |
| 25,31 | Lakeville, Petersham | 5, 1 | W.Petersen#, M.Lynch# | |
| Broad-winged Hawk: | | | | |
| 1,10 | E.Middleboro, E.Boston | 2, 1 | K.Anderson, S.Zendeh | |
| 18,27 | Harwich, Nantucket | 1, 1 ad. | D.Morimoto#, S.Perkins | |
| 29 | Wellfleet, F'town | 9, 15+ | B.Nikula, P.Trimble | |
| 31 | Wareham | 2 | L.Robinson | |
| Red-tailed Hawk: | | | | |
| thr. | Lexington, Newbypt | pr.nesting, pr.nesting | R.Stymeist# | |
| 17 | Newbury | ad.at nest w/yg. | D.Morimoto# | |
| Rough-legged Hawk: | | | | |
| 1,6 | Saugus | 1 lt. | J.Berry | |
| Golden Eagle: | | | | |
| 3,21 | P.I. | 1 ad. | J.Barton, BBC(A.Blaisdell) | |
| American Kestrel: | | | | |
| 3;10,11 | Newbypt; E.Boston | 7; 6, 1 | BBC(G.d'Entremont); S.Zendeh | |
| Merlin: | | | | |
| 1,10,11,21 | Mt.A. | 1+ | v.o. | |
| 2,6 | P'town, Marshfield | 2, 3 | R.Humphrey#, D.Clapp | |
| 5,6 | W.Bridgewater, Bolton Flats | 1, 1 | K.Ryan, E.Salmela | |
| 8,9;8 | Marshfield (DWWS), N.Scituate | 1, 1; 1 | D.Clapp; M.+B.Litchfield | |
| 17 | P.I., Ipswich | 1, 1 | R.Stymeist#, R.Forster# | |
| Peregrine Falcon: | | | | |
| 2,5 | Truro, W.Bridgewater | 1, 1 | R.Humphrey#, K.Ryan | |
| 6 | Squantum, Eastham | 1, 1 | K.Ryan, L.MacIvor# | |
| 8,30;8 | Boston; P.I. | 1, 1-2; 1 | J.Berry; J.Gruan | |
| 9,11 | S.Dartmouth, N.Monomoy | 1, 1 | A.Hankin, R.Humphrey | |
| 17,21 | N.Scituate, Sharon | 1, 1 | SSBC(K.Anderson), K.Ryan | |

RUFFED GROUSE THROUGH AMERICAN COOT

Two Clapper Rails were found in South Dartmouth, and King Rails were reported from Lynnfield Marsh, Plum Island (Hellcat Swamp), and South Dartmouth. Soras and Virginia Rails were noted in far fewer numbers than in previous years, an observation made by observers in the Lynnfield area. Only two American Coot were reported all month.

R.H.S.

| | | | | |
|--------------------|------------------------|-------------------|------------------------------------|--|
| Ruffed Grouse: | | | | |
| 24 | Uxbridge | 3 | M.Lynch# | |
| Wild Turkey: | | | | |
| 29 | Petersham | 1 ad. m. | E.Nielsen | |
| Northern Bobwhite: | | | | |
| thr. | Belmont, Waltham | 1-2+, 1-2+ | L.Taylor | |
| 4,11 | Halifax, Holliston | 3, 1 | K.Holmes, R.Hildreth | |
| Clapper Rail: | | | | |
| 27 | S.Dartmouth | 2 | R.Marshall | |
| King Rail: | | | | |
| 17,24-25 | Lynnfield, P.I. | 1, 1 | H.Wiggin#, L.Crofoot + v.o. | |
| 26 | S.Dartmouth | 1 | T.Raymond | |
| Virginia Rail: | | | | |
| thr. | W.Harwich, P.I. | max.2, max.3+ | fide B.Nikula, v.o. | |
| thr. | Marshfield, Lynnfield | max.4, max.8 | W.Petersen#+v.o., R.Stymeist#+v.o. | |
| 18,26 | Milton(FM), Marshfield | 1, pr.w/downy yg. | L.Taylor#, W.Petersen | |
| Sora: | | | | |
| thr. | Lynnfield | max.4 (5/13) | J.Berry + v.o. | |
| 10 | Marshfield, Randolph | 2, 1 | W.Petersen#, G.d'Entremont# | |
| 17,28 | Oxbow NWR, P.I. | 1, 1 | BBC(M.Lynch), D.Morimoto# | |
| Common Moorhen: | | | | |
| thr. | P.I. | max.4+ (5/31) | R.Heil + v.o. | |
| 16,17 | Wayland, Milford | 1, 3 | R.Forster#, R.Hildreth | |
| 17,19;17,24 | S.Monomoy; W.Newbury | 1; 1, 2 | R.Humphrey#; D.Morimoto#, R.Heil | |
| 25 | Westboro(Suasco) | 1 | C.Quinlan | |
| American Coot: | | | | |
| 17,29;17 | S.Monomoy; Lynnfield | 1; 1 | B.Nikula#; R.Forster# | |

SHOREBIRDS THROUGH ALCIDS

Lesser Golden-Plovers, not numerous as a spring migrant here, were seen in three locations. Piping Plovers were reported from seven locations that they possibly use as breeding territory. Ten pairs of American Oystercatcher were counted at North Monomoy, and

eight pairs were on South Monomoy. There were also sixteen Willet nests found on North Monomoy. Greater and Lesser yellowlegs arrived in good numbers, and twenty Solitary Sandpipers were noted in the Topsfield area. A Black-necked Stilt was reported from Belle Isle Marsh, East Boston, with details submitted. However, there were no accompanying details with the report of a "Eurasian" Whimbrel on Nantucket. This European subspecies has whitish wing linings and a white V up the rump, similar to a dowitcher. Normally high counts of the following species were reported, mostly at North Monomoy: Ruddy Turnstone, Red Knot, Sanderling, Semipalmated Sandpiper, Least Sandpiper, White-rumped Sandpiper, and Dunlin. The only Pectoral Sandpipers reported were a flock of twenty in Lakeville. A Curlew Sandpiper in breeding plumage was seen for two days at Newburyport Harbor, where a Ruff and a Reeve were seen earlier in the month on separate occasions.

A Pomarine Jaeger was noted at Nantucket, and several Parasitic Jaegers were reported from three locations on Cape Cod. Iceland and Glaucous gulls were present to the month's end. Caspian Terns continued migrating at the beginning of the month with reports from four locations, and a Royal Tern appeared at North Monomoy where an adult Sandwich Tern was also seen. Reported numbers and occurrences of Roseate Terns were very low, perhaps indicating that observers were not present in the areas where this tern species was - and not necessarily indicating a decline of this species. Four Arctic Terns were seen at Plymouth Beach and ten were reported from Monomoy. Black Skimmers were present in the Chatham-Orleans area, possibly as breeders, and two were reported from Martha's Vineyard. A Razorbill was seen in Nantucket Sound on May 23.

G.W.G.

| SPECIES/DATE | LOCATION | NUMBER | OBSERVERS | MAY 1986 |
|---|---------------------------|-------------------|--------------------------------|----------|
| <u>Black-bellied Plover:</u> | | | | |
| thr. | N.Monomoy | max.900 (5/10) | fide B.Nikula | |
| 23,25 | M.V., E.Boston | 100, 770 | BBC(J.Barton), S.Zendeh | |
| 31 | Newburyport | 500 | J.Berry | |
| <u>Lesser Golden-Plover:</u> | | | | |
| 10,16,19;23 | N.Monomoy; M.V. | 1; 1 | B.Nikula#; BBC(J.Barton) | |
| 26 | Newbypt | 1 | G.d'Entremont# | |
| <u>Semipalmated Plover:</u> | | | | |
| 11,17 | E.Boston, Scituate | 6, 150 | S.Zendeh, SSBC(K.Anderson) | |
| 19,21 | N.Monomoy, Newbypt | 50, 30 | B.Nikula, BBC(A.Blaisdell) | |
| <u>Piping Plover:</u> | | | | |
| thr. | Nantucket, S.Dartmouth | max.4, max.6 | M.Litchfield, R.Marshall | |
| 10,20 | N.Monomoy, Duxbury | 9, 11 | R.Humphrey, R.Forster# | |
| One or two others from three locations. | | | | |
| <u>Killdeer:</u> | | | | |
| 3,7 | Newbypt, Halifax | 10, 1 ad.+ 4 yg. | BBC(G.d'Entremont), K.Anderson | |
| 25 | P.I., Plymouth | 15, 6 | M.Lynch#, BBC(R.Timberlake) | |
| <u>American Oystercatcher:</u> | | | | |
| thr. | N.Monomoy, S.Monomoy | 10 pr., 8 pr. | R.Humphrey# | |
| thr. | Orleans | max. 6 | v.o. | |
| 27,31 | S.Dartmouth, M.V. | 2, 4 | R.Marshall, D.Davis# | |
| <u>Black-necked Stilt:</u> | | | | |
| 25 | E.Boston | 1 | C.Foley (fide S.Zendeh) | |
| <u>Greater Yellowlegs:</u> | | | | |
| 3,11 | Newbypt, Squantum | 50, 85 | J.Berry, SSBC(A.Leggett) | |
| 10 | W.Newbury, Newbypt | 70, 430 | BBC(S.Bolton), R.Forster | |
| 10,26 | E.Boston, Newbypt | 30, 35 | S.Zendeh, M.Lynch# | |
| <u>Lesser Yellowlegs:</u> | | | | |
| 10,18 | Newbypt, Squantum | 20, 40 | R.Forster, D.Morimoto# | |
| 21,25 | Newbypt, P.I. | 3, 1 | BBC(A.Blaisdell), M.Lynch# | |
| <u>Solitary Sandpiper:</u> | | | | |
| 4,11 | Bridgewater, Petersham | 1, 2 | K.Holmes, G.Gove# | |
| 17 | ONWR, Middleton-Topsfield | 3,20 | M.Lynch#, J.Berry | |
| 26 | Belmont | 2 | L.Taylor | |
| 6-25 | 7 locations | 1-2 ind. | v.o. | |
| <u>Willet:</u> | | | | |
| thr. | N.Monomoy | 40±(16 nests) | B.Nikula# + L.MacIvor | |
| thr. | Newburyport | max.8 (5/17) | v.o. | |
| <u>Spotted Sandpiper:</u> | | | | |
| 3,11 | Waltham, Petersham | 1, 2 | L.Taylor, G.Gove# | |
| 18,25 | P'town, Wareham | 1, 2 | D.Morimoto#, L.Robinson | |
| <u>Upland Sandpiper:</u> | | | | |
| 3 | Newbypt, Middleboro | 4, 2 | BBC(G.d'Entremont), K.Holmes | |
| 6,17 | Bolton, E.Middleboro | 1, 1 | E.Salmela, K.Anderson | |
| 18,25 | N.Monomoy, Halifax | 1, pr. displaying | R.Humphrey, W.Petersen | |
| <u>Whimbrel:</u> | | | | |
| 29,31 | N.Monomoy | 1 | J.Russell#, W.Petersen# | |

| SPECIES/DATE | LOCATION | NUMBER | OBSERVERS | MAY 1986 |
|-----------------------------------|----------------------|----------------------|---------------------------------|----------|
| "Eurasian" Whimbrel (no details): | | | | |
| 27 | Nantucket | 1 | S.Perkins | |
| Ruddy Turnstone: | | | | |
| thr. | N.Monomoy | max.300 (5/31) | B.Nikula# + v.o. | |
| 23-25,25 | M.V., E.Boston | 24, 15 | BBC(J.Barton), S.Zendeh | |
| Red Knot: | | | | |
| thr. | N.Monomoy | max.150 (5/27) | B.Nikula# + v.o. | |
| Sanderling: | | | | |
| thr. | N.Monomoy | max.2500 (5/31) | B.Nikula# + W.Petersen# | |
| 23,25 | M.V., Nahant | 300, 120 | BBC(J.Barton), R.Stymeist# | |
| Semipalmated Sandpiper: | | | | |
| thr. | N.Monomoy | max.3000 (5/31) | B.Nikula# + W.Petersen# | |
| 21 | Newburyport | 200 | BBC(A.Blaisdell) | |
| Least Sandpiper: | | | | |
| thr. | N.Monomoy | max.150 (5/19) | B.Nikula# | |
| thr. | S.Dartmouth | max.213 (5/13) | R.Marshall | |
| 10,17 | Newbypt, Halifax | 300, 46 | R.Forster, SSBC(K.Anderson) | |
| 25,26 | P.I., Ipswich | 50, 7 | M.Lynch#, J.Berry | |
| White-rumped Sandpiper: | | | | |
| 6,10-31 | Scituate, N.Monomoy | 1, max.50 (5/31) | D.Clapp, B.Nikula# + W.Petersen | |
| 25,31 | Newbypt, P.I. | 10, 20 | G.Gove#, R.Heil | |
| Pectoral Sandpiper: | | | | |
| 16 | Lakeville | 20 | K.Holmes | |
| Purple Sandpiper: | | | | |
| 17,26 | Lynn | 250, 35 | R.Forster | |
| Dunlin: | | | | |
| thr. | N.Monomoy | max.1000 (5/10) | B.Nikula# | |
| 10,23 | Newbypt, M.V. | 200, 50 | R.Forster, BBC(J.Barton) | |
| Curlew Sandpiper: | | | | |
| 25,26 | Newbypt | 1 ad. br.pl. | G.Gove#, C.Floyd# | |
| Stilt Sandpiper: | | | | |
| 6,21 | Eastham, Newbypt | 1, 1 | .MacIvor#, BBC(A.Blaisdell) | |
| 26,31 | P.I. | 1, 1 br.pl. | M.Lynch#, R.Heil | |
| Ruff: | | | | |
| 3,10 | Newburyport | 1 f., 1 m. | J.Berry, R.Forster | |
| Short-billed Dowitcher: | | | | |
| 16-31 | N.Monomoy | max.10 (5/27) | B.Nikula# | |
| 25 | E.Boston, Plymouth | 60, 50 | S.Zendeh, BBC(R.Timberlake) | |
| 31 | Newburyport | 40 | J.Berry | |
| Common Snipe: | | | | |
| 10 | W.Newbury, Whitman | 3, 1 | BBC(S.Bolton), W.Petersen# | |
| 17 | ONWR | 2 | BBC(M.Lynch) | |
| American Woodcock: | | | | |
| 17,21 | P.I., S.Wellfleet | 6, 6 | R.Stymeist#, A.Williams# | |
| Wilson's Phalarope: | | | | |
| 10-17 | P.I. | max. 5 | v.o. | |
| 25,31 | E.Boston, N.Monomoy | 2, 1 m. | S.Zendeh, V.Rosenberg# | |
| Red-necked Phalarope: | | | | |
| 14 | S.Monomoy, N.Monomoy | 25, 11 | B.Nikula# | |
| 15 | 10 mi. E of Chatham | 250 | R.Humphrey# | |
| 12 | Nantucket | 84 | S.Perkins | |
| 25 | Stellwagen Bank | 1 | C.Leahy | |
| Red Phalarope: | | | | |
| 12,30 | Nant., N.Monomoy | 126, 1 m. | S.Perkins, B.Nikula | |
| Phalarope sp.: | | | | |
| 12 | Dennis | 6 | B.Nikula | |
| Pomarine Jaeger: | | | | |
| 17 | Nantucket | 1 | M.Litchfield# | |
| Parasitic Jaeger: | | | | |
| 14,16,27 | N.Monomoy | 4, 1, 1 | R.Humphrey# | |
| 15,29 | Chatham, P'town | 5, 3 | R.Humphrey#, B.Nikula# | |
| Laughing Gull: | | | | |
| 17,23 | N.Monomoy, M.V. | 50, 4 | BBC(J.Barton) | |
| Little Gull: | | | | |
| thr. | Newbypt | max. 2 | v.o. | |
| 17 | S.Monomoy | 1 (1S) | B.Nikula# | |
| Common Black-headed Gull: | | | | |
| 4,10,26 | Newburyport | 2(1S), 1 imm., 1 ad. | W.Petersen#, R.Heil, D.Brown# | |
| 17,27-31 | Nahant, N.Monomoy | 1 ad., 1 ad. | G.Gove#, B.Nikula# | |
| Iceland Gull: | | | | |
| 4,19 | P.I., Nantucket | 3, 5 | W.Petersen#, M.Litchfield | |
| 29 | P'town | 1 | B.Nikula# | |

| <u>SPECIES/DATE</u> | <u>LOCATION</u> | <u>NUMBER</u> | <u>OBSERVERS</u> | <u>MAY 1986</u> |
|---------------------------|-----------------------|-----------------|----------------------------------|-----------------|
| Lesser Black-backed Gull: | | | | |
| 12 | Nantucket | 3 | S.Perkins | |
| Glaucous Gull: | | | | |
| 24 | N.Monomoy | 1 | L.MacIvor# | |
| Black-legged Kittiwake: | | | | |
| 20,29 | Orleans, P'town | 12, 4 | B.Nikula, P.Trimble# | |
| Caspian Tern: | | | | |
| 2,10 | N.Scituate, Dartmouth | 2, 1 | D.Clapp, G.d'Entremont# | |
| 11,13 | Newbypt, P.I. | 2, 1 | J.Grugan, T.Collins# | |
| Royal Tern: | | | | |
| 20 | N.Monomoy | 1 | B.Nickerson, R.Humphrey | |
| <u>Sandwich Tern:</u> | | | | |
| 30 | S.Monomoy | 1 ad. | R.Humphrey# | |
| Roseate Tern: | | | | |
| 23,25 | M.V., Plymouth | 4, 3 | BBC(J.Barton), BBC(R.Timberlake) | |
| Common Tern: | | | | |
| 17,23 | N.Monomoy, M.V. | 2000, 1000 | BBC(J.Barton) | |
| 25 | Plymouth, E.Boston | 500, 4 | BBC(R.Timberlake), S.Zendeh | |
| Arctic Tern: | | | | |
| 25,31 | Plymouth, N.Monomoy | 4, 10 | BBC(R.Timberlake), W.Petersen# | |
| Least Tern: | | | | |
| 6,13,20,27 | S.Dartmouth | 1, 2, 11, 75 | R.Marshall | |
| 14,17 | Scituate, N.Monomoy | 20+, 12 | D.Clapp, BBC(J.Barton) | |
| 24,25 | M.V., Plymouth | 24, 30 | BBC(J.Barton), BBC(R.Timberlake) | |
| 25 | E.Boston(Belle Isle) | 2 | S.Zendeh | |
| Black Tern: | | | | |
| 8,14 | P.I., S.Monomoy | 1, 1 | M.Baird#, B.Nikula# | |
| 24,25 | Newbypt, Edgartown | 2, 1 | R.Heil, P.Hallowell | |
| Black Skimmer: | | | | |
| 16-31,24-31 | Orleans, N.Monomoy | 2, max.5 (5/30) | v.o., R.Humphrey# | |
| 24,30 | M.V., S.Monomoy | 2, 1 | BBC(J.Barton), R.Humphrey# | |
| Razorbill: | | | | |
| 23 | Nantucket Sound | 1 | M.Litchfield | |

CUCKOOS THROUGH WOODPECKERS

Reports of both species of cuckoos were very scant with only fifteen Black-billeds and just four Yellow-billeds noted all month.

Nesting owl reports included the usual Common Barn-Owl at the Nip and Tuck Farm on Martha's Vineyard. In Concord, young Barred Owls hatched on May 12, and Long-eared Owls successfully raised young on Plum Island. Short-eared Owls nested on both North and South Monomoy, and young were flying by May 14.

Nighthawks were not reported in any significant numbers and were conspicuously absent from traditional haunts in some parts of Cambridge and Brookline. A Chuck-will's-widow was heard calling in Brewster, where Whip-poor-wills were also heard.

Hummingbird reports were very few, and this observer, for the first time in twenty-five years missed seeing one in Mt. Auburn Cemetery. Interesting woodpecker reports included six Red-headed, four Red-bellied, and six Pileated. A late sapsucker was found in Newburyport on May 10, the only report in May. R.H.S.

| | | | | |
|-----------------------|-----------------------|-----------------------------|-----------------------------|--|
| Black-billed Cuckoo: | | | | |
| 7,17 | Brookline, Hingham | 1, 1 | H.Wiggin, S.Smith | |
| 18,19 | Mt.A. | 4, 1 | R.Stymeist# + v.o. | |
| 18,24 | Bridgewater, Belmont | 1, 1 | K.Holmes, L.Taylor | |
| 24 | Milton, Uxbridge | 3, 1 | R.Forster#, M.Lynch# | |
| 26 | Acoaxet | 2 | BBC(T.Athearn) | |
| Yellow-billed Cuckoo: | | | | |
| 17,21 | Uxbridge | 1, 1 | R.Hildreth, M.Lynch# | |
| 24,25 | Milton, Plymouth | 1, 1 | R.Forster,BBC(R.Timberlake) | |
| Common Barn-Owl: | | | | |
| 17,24 | Middleboro, W.Tisbury | 1, 2 | K.Anderson#, BBC(J.Barton) | |
| Eastern Screech-Owl: | | | | |
| thr. | Mt.A. | 1 | H.Hoffman + v.o. | |
| Great Horned Owl: | | | | |
| early May | Manchester | 2 ad.+ 2 yg. | G.Hotz# | |
| 8,27 | Concord | 2 ad.+ 1 yg. fledged, 1 pr. | N.Clayton#, J.Berry | |
| Barred Owl: | | | | |
| 12 | Concord | pr.nesting-eggs hatched | B.Rosato | |
| 17,25 | E.Middleboro, Boxford | 2, 1 | K.Anderson, R.Stymeist# | |

| SPECIES/DATE | LOCATION | NUMBER | OBSERVERS | MAY 1986 |
|--|---|--|--|----------|
| Long-eared Owl: thr. | P.I. | pr. w/yg. | D.Abbott | |
| Short-eared Owl: thr. 14 | N.Monomoy S.Monomoy | 2+ nests + 5 eggs (5/3) 6 including 2 fledged yg. | R.Humphrey# + v.o. R.Humphrey# | |
| Northern Saw-whet Owl: 16 | Hamilton | 1 | G.d'Entremont + D.Brown | |
| Common Nighthawk: 17 17,18;18 20,28 28,30 | Bolton Flats, Milford Marshfield; Milton Worcester, Whitman Ipswich, Watertown | 1, 2 1, 1; 13 1, 1 1, 2 | M.Lynch#, R.Hildreth D.Clapp; L.Taylor# S.Carroll#, W.Petersen# J.Berry, R.Stymeist | |
| Chuck-will's-widow: 19,22 | Brewster | 1, 1 | R.Finch, D.,E.,+A.Williams | |
| Whip-poor-will: 7,8 17 17 19 21,22 26,31 | Mt.A., P.I. Clinton, P.I. E.Middleboro Hopkinton, S.Dartmouth S.Wellfleet, Brewster Rowley, Medway | 1, 1 2, 1 1 3, 4 5, 2 2, 1 | M.Hall, J.Grujan M.Lynch#, R.Stymeist# K.Anderson S.Stapleton, LCES(R.Maker) D.,E.,+A.Williams R.Forster, S.Stapleton | |
| Chimney Swift: thr.,7 13 | Mt.A., Ipswich Watertown | max.15 (5/6), 20 40 (flock) | v.o., J.Berry R.Stymeist | |
| Other reports of one to six birds only. | | | | |
| Ruby-throated Hummingbird: 6,11 11;17 25 26,30 | Middleboro, Mt.A. Salem; P.I., Boxford Plymouth, P.I. Westport, Mt.A. | 1, 1 3; 1, 1 1, 6 1, 1 | K.Holmes, S.Koffman C.Blasczak; D.Morimoto# BBC(R.Timberlake), M.Lynch# BBC(T.Athearn), v.o. | |
| Belted Kingfisher: thr. | Mt.A. | pr. nesting | R.Stymeist# | |
| Red-headed Woodpecker: 4,5 24 | Dover, Hyde Park Uxbridge, Millis | 3, 1 1, 1 | P.Hallowell, K.Kilduff M.Lynch#, BBC(B.Cassie) | |
| Red-bellied Woodpecker: thr. 12,19 24 | Randolph Chelmsford, Milton M.V. | 1 1, 1 1 | R.Langley + v.o. S.Musgrave, W.+O.Kerr BBC(J.Barton) | |
| Yellow-bellied Sapsucker: 10 | Newburyport | 1 | BBC(S.Bolton) | |
| Pileated Woodpecker: 3,20 26 | Boxford, Littleton Royalston, Quabbin(G-43) | 1, 1 1, 3 | M.Hall, W.Vales M.Lynch#, M.Hall | |

FLYCATCHERS THROUGH VIREOS

A walk along the Gate 40 road to Dana Center at Quabbin on May 11 produced the remarkable count of thirty-nine calling Least Flycatchers. In Harwich on the previous day (and only then), a Scissor-tailed Flycatcher was seen. Interesting corvid reports from the southeastern part of the state included a Blue Jay on South Monomoy. This species is decidedly uncommon at that location. The Nantucket jackdaw continued and again this spring was looking rather poorly as a result of molt. Last year it was a crooked bill; this year a limp! Worth noting also was the successful nesting of a pair of Carolina Wrens in Winchester (in a newly-opened bag of peat moss) with the fledging of three young on May 23 - a first breeding record for Middlesex County of a species rarely, if ever, recorded nesting this far north in the state. Most of the more-watched-for migrant species were observed in typical numbers compared to the May records for the last ten years. Included in this category were Blue-gray Gnatcatcher at forty-five individuals total, Gray-cheeked Thrush at eleven, White-eyed Vireo at thirteen, and Philadelphia Vireo at four.

L.E.T.

| | | | |
|---|---|-----------------------|--|
| Olive-sided Flycatcher: 18,25 25,30 | Dover, Marblehead Newbypt, Annisquam | 1, 1 1, 1 | P.Hallowell, G.Gove# v.o., H.Wiggin |
| Eastern Wood-Pewee: 13,15-31 25,31 | Lynnfield, Mt.A. Boxford, Petersham | 1, 1-2 daily 12, 6 | BBC(W.VanCor), v.o. R.Stymeist#, S.Carroll# |
| Yellow-bellied Flycatcher: 23-30,25 25,31 | Mt.A., MNWS P.I., Annisquam | 3+, 1 1, 1 | v.o., R.Stymeist# M.Lynch#, H.Wiggin |

| <u>SPECIES/DATE</u> | <u>LOCATION</u> | <u>NUMBER</u> | <u>OBSERVERS</u> | <u>MAY 1986</u> |
|--------------------------------|---------------------------------|-------------------|-------------------------------|-----------------|
| Acadian Flycatcher: | | | | |
| 24,26 | MNWS, Royalston Falls | 1, 2 | R.Heil, S.Carroll# | |
| 29,31 | Mt.A. | 2, 2 | E.Nielsen#, v.o. | |
| 31 | Nahant, Hamilton | 1, 1 | R.Stymeist, J.Berry | |
| Alder Flycatcher: | | | | |
| 17,18-30 | ONWR, 6 loc. | 1, 6 ind. | BBC(M.Lynch), v.o. | |
| 25-29 | Lynnfield | 1 | v.o. | |
| Willow Flycatcher: | | | | |
| 6,17 | Cambridge, Lancaster | 1, 3 | J.Barton, S.Carroll# | |
| 17-31 | Lynnfield | max. 4 (5/25) | v.o. | |
| Least Flycatcher: | | | | |
| 6,7 | Boxford, Brookline | 2, 1 | W.Drummond#, H.Wiggin | |
| 11,14-31 | Quabbin(G-40), 9 loc. | 39, 18 total | L.Taylor#, v.o. | |
| 18,31 | Provincetown, Mt.A. | 4, 2 | B.Nikula, v.o. | |
| Eastern Phoebe: | | | | |
| thr.,24 | 6 loc., Uxbridge | 14 total, 8 | v.o., M.Lynch# | |
| Great Crested Flycatcher: | | | | |
| thr.,15 | Mt.A., Manchester | max. 6 (5/10), 6 | v.o., BBC(G.Hotz) | |
| 17,23 | Topsfield, M.V. | 20, 6 | J.Berry, BBC(J.Barton) | |
| Eastern Kingbird: | | | | |
| 1,4 | Bridgewater, Marshfield | 1, 1 | K.Holmes, D.Clapp | |
| 10-31,17 | Mt.A., ONWR | max. 6 (5/21), 12 | v.o., BBC(M.Lynch) | |
| 21,24 | Newbypt, Uxbridge | 15, 10 | BBC(A.Blaisdell), S.Carroll# | |
| Scissor-tailed Flycatcher: | | | | |
| 10 | Harwich | 1 | P.Gove | |
| Horned Lark: | | | | |
| 17 | P.I., N.Monomoy | 1, 20 | G.d'Entremont, BBC(J.Barton) | |
| 23,26 | M.V., Gardner | 2, 2 | BBC(J.Barton), M.Lynch# | |
| Purple Martin: | | | | |
| thr.,29 | P.I., P'town | max. 40 (5/17), 1 | v.o., P.Trimble# | |
| Northern Rough-winged Swallow: | | | | |
| 1,6 | Fall River, Peabody | 4, 11 | K.Holmes, R.Heil | |
| 24,25 | Uxbridge, Plymouth | 9, 5 | S.Carroll#, BBC(R.Timberlake) | |
| Bank Swallow: | | | | |
| 17 | ONWR, Lakeville | 2, 15 | BBC(M.Lynch), K.Holmes | |
| 18,23 | Manomet, M.V. | 50, 12 | D.Morimoto#, BBC(J.Barton) | |
| Cliff Swallow: | | | | |
| 3,7 | Topsfield, Concord | 1, 1 | H.Wiggin#, T.Carrolan | |
| 17-19,25 | Rowley, P'town | max. 18, 1 | v.o., L.MacIvor# | |
| Blue Jay: | | | | |
| 29 | S.Monomoy | 1 | J.Russell# | |
| Eurasian Jackdaw: | | | | |
| thr. | Nantucket | 1 | E.Andrews# | |
| American Crow: | | | | |
| 11 | Gloucester | 55 | BBC(J.Nove) | |
| Fish Crow: | | | | |
| thr. | Mt.A., E.Middleboro | 2 pr., 2 | v.o., K.Anderson | |
| 2-18 | 8 loc. | 8 ind. | v.o. | |
| Common Raven: | | | | |
| 11 | Quabbin(G-40) | 2 | G.Gove# | |
| Red-breasted Nuthatch: | | | | |
| 1-19,thr. | Mt.A., E.Middleboro | max. 3 (5/7), pr. | v.o., K.Anderson | |
| 10-26,23 | Boxford, M.V. | max. 4 (5/10), 2 | v.o., BBC(J.Barton) | |
| 31 | Plymouth, Hamilton | 1 m., 2 | G.d'Entremont, J.Berry | |
| Brown Creeper: | | | | |
| 1,6 | Mt.A., Bridgewater | 1, 2 | v.o., K.Holmes | |
| 10 | Boxford, Wareham | 3, pr. nesting | J.Berry, L.Robinson | |
| Carolina Wren: | | | | |
| thr. | S.Dartmouth | max. 5 (5/26) | R.Maker | |
| 17;24,25 | Nant.; Whitman, Scituate | 1; 1, 1 | M.Litchfield; W.Petersen | |
| 3-23 | Winchester nesting pr. w/ 3 yg. | fledged | L.Abers-Nunno + T.Nunno | |
| House Wren: | | | | |
| 1,1 | Lakeville, Mt.A. | 1, 1 | K.Holmes, v.o. | |
| thr., 3 | S.Dartmouth, Waltham | max. 6 (5/10), 2 | LCES, L.Taylor | |
| Marsh Wren: | | | | |
| 6,10 | S.Dartmouth, Marshfield | 1, 3 | R.Maker, W.Petersen# | |
| 13-31,25 | Lynnfield, P.I. | max. 15 (5/17), 8 | v.o., S.Carroll# | |
| Golden-crowned Kinglet: | | | | |
| 23 | M.V. | 4 | BBC(J.Barton) | |
| Ruby-crowned Kinglet: | | | | |
| 1-13,3 | Mt.A., Manchester | max. 6 (5/6), 3 | v.o., BBC(G.Hotz) | |
| 10,11 | Boxford, Gloucester | 4, 4 | J.Berry, BBC(J.Nove) | |

| <u>SPECIES/DATE</u> | <u>LOCATION</u> | <u>NUMBER</u> | <u>OBSERVERS</u> | <u>MAY 1986</u> |
|------------------------|------------------------|------------------------|-------------------------------|-----------------|
| Blue-gray Gnatcatcher: | | | | |
| thr. | P'town, Boxford | max. 3, 3 pr. | B.Nikula#, v.o. | |
| thr., 1-10 | Petersham, Mt.A. | max. 6, 1-2 | v.o. | |
| 1-20, 4-17 | 10 loc., Waltham | 18 total, max. 2 | v.o., L.Taylor | |
| 17, 31 | Topsfield, Lincoln | 7, 1 | J.Berry, M.Baird# | |
| Eastern Bluebird: | | | | |
| thr., 2 | E.Middleboro, Millis | pr. nesting, pr. | K.Anderson, B.Cassie | |
| 3, 4-31 | Milton, Plymouth | pr., max. 8 (5/14) | R.Stymeist#, v.o. | |
| 11, 11-18 | Quabbin(G-40), Milford | 3 pr., 2 | L.Taylor#, R.Hildreth | |
| 18, 23 | E.Dartmouth, M.V. | 1 m., 3 | J.Swenson, BBC(J.Barton) | |
| 25 | Lincoln | 3 | P.Swift | |
| Veery: | | | | |
| 1-27, 3 | Mt.A., Milton | max. 4 (5/10), 1 | v.o., R.Stymeist# | |
| 18, 24 | MNWS, Uxbridge | 6, 7 | C.Blasczak, M.Lynch# | |
| Gray-cheeked Thrush: | | | | |
| 17 | Newbypt, Milford | 1, 1 | BBC(W.Drummond), R.Hildreth | |
| 17, 18-30 | Nahant, Mt.A. | 2, 3 total | H.Coolidge#, v.o. | |
| 18-25 | 4 loc. | 4 total | v.o. | |
| Swainson's Thrush: | | | | |
| 13, 15 | Nantucket, Manchester | 1, 1 | E.Andrews, BBC(G.Hotz) | |
| 17 | Nahant, MNWS | 20, 15 | R.Stymeist# | |
| 18-30, 23 | Mt.A., M.V. | max. 10 (5/19), 4 | v.o., BBC(J.Barton) | |
| Hermit Thrush: | | | | |
| 1-11, 17 | Mt.A., Boxford | 16 total, 1 | v.o. | |
| 18 | Provincetown | 1 | D.Morimoto# | |
| Wood Thrush: | | | | |
| 3-24, 6 | Waltham, Boxford | max. 3 (5/24), 2 | L.Taylor, W.Drummond# | |
| 6-23, 23 | Mt.A., M.V. | max. 5 (5/18), 12 | v.o., BBC(J.Barton) | |
| Gray Catbird: | | | | |
| 3, 4-31 | Belmont, Mt.A. | 1, max. 13 (5/25) | L.Taylor, v.o. | |
| 23, 25 | M.V., P.I. | 50, 20 | BBC(J.Barton), S.Carroll# | |
| Northern Mockingbird: | | | | |
| thr. | Mt.A. | 5 pr. | v.o. | |
| Brown Thrasher: | | | | |
| 2, 4 | Waltham, Marshfield | 2, 1 | L.Taylor, D.Clapp | |
| 4, 11 | Wareham, Ipswich | 1, 5 | L.Robinson, J.Berry | |
| 25 | P.I. | 6 | R.Stymeist# | |
| Water Pipit: | | | | |
| 10, 17 | Mt.A., P.I. | 1, 2 | R.Stymeist, S.Surner | |
| Cedar Waxwing: | | | | |
| thr., 1 | Mt.A., E.Middleboro | max. 150 (5/11), 13 | D.Arvidson+v.o., K.Anderson | |
| 3, 17 | Newbypt, Belmont | 40, 15 | BBC(G.d'Entremont), L.Taylor# | |
| 30, 31 | Annisquam, Ipswich | 8, 7 | H.Wiggin, J.Berry | |
| White-eyed Vireo: | | | | |
| 1, 6, 9-13 | Mt.A. | 1, 1, 1-2 | v.o. | |
| 9-18, 17-31 | 7 loc., Nahant | 7 total, max. 2 (5/31) | v.o. | |
| Solitary Vireo: | | | | |
| 1-18, 6 | Mt.A., Peabody | max. 4, 18 | v.o., R.Heil | |
| 10, 25 | Waltham, P.I. | 6, 1 | L.Taylor, M.Lynch# | |
| 26 | Royalston | 2 | S.Carroll# | |
| Yellow-throated Vireo: | | | | |
| 6, 17 | Peabody, Westport | 1, 1 | R.Heil, K.Holmes | |
| 17, 18 | ONWR, Mt.A. | 4, 1 | BBC(M.Lynch), S.Carroll# | |
| 22-31, 24 | Topsfield, Uxbridge | pr.+ nest, 1 | v.o., M.Lynch# | |
| Warbling Vireo: | | | | |
| thr. | Mt.A., Cambridge(F.P.) | 5 total, max. 12(5/20) | v.o., J.Barton | |
| 17, 18-19 | ONWR, Nantucket | 12, 1 | BBC(M.Lynch), E.Andrews | |
| 19, 25 | Wayland, Lynnfield | 10, 6 pr. | L.Robinson, R.Stymeist# | |
| Philadelphia Vireo: | | | | |
| 17, 18 | Fall River, P'town | 1, 2 | K.Holmes#, D.Morimoto# | |
| 23-24 | M.V. | 1 | BBC(J.Barton) | |
| Red-eyed Vireo: | | | | |
| 11-31, 17 | Mt.A., ONWR | max. 15 (5/18), 8 | v.o., S.Carroll# | |
| 17, 23 | P.I., M.V. | 8, 12 | D.Morimoto#, BBC(J.Barton) | |
| 24, 30 | Uxbridge, Annisquam | 15, 5 | M.Lynch#, H.Wiggin | |

WARBLERS

The total of thirty-six warbler species, including reports of both "winged" warbler hybrids, was typical for May. Of the thirty-six, twenty-nine species made their first appearances for the year in the month of May, an unusually high number caused by the lack of proper migrating weather in late April. The two biggest days for warbler first

arrivals were May 1 and May 6, with four species each; weather was a definite factor on the sixth, with warm, southwesterly flow prevailing the night before. Days showing first arrivals for three warbler species each were May 4, 10, 11, and 14, consisting mostly of weekends. The latest first arrival for a species which might regularly be expected was May 17 for Mourning Warbler, which staged a generally unremarkable flight this year. Nashville Warblers, both in total count for the month and in the highest single-day count, were reported in comparatively good numbers. There was also a spectacular single-day count of over one hundred Black-and-white Warblers in Peabody on May 6. The expected "southern" warbler species were all represented this May, and in typical numbers. This is the fourth year in a row for the male Kentucky Warbler at Fowl Meadow in Milton.

L.E.T.

| SPECIES/DATE | LOCATION | NUMBER | OBSERVERS | MAY 1986 |
|-------------------------------|---------------------------|------------------------------|--------------------------------|----------|
| Blue-winged Warbler: | | | | |
| 6 | Boxford, Lexington | 2, 1 | W.Drummond#, M.Baird | |
| 6-15,15 | Mt.A., Scituate | max. 4 (5/11), 3 | v.o., M.Litchfield# | |
| 17,17-31 | ONWR, W.Newbury | 3, max. 3 | BBC(M.Lynch), v.o. | |
| 24 | Uxbridge | 5 | S.Carroll# | |
| Golden-winged Warbler: | | | | |
| 14,21 | Wayland, Lincoln | 1, 1 | J.Huntington, M.Baird | |
| 21,22-31 | W.Newbury, Topsfield | 1, 1 | BBC(A.Blaisdell), v.o. | |
| 26-31 | Ipswich | max. 3 (5/26) | G.d'Entremont+D.Brown+v.o. | |
| "Brewster's" Warbler: | | | | |
| 25 | Ipswich | 2 | M.Lynch# | |
| "Lawrence's" Warbler: | | | | |
| 6,17-31 | WBWS, W.Newbury | 1 ph. b., 1 | D.Reynolds#, v.o. | |
| Tennessee Warbler: | | | | |
| 11-30,17 | Mt.A., P.I. | max. 20 (5/18), 6 | v.o., BBC(W.Drummond) | |
| Orange-crowned Warbler: | | | | |
| 7,17 | Brookline, MNWS | 2, 1 | B.Hallett, C.Floyd# | |
| Nashville Warbler: | | | | |
| 1-23,3 | Mt.A., Newbypt | max.20 (5/7), 2 | v.o., BBC(G.d'Entremont) | |
| 6,10 | Peabody, Middleboro | 17, 4 | R.Heil, K.Holmes | |
| 17,26 | P.I., Royalston | 4, 3 | BBC(W.Drummond), S.Carroll# | |
| Northern Parula: | | | | |
| 1-30,6 | Mt.A., Peabody | max.20 (5/7), 40 | v.o., R.Heil | |
| 17,25 | P.I., Middleboro | 12, 1 m. | S.Surner, W.Petersen# | |
| 30 | Annisquam | 2 m. | H.Wiggin | |
| Yellow Warbler: | | | | |
| 1,thr. | Wayland, Cambridge (F.P.) | 2, max.21 (5/15) | G.Gove, J.Barton | |
| 6-31,11 | Mt.A., Gloucester | max.7 (5/10), 26 | v.o., BBC(J.Nove) | |
| 17 | ONWR, Topsfield | 20, 42 | BBC(M.Lynch), J.Berry | |
| Chestnut-sided Warbler: | | | | |
| 6-18,17 | Mt.A., ONWR | max.5 (5/11), 8 | v.o., BBC(M.Lynch) | |
| 26,27 | Royalston, Mt.A. | 9, 1 | S.Carroll#, R.Stymeist# | |
| Magnolia Warbler: | | | | |
| 4,6-27 | Holliston, Mt.A. | 1, max.10 (5/18) | R.Hildreth, v.o. | |
| 17,31 | P.I., Marshfield | 12, 1 | BBC(W.Drummond), D.Morimoto | |
| Cape May Warbler: | | | | |
| 6-13,17 | Mt.A., Newbypt | max.6 (5/16), 1 | v.o., BBC(W.Drummond) | |
| Black-throated Blue Warbler: | | | | |
| 6-29,17 | Mt.A., P.I. | max.3, 3 | v.o., D.Morimoto# | |
| 18 | Nantucket | 2 | J.Andrews# | |
| Yellow-rumped Warbler: | | | | |
| 1-25,6 | Mt.A., Peabody | max.100 (5/6), 200 | v.o.; R.Heil | |
| 6,26 | P'town, Wareham | 200, 1 m. | B.Nikula#, L.Robinson | |
| Black-throated Green Warbler: | | | | |
| 2-27,4-17 | Mt.A., Boxford | max.12 (5/10), max.11 (5/10) | v.o. | |
| 7,30 | Ipswich, Annisquam | 10, 1 m. | J.Berry, H.Wiggin | |
| Blackburnian Warbler: | | | | |
| 3-17,7-27 | Boxford, Mt.A. | max.2, max.9 (5/18) | v.o. | |
| 17 | Waltham, P.I. | 6, 6 | L.Taylor, BBC(W.Drummond) | |
| 18,30 | Marshfield, Annisquam | 5, 2 | BBC(R.Timberlake), H.Wiggin | |
| Yellow-throated Warbler: | | | | |
| 2 | Norfolk | 1 m. | B.Cassie# | |
| Pine Warbler: | | | | |
| thr.,7 | E.Middleboro, Ipswich | 2 m., 6 m. | K.Anderson, J.Berry | |
| 17,23 | ONWR, M.V. | 2, 6 | BBC(M.Lynch), BBC(J.Barton) | |
| 25 | Plymouth, Boxford | 3, 3 pr. | BBC(R.Timberlake), R.Stymeist# | |
| Prairie Warbler: | | | | |
| 4-18,11 | Holliston, Mt.A. | max.17 (5/18), 2 | R.Hildreth, v.o. | |
| 17,23 | W.Newbury, M.V. | 2, 12 | D.Morimoto#, BBC(J.Barton) | |

| SPECIES/DATE | LOCATION | NUMBER | OBSERVERS | MAY 1986 |
|--------------------------|----------------------------|--------------------|--------------------------------|----------|
| Palm Warbler: | | | | |
| 1-11, 3 | Mt.A., Boxford | max.6 (5/4), 2 | v.o., BBC(G.d'Entremont) | |
| 3, 10 | Manchester, Newton | 4, 2 | BBC(G.Hotz), BBC(D.+A.Bandes) | |
| Bay-breasted Warbler: | | | | |
| 11-23, 17 | Mt.A., P.I. | max.10 (5/18), 6 | v.o., BBC(W.Drummond) | |
| 17, 18 | MNWS, Newbypt | 6, 8 | R.Forster | |
| 18 | Cambridge(F.P.), P'town | 18, 4 | J.Barton, D.Morimoto# | |
| Blackpoll Warbler: | | | | |
| 11-31, 23 | Mt.A., M.V. | max.40 (5/20), 6 | v.o., BBC(J.Barton) | |
| 24 | Marshfield, MNWS | 20, 40 | D.Morimoto#, R.Heil | |
| 29, 30 | Nant., Annisquam | 1, 3 | E.Andrews, H.Wiggin | |
| Cerulean Warbler: | | | | |
| 4, 14 | Cambridge(F.P.), Worcester | 1 m., 1 m. | J.Barton, S.Carroll# | |
| 17, 25 | Mashpee, Newbypt | 1, 1 | B.Nikula, B.Drummond# | |
| Black-and-white Warbler: | | | | |
| 1-25, 6 | Mt.A., Peabody | max.27 (5/6), 100+ | v.o., R.Heil | |
| 17, 23 | P.I., M.V. | 16, 6 | S.Surner, BBC(J.Barton) | |
| American Redstart: | | | | |
| 10, 11-31 | Bridgewater, Mt.A. | 1, max.20 (5/21) | K.Holmes, v.o. | |
| 14-31 | Cambridge(F.P.) | max.17 (5/18) | J.Barton | |
| 18 | Marshfield | 16 | BBC(R.Timberlake) | |
| Prothonotary Warbler: | | | | |
| 10, 12 | WBWS, Nantucket | 1, 1 | S.Clifton#, S.Perkins | |
| Worm-eating Warbler: | | | | |
| 7; 10, 19 | Brookline; MNWS | 1; 1, 1 | H.Wiggin; R.Heil | |
| 10, 17 | Westport, ONWR | 1, 1 | D.Brown#, M.Lynch# | |
| 18, 18-31 | Bridgewater, Dover | 1, 1 | K.Holmes, P. Hallowell | |
| 19, 22 | Mt.A. | 1, 2 | v.o. | |
| Ovenbird: | | | | |
| 3, 6-20 | Manchester, Mt.A. | 1, max.6 (5/11) | BBC(G.Hotz), v.o. | |
| 10, 11 | Boxford, Wareham | 20 m., 25 | J.Berry, L.Robinson | |
| 18, 24 | MNWS, Uxbridge | 11, 14 | C.Błaszcak, S.Carroll# | |
| Northern Waterthrush: | | | | |
| thr. | Mt.A., Brookline | 1-2, pr. | v.o., R.Stymeist# | |
| 17, 18 | P.I., P'town | 3, 3 | D.Morimoto# | |
| Louisiana Waterthrush: | | | | |
| thr., 10 | Boxford, Bridgewater | 3 pr., 1 | v.o., K.Holmes | |
| 21, 31 | Uxbridge, Petersham | 2, 3 | S.Carroll# | |
| Kentucky Warbler: | | | | |
| 8, 11 | Millis, Salem | 1 m., 1 | B.Cassie, C.Błaszcak | |
| 11, 13-31 | Whitman, Milton(F.M.) | 1, 1 m. | K.Holmes, v.o. | |
| 17, 22 | Boston, MNWS | 1 m., 1 | P.Thayer, R.Heil | |
| 23, 28; 25 | Mt.A.; Milford | 1, 1; 1 | v.o.; C.Quinlan | |
| Mourning Warbler: | | | | |
| 17, 17-31 | FallRiver, Nahant | 1, max.4 (5/20) | K.Holmes#, v.o. | |
| 18-31, 23-30 | 9 loc., Mt.A. | 9 ind., 4 total | v.o. | |
| 24, 30 | MNWS, Annisquam | 3, 2 m. | R.Heil, H.Wiggin | |
| Common Yellowthroat: | | | | |
| thr., 3 | Mt.A., Bridgewater | max.5 (5/18), 1 | v.o., K.Holmes | |
| 17, 18 | Topsfield, MNWS | 52, 27 | J.Berry, C.Błaszcak | |
| Hooded Warbler: | | | | |
| 14-20, 19 | Mt.A., MNWS | 1, 1 f. | v.o., R.Heil | |
| 20, 24 | Chatham, Newbury | 1, 1 m. | R.Clem, R.Heil | |
| Wilson's Warbler: | | | | |
| 14, 15 | Mt.A., Manchester | 2, 2 | v.o., BBC(G.Hotz) | |
| 18 | Marshfield, P'town | 3, 2 | BBC(R.Timberlake), D.Morimoto# | |
| 28, 29 | P.I., Mt.A. | 2, 2 | D.Morimoto, v.o. | |
| Canada Warbler: | | | | |
| 10, 15 | Halifax, Manchester | 1, 1 | K.Holmes, BBC(G.Hotz) | |
| 16-30, 18 | Mt.A., Marshfield | max.12 (5/18), 5 | v.o., BBC(R.Timberlake) | |
| 27, 30 | Cambridge(F.P.), Annisquam | 6, 2 | J.Barton, H.Wiggin | |
| Yellow-breasted Chat: | | | | |
| 23 | M.V. | 1 | BBC(J.Barton) | |

SUMMER TANAGER THROUGH EVENING GROSBEAK

A total of six Summer Tanagers on Nantucket was excellent for a single locale, though the regional total of thirteen individuals for the month of May was about average. Most of the Nantucket birds were at feeders and were verified as being different individuals. Blue Grosbeak numbers remained on the low side, as in April. Uncommon sparrow species reported included a Clay-colored at Peabody on May 5 and two male Lark Buntings, one in Sandwich, May 16-21, and one at Plum Island on May 20. The White-crowned

Sparrow flight was the best in ten years - a total of over eighty individuals, many at feeders, at widely dispersed locations. Dark-eyed Juncos departed early and thoroughly this year with only a single individual reported during the first week of the month. Contrarily, a lone Snow Bunting tarried over a month beyond departure norms on Monomoy. There are recent precedents for late spring reports of this species on Monomoy.

Maximum daily counts of Bobolinks were low compared to recent years. The report of 190 Rusty Blackbirds at Millis was very high for so late in the season. Several winter finch species lingered well into May. It was unusual to record both crossbill species in May, with Reds apparently breeding successfully on Cape Cod. The Middleboro White-winged Crossbills were observed in three separate flocks, and tape recordings were made. Origins are unknown for the European Goldfinch observed for two days at a feeder in Sudbury.

L.E.T.

| <u>SPECIES/DATE</u> | <u>LOCATION</u> | <u>NUMBER</u> | <u>OBSERVERS</u> | <u>MAY 1986</u> |
|-------------------------|----------------------------------|---------------------|---------------------------------|-----------------|
| Summer Tanager: | | | | |
| 1-17,7 | Nantucket, Brewster | 5 m. + 1 f., 1 | E.Andrews#, P.Trull | |
| 11,14 | Dennis, Newbypt | 1, 1 f. | fide W.Bailey, J.Grujan | |
| 16,17 | Mt.A., P.I. | 1 m., 1 m. | v.o., BBC(W.Drummond) | |
| .19,25 | Chatham, Newbypt | 1, 1 m. | R.Clem, T.Martin | |
| Scarlet Tanager: | | | | |
| 10,11 | Newton, Nantucket | 2, 1 m. | BBC(D.+A.Bandes), F.Reed# | |
| 12-15,18-31 | Bridgewater, Mt.A. | max.2, max.4 (5/22) | K.Holmes#, v.o. | |
| Rose-breasted Grosbeak: | | | | |
| 6-31,8 | Mt.A., Millis | max. 6, 14 | v.o., B.Cassie | |
| 17 | Middleton, Waltham | 15, 30 | J.Berry, L.Taylor | |
| Blue Grosbeak: | | | | |
| 10,17 | Mt.A., Nantucket | 1, 1 m. | J.Barton, R.Tate# | |
| 21-22 | Mt.A. | 1 f. | R.Stymeist# | |
| Indigo Bunting: | | | | |
| 4,5-6 | Dedham, Marion | 1, 1 | Mrs.Hobbs, A.Hegyi | |
| 5-7,18-31 | Norwood, Mt.A. | 2, max.3 (5/18) | C.Polito, v.o. | |
| Rufous-sided Towhee: | | | | |
| thr.,1 | Mt.A., E.Middleboro | max.5 (5/10), 1 | v.o., K.Anderson | |
| 10,23-25 | W.Newbury, M.V. | 20, 100 | BBC(S.Bolton),BBC(J.Barton) | |
| Chipping Sparrow: | | | | |
| thr. | Mt.A. | max.18 (5/10) | v.o. | |
| Clay-colored Sparrow: | | | | |
| 5 | Peabody | 1 | R.Heil | |
| Vesper Sparrow: | | | | |
| 4,26 | Plymouth, Gardner | 6, 1 | R.Abrams, S.Carroll# | |
| Lark Sparrow: | | | | |
| 9 | Chatham | 1 m. | P.Trull | |
| Lark Bunting (details): | | | | |
| 16-21+, 20 | Sandwich, P.I. | 1 ad. m., 1 ad. m. | R.Pease + v.o., H.Parker | |
| Savannah Sparrow: | | | | |
| 1,6 | Mt.A., Peabody | 5, 112 | R.Stymeist, R.Heil | |
| Grasshopper Sparrow: | | | | |
| 13-20,19 | S.Dartmouth, Nant. | max. 3, 1 | R.Marshall, L.Jodrey# | |
| 25 | M.V. | 2 | R.Culbert# | |
| Sharp-tailed Sparrow: | | | | |
| 6,27;23 | S.Dartmouth; M.V. | 2, 34; 4 | R.Marshall; BBC(J.Barton) | |
| 25,26 | E.Boston, P.I. | 3, 3 | S.Zendeh, M.Lynch# | |
| Seaside Sparrow: | | | | |
| 6-31,26 | S.Dartmouth, Westport | max. 5, 1 | R.Marshall, BBC(T.Athearn) | |
| Lincoln's Sparrow: | | | | |
| 6,10 | Brookline, Mt.A. | 1, 1 | D.Arvidson#, v.o. | |
| 17-19,24 | 3 loc., Belmont | 4 total, 1 | v.o., L.Taylor | |
| Swamp Sparrow: | | | | |
| 13,17 | Lynnfield, ONWR | 20+ m., 18 | J.Berry, BBC(M.Lynch) | |
| White-throated Sparrow: | | | | |
| 1-14,6 | Mt.A., Peabody | max. 150 (5/8), 810 | v.o., R.Heil | |
| 16,18 | Boston, Quincy | 2, 2 | J.Berry, D.Morimoto# | |
| White-crowned Sparrow: | | | | |
| 1(from April),3-13 | Ipswich, Mt.A. | 1, max.4 (5/10) | J.Berry, v.o. | |
| 3-15 | 26 loc. (1-5 each) | totaling 58 | v.o. | |
| 6,11 | Peabody, Wayland | 6, 9 | R.Heil, R.Forster | |
| Dark-eyed Junco: | | | | |
| 7 | Holliston | 1 | R.Hildreth | |
| Snow Bunting: | | | | |
| <u>10</u> | N.Monomoy | 1 | R.Humphrey | |
| Bobolink: | | | | |
| 7,15,17 | Easton, Cambridge(F.P.), Harwich | 6, 36, 40 | K.Ryan, J.Barton, B.Nikula | |
| 17,18 | ONWR, Marshfield | 30, 40 | BBC(M.Lynch), BBC(R.Timberlake) | |

| <u>SPECIES/DATE</u> | <u>LOCATION</u> | <u>NUMBER</u> | <u>OBSERVERS</u> | <u>MAY 1986</u> |
|--------------------------|-------------------------|---------------------|--------------------------|-----------------|
| Eastern Meadowlark: | | | | |
| 2,3 | Waltham, Newbypt | 1, 5 | L.Taylor, J.Berry | |
| Yellow-headed Blackbird: | | | | |
| 15 | Marblehead | 1 ad. m. singing | R.Heil | |
| Rusty Blackbird: | | | | |
| 4,6 | Wayland, Millis | 5, 190 | R.Forster, B.Cassie# | |
| Orchard Oriole: | | | | |
| 1;3,18 | Millis; Mt.A. | 1 m.; 1, 1 | B.Cassie; v.o. | |
| 6-31,14 | 19 loc., Peabody | 36 total, 7 | v.o., R.Heil | |
| Northern Oriole: | | | | |
| 2,6-31 | S.Middleboro; Mt.A. | 1 m., max.20 (5/13) | K.Anderson, v.o. | |
| 10,17 | Nant., Topsfield | 15 b., 50 m. | E.Andrews, J.Berry | |
| Purple Finch: | | | | |
| thr.;3,9 | S.Middleboro, Mt.A. | 1 pr.; 1, 1 | K.Anderson; v.o. | |
| 4,10 | Waltham, Boxford | 3, 5 | L.Taylor, J.Berry | |
| Red Crossbill: | | | | |
| 28 | Harwich | 1 pr. + 2 yg. | K.Hamilton | |
| White-winged Crossbill: | | | | |
| 7-12,19 | Watertown, E.Middleboro | 1 f., 27 | F.Epstein, P.Donahue | |
| Pine Siskin: | | | | |
| thr. | P'town, Hopkinton | max.4 (5/11), 1 | B.Nikula, J.Gordon | |
| 6,19-23 | Boxford, Mt.A. | 10, max.6 (5/19) | B.Drummond#, R.Stymeist# | |
| European Goldfinch: | | | | |
| 9-10 | Sudbury | 1 at feeder | E.Bennett | |
| Evening Grosbeak: | | | | |
| 1-16,4 | Mt.A., Wayland | 61 total, 35 | v.o., R.Forster | |
| 7,13 | Townsend, Billerica | 40, 20 | Mrs. Edfors, K.Griffis | |

UPDATE ON THE COMMON LOON (*Gavia immer*)

The Loon Preservation Committee (LPC), a self-funded project of the Audubon Society of New Hampshire, published in June of 1986 a summary of its findings over the past decade. In 1976, the LPC's first survey of 250 New Hampshire lakes and ponds produced the grim result that only 214 adults and 51 chicks survived that year and that chicks had hatched on less than two dozen of the bodies of water surveyed. The current (1986) report reveals that the total New Hampshire Common Loon population in 1985 rose to 435. "Nesting success is increasing along with the number of nesting pairs and the number of lakes on which they nest. The number of fledgling chicks peaked in 1983 at 83 (compared to 36 in 1978) and has remained well above the range of the first five years." However, the LPC workers had discovered by 1977 that "at least half of the lakes which formerly sustained nesting loons no longer did. In 1979 this evidence helped justify...the Common Loon's classification as a threatened species."

The 1986 report states that egg predation by raccoons is the chief cause of nest loss but that egg predation by gulls also has increased recently. Both raccoon and gull populations benefit from human refuse, and hence, man is responsible for the increase in these predators as well as for the other major threats to the loon - habitat loss and direct disturbance. Another factor in the decline of loons cited in the report is the fluctuating water level in lakes and ponds - an increase of six inches or a decrease of one foot will cause nest loss to the shore-nesting loons. The LPC confronted this problem by regulating water level where possible and by providing anchored artificial islands of cedar logs notched together and floored with wire mesh supporting sodded shoreline vegetation and duff to provide nesting material. As many as one-fifth to one-third of the fledged loons have come from raft nests. However, these are useful only where shoreline nesters have tried and failed; they do not seem to attract new nesters.

Other discoveries of the LPC studies are that loon eggs failed despite the absence of DDT and the presence of organochlorine residues at levels believed too low to affect reproduction. However, it seems certain that two heavy metals (mercury and lead) are playing havoc with the Common Loon. Mercury poisoning "played a major role in the death of over 2,500 loons in their wintering area in 1983." Also, of the adult loon carcasses examined by the LPC, "nearly one-fourth...died due to lead poisoning....invariably a small fishing sinker picked up with gravel....Another one-fourth of the loons killed were found tangled in monofilament or with ingested fishing lures." Thus, in contrast to the early days when hunters and egg collectors caused loon deaths by the hundreds, more loons today succumb to fishermen!

Despite the relative success of New Hampshire's LPC, an article in the 9/86 issue of *Natural History* by Canadian researcher Robert Alvo ("Lost Loons of the Northern Lakes" 95: 58) adds a very sad note to the loon story. He found that 62 percent of loon broods on acidic lakes dies compared to 14 percent on healthy lakes. Thus, acid rain is one more factor of an increasingly hostile environment that may still forever the lovely tremolo of the Common Loon.

D.R.A.