

Florida Field Nat. 26(3):94-96, 1998.

GULL-BILLED TERN FLEDGED ON A ROOF IN NORTHWEST FLORIDA

GARY L. SPRANDEL¹, HEATHER A. BOLTE², AND DAVID T. COBB²

¹Florida Game and Fresh Water Fish Commission, Route 7 Box 3055
Quincy, Florida 32351

²Florida Game and Fresh Water Fish Commission, 620 S. Meridian Street
Tallahassee, Florida 32399

Several species of larids, including Least Terns (*Sterna antillarum*), Roseate Terns (*S. dougallii*), and Black Skimmers (*Rynchops niger*), have fledged young on roofs in Florida (Greene and Kale 1976, Fisk 1978, Hovis and Robson 1989, Gore 1991, Gore and Kinnison 1991, Cimbaro 1993, Coburn 1995, Smith 1996, Hovis and Gore 1997, Zambrano et al. 1997, Bolte and Cobb 1998). Additionally Royal Terns (*S. maxima*) (Toland and Gilbert 1987) and Gull-billed Terns (*S. nilotica*) (Coburn 1996) have unsuccessfully attempted roof-nesting in this state. This note reports the successful fledging of a Gull-billed Tern chick from a roof in Panama City, Florida.

In 1996, we monitored a colony of Black Skimmers on the roof of the Sears department store, Panama City, Bay County (30°11.1'N, 85°38.9'W). The 1.0 ha, 10-m high roof was covered with approx. 1 to 2 cm of light-brown gravel of 0.5 to 1.0 cm diameter. On 2 July, 17 Black Skimmer nests were distributed over 75 percent of the roof. One adult Gull-billed Tern was on a scrape with two undamaged eggs and one slightly dented (<0.5 cm) egg in the middle of the roof near a rubber partition. The adult tern flushed from the nest on our approach. We also heard two Gull-billed Terns calling, saw three flying, and observed one mobile chick. Although we found no other Gull-billed Tern nests, we cannot say if the chick was from the same nest as that with eggs. On 12 July, an adult Gull-billed Tern was standing by the nest, one was circling overhead, and three eggs were still present. The chick appeared about ready to fledge (Fig. 1). The lower mandible on the chick was predominantly orange, the upper mandible black, and the legs a dirty orange-pink. On 17 July, one adult was standing by the three eggs, of which one had a crack approximately 1 cm in length. The fledgling was flying over the roof, suggesting it was at least 28 to 35 days old (Clapp et al. 1983, Parnell et al. 1995). On 24 July, all three eggs had cracks >1 cm long, an adult Gull-billed Tern was observed flying, and the chick was not seen.

Gull-billed Terns have produced young on roofs in Texas and Louisiana (Purrington 1990, Smalley et al. 1991), on causeways (Smith et al. 1993), dredged-material islands, and disturbed inland habitats (e.g. phosphate mines) (Layne et al. 1977, Smith and Alvear 1997). To our knowledge, however, this is the first Gull-billed tern chick to fledge from a roof in Florida. The roof nesting sites in Bay County reported here and by Coburn (1996) were 3.8 and 2.5 km, respectively, from the bay and even closer to small bayous. In 1997, we returned and saw an adult flying over the Sears roof on 9 July, but no nests were located during the summer. As in previous reports from northwest Florida, Gull-billed Terns show a strong affinity for nesting with Black Skimmers (Smith et al. 1993, Smith and Gore 1996, Coburn 1996). Since Black Skimmers have continued to use roofs in recent years (Hovis and Gore 1997, Zambrano et al. 1997, Bolte et al. 1998) perhaps the Gull-billed Tern will also continue to be found on roofs.

Three Gull-billed Tern eggs probably cracked on the gravel substrate. Gore (1987) and Coburn (1995) noted that relatively large birds like the Black Skimmer are more likely to crack their eggs on roofs than the smaller Least Tern, because the thin layer of gravel prevents skimmers from making a scrape deep enough to support the weight of



Figure 1. Gull-Billed Tern chick, estimated 23-30 days old, at Sears Roof, Panama City, Bay County, Florida, 12 July 1996.

the incubating adult (Fisk 1978). On rooftops, perhaps eggs of Gull-billed Terns, which are intermediate in size to Least Terns and Black Skimmers, will crack more often than eggs of Least Terns but less often than Black Skimmers. In order to understand what steps may be taken to avoid excessive egg loss on roofs, studies of the effect of substrate type, color and gravel depth on reproductive success should continue as proposed by Coburn (1995), Coburn et al. (1997), and Bolte and Cobb (1998). Future observations of nesting on artificial sites should report substrate depth, substrate size and color, and nest height above ground.

We thank the management at Sears in Panama City for allowing us access to the roof. Todd Engstrom, Jeff Gore, Hank Smith, Brian Toland, Glenn Reynolds, and an anonymous reviewer provided helpful comments on earlier drafts of this manuscript.

LITERATURE CITED

- BOLTE, H. A., AND D. T. COBB. 1998. An evaluation of management techniques to increase the reproductive success of roof-nesting Black Skimmers in northwest Florida. Florida Game and Fresh Water Fish Comm. Annual Perf. Rept. Tallahassee.
- CIMBARO, J. S. 1993. Hatching, fledging success and thermoregulatory behaviors of roof-nesting Least Terns (*Sterna antillarum*). M.S. thesis. Florida Atlantic Univ., Boca Raton.
- CLAPP, R. B., D. MORGAN-JACOBS, AND R. C. BANKS. 1983. Marine birds of the southeastern United States and Gulf of Mexico. Part III. Charadriiformes. U.S. Fish. Wildl. Serv., Div. Biol. Serv., FWS/OBS-83/30. Washington, D.C.
- COBURN, L. M. 1995. Reproductive success of roof and ground nesting Black Skimmers (*Rynchops niger*) in northwest Florida. M.S. thesis. Univ. W. Florida, Pensacola.
- COBURN, L. M. 1996. Gull-billed Tern nesting on a roof in northwest Florida. Florida Field Nat. 24:76-77.

- COBURN, L. M., D. COBB, AND J. GORE. 1997. Management opportunities and techniques for roof- and ground-nesting Black Skimmers in Florida. Florida Game and Fresh Water Fish Comm. Final Perf. Rept. Tallahassee.
- FISK, E. J. 1978. Roof-nesting terns, skimmers, and plovers in Florida. Florida Field Nat. 6:1-8.
- GORE, J. A. 1987. Black Skimmers nesting on roofs in northwest Florida. Florida Field Nat. 15:77-79.
- GORE, J. A. 1991. Distribution and abundance of nesting Least Terns and Black Skimmers in northwest Florida. Florida Field Nat. 19:65-72.
- GORE, J. A., AND M. J. KINNISON. 1991. Hatching success in roof and ground colonies of Least Terns. Condor 93:759-762.
- GREENE, L. L., AND H. W. KALE. 1976. Roof nesting by Black Skimmers. Florida Field Nat. 4:15-17.
- HOVIS, J. A., AND J. A. GORE. 1997. Nesting shorebird survey. Florida Game and Fresh Water Fish Comm. Final Perf. Rept. Tallahassee.
- HOVIS, J. A., AND M. S. ROBSON. 1989. Breeding status and distribution of the Least Tern in the Florida Keys. Florida Field Nat. 17:61-66.
- LAYNE, J. N., J. A. STALLCUP, G. E. WOOLFENDEN, M. N. MCCAULEY, AND D. J. WORLEY. 1977. Fish and wildlife inventory of the seven-county region included in the central Florida phosphate industry areawide environmental impact study. U.S. Fish. Wild. Serv. Contract No. 14-16-0097-77-005, Washington, D.C.
- PARNELL, J. F., R. M. ERWIN, AND K. C. MOLINA. 1995. Gull-billed Tern (*Sterna nilotica*). In The Birds of North America, No 140. A. Poole and F. Gill (Eds.). Amer. Ornithol. Union and Philadelphia Acad. of Nat. Sci., Philadelphia.
- PURINGTON, R. D. 1990. Central Southern Region. Audubon Field Notes: 44(5):1143-1147.
- SMALLEY, A. E., G. B. SMALLEY, A. J. MUELLER, AND B. C. THOMPSON. 1991. Roof-nesting Gull-billed Terns in Louisiana and Texas. J. La. Ornith. 2(1):18-20.
- SMITH, H. T. 1996. Roseate Tern. Pp. 247-257 In Rare and endangered biota of Florida: Volume V. Birds. (J. A. Rodgers, H. W. Kale II, and H. T. Smith, Eds.). Univ. Press of Florida, Gainesville.
- SMITH, H. T., AND E. M. ALVEAR. 1997. Recent breeding reports of the Gull-billed Tern in Florida . . . status undetermined. Florida Naturalist 70(1):22-23.
- SMITH, H. T., AND J. A. GORE. 1996. Gull-billed Tern. Pp. 624-632 In Rare and endangered biota of Florida: Volume V. Birds. J. A. Rodgers, H. W. Kale II, and H. T. Smith (Eds.). Univ. Press of Florida, Gainesville.
- SMITH, H. T., J. A. GORE, W. W. MILEY, H. L. EDMISTON, AND J. A. RODGERS. 1993. Recent nesting of Gull-billed Terns in northwest Florida. Florida Field Nat. 21:80-82.
- TOLAND, B., AND T. GILBERT. 1987. Roof nesting by Royal Terns in Vero Beach, Florida. Florida Field Nat. 15:80-82.
- ZAMBRANO, R., M. S. ROBSON, D. Y. CHARNETZKY, AND H. T. SMITH. 1997. Distribution and status of Least Tern nesting colonies in southeast Florida. Florida Field Nat. 25:85-91.