# News, Notes, Comments

## ERRATA

Please make the following corrections to the Ms entitled "Observations of the Decline Over 47 Years, 1970-2016 in an Adirondack Breeding Population of Purple Finches Attributable to Climate Change" Apr-Sep 2018, Vol.43 No.2&3.

- 1). P. 42, R col., L 7 "11 km" should be "50 km"
- 2). P. 43, R col., L 4: "16-29" should be "16-20"
- 3). P. 44, L col., L 6 from bottom: "11 km" should be "50 km"
- 4). P. 44, L col., L 2 from bottom: Remove the "." after Schenectady" and place it after the word "captures" on the next line, then insert two spaces before the word "Even...."
- 5) P. 44, R col., L 15: insert the word "from" between "1994" and "which."
- 6) P. 46, L col., L 19: "61.5/58.5" should be "61.5/38.5."
- 7) P. 46, R col., L 12 from bottom: change "Apr 1959" to "Apr 1954."

On the last cover of *NABB*: Oct - Sept 2018 Vol.43 No. 4, Sept should be **Dec.** 

### ACKNOWLEDGEMENTS and LITERATURE CITED were inadvertenly omitted from the Manuscript "Capture Efficiency of Three Mesh Sizes of Mist Nets" in the last issue of *NABB*. Please place The Following:

## ACKNOWLEDGEMENTS

I thank the volunteers who helped collect this data, and the City of La Verne for allowing me to conduct this study on city-owned land. For assistance in getting copies of articles, I thank Molly Garson at Avinet, James Karr at the University of Washington, and Susanne Kreutzer at Landesamt für Umwelt, Naturschutz und Geologie in Mecklenburg-Vorpommern, Germany. I especially thank my master bander, Walter Sakai, who guided me in cleaning up a disorganized manuscript. All faults in research design, execution and analysis are of course my own.

#### LITERATURE CITED

- Avinet. 2018. www.avinet.com. Accessed 6 May 2018.
- Bleitz, D. 1970. Mist nets and their use. *Inland Bird Banding News* 42:1-14.
- Bub, H. 1991. Bird trapping and bird banding:
  a handbook for trapping methods all over the world. Cornell University Press, Ithaca, NY. (translation by F. Hamerstrom and K. Wuertz-Schaefer of Bub, H. 1978. Vogelfang und Vogelberingung. A. ZiemsenVerlag Wittenberg Lutherstadt, East Germany.)
- DeSante, D.F., J.F. Saracco, C.R.d.V. Alvarez, and S. Morales. 2009. 2009-10 MoSI (Monitoreode Sobrevivencia Invernal) manual: instructions for establishing and operating bird-banding stations as part of the MoSI program. Pt. Reyes Station, CA: The Institute for Bird Populations. http://www. birdpop.org/docs/misc/MoSI-Manual-English. pdf. Accessed 6 May 2018.
- DeSante, D.F., J.F. Saracco, D.R. O'Grady,
  K.M. Burton, and B.L. Walker.
  2004. Methodological considerations of the Monitoring Avian Productivity and Survivorship (MAPS) program. *Studies in Avian Biology* 29:28-45.
- Dorsch, H. 1983. Die Fangeffizienz zweier Vogelfangnetztypen. Berichte aus der Vogelwarte Hiddensee 4:129-132.
- Faaborg, J., W.J. Arendt, and K.M. Dugger. 2004. Bird population studies in Puerto Rico using mist nets: general patterns and comparisons with point counts. *Studies in Avian Biology* 29:144-150.
- Heimerdinger, M.A. and R.C. Leberman. 1966. The comparative efficiency of 30 and 36 mm mesh in mist nets. *Bird-Banding* 37:280-285.
- Jenni, L., M. Leuenberger, and F. Rampazzi.1996. Capture efficiency of mist nets with comments on their role in the assessment of passerine habitat use. *Journal of Field Ornithology* 67:263-274.
- Karr, J.R. 1979. On the use of mist nets in the study of bird communities. *Inland Bird Banding* 51:1-10.
- Karr, J.R. 1981. Surveying birds with mist nets. *Studies in Avian Biology* 6:62-67.

Apr - Jun & Jul - Sep 2019

Keyes, B.E. and C.E. Grue. 1982. Capturing birds
with mist nets: a review. North American Bird
<i>Bander</i> 7:2-14.
Ludwig, F.E. 1969. More about nets. Inland Bird
Banding News 41:48-49.
McClure, E. 1984. Boxwood Press Pacific-
Grove, CA.
North American Banding Council. 2001
The North American bander's study guide.
Point Reyes Station, CA.
http://www.nabanding.net/wp-content/
uploads/2012/04/STUDYGUIDE1.pdf.
Pardieck, K. and R. Waide. 1992.
Mesh size as a factor in avian community
studies using mist nets. Journal of Field
Ornithology 63:250-255.
Piratelli, A. 2003. Mesh size and bird capture rates in Mato Grosso do Sul State, Brazil. <i>Brazilian</i>
Journal of Biology 63:105-111.
http://dx.doi.org/10.1590/S15196984200300014.
Accessed 6 May 2018.
Pyle, P. 1997. Identification Guide to North
American Birds, part 1: Columbidae to
Ploceidae. Slate Creek Press, Bolinas, CA.
Ralph, C.J., G.R. Geupel, P. Pyle, T.E. Martin, and
D.F.DeSante. 1993. Handbook of field
methods for monitoring landbirds.
Albany, CA: United States Forest Service,
Pacific Southwest
Research Station, General Technical
ReportPSW-GTR-144-www.
http://dx.doi.org/10.3996/012016-
JFWM-002.S5. Accessed 5 May 2018.
Sibley, D.A. 2016. Sibley Birds West: Field Guide to
Birds of Western North America,
Knopf. New York, N.Y.
Whitman, A.A. 2004. Use of mist nets for study of
neotropical bird communities. Studies in
Avian Biology 29:161-167.
The Table of Contents Oct -Sept 2018 Vol.43
No. 4, Jan - Mar 2019 Vol.44 No. 1 should be as
follows under:
Western Regional News
147 Western Bird Banding Association

Annual Meeting - Brighton, Colorado 147 Annual Summary of Birds Banded in 2017, in the WBBA Region *Walter H. SAKAI*, WBBA EDITOR

## Two Same-season, Long-distance Recaptures of Alder Flycatchers in Alaska

#### Introduction

The accurate identification of Alder and Willow flycatchers in the hand by banders has been problematic where the two co-occur during migration. Species' identification is easier with breeding populations which are, for the most part, geographically separated.

*Captures in Nebraska*: WCS banded an Alder Flycatcher at Cedar Point Biological Station in Nebraska on 28 May 1993 that was recaptured in Alaska. This recapture was reported in Brown et al. (1996) and Sharpe et. al (2001), but without specifics on sites, dates, lat/long, and band number. Another Alder Flycatcher banded by WCS in central Nebraska at Cottonwood Ranch on 5 Jun 2001 reported in Johnsgard (2018) without specifics was also recaptured in Alaska. Our total capture effort in Nebraska amounted to 53,536.6 net-hr with 68.9% accomplished during seven seasons at Cedar Point Biological Station, and 31.1% during six seasons in central Nebraska.

In submitting banding records, WCS initially identified these Empidonax species based mostly on the color of upperparts (before the publication of Pyle 1997). This prompted a terse letter from M. Kathleen Klimkiewicz (Banding Lab Biologist). She changed all Willow and Alder flycatcher identifications to Traill's Flycatchers because of the difficulty in separating these species. That policy apparently continues to be followed by the Banding Lab. WCS was encouraged to continue recording the color-based identity of the two species by Linda R. Brown, who, with Paul A. Johnsgard, had recently finished a museum project on separating the two species. Johnsgard describes the Alder Flycatcher as "apparently a regular migrant in Nebraska," (Johnsgard 2018), and Ely (1970) documents heavy migrations of Traill's Flycatchers through adjacent west-central Kansas. WCS continued recording the color types as separate species in his notes. At Cedar Point Biological Station, 177 birds were banded as