

# ORNITOLOGIA NEOTROPICAL

EDITOR

ANDRÉ-ALEXANDER WELLER

## INDEX TO VOLUME 22, 2011

(Authors, key words, families, scientific names)

- Acoustic adaptation, 219–227  
 Adaptations, 553–568  
*Adenostemma platyphyllum*, 207–217  
*Agelaius cyanopus*, 387–404  
 Agricultural landscape, 539–551  
*Agriornis andicola*, 517–537  
 Albardón, 387–404  
*Alectrurus risora*, 517–537  
*Alectrurus tricolor*, 517–537  
*Alipiopsitta xanthops*, 79–88  
 Allometric equation, 553–568  
 Álvarez, M. E., 615–621  
*Amazilia beryllina*, 229–246  
*Amazilia decora*, 421–436  
*Amazilia tzacatl*, 421–436  
*Amazona aestiva*, 459–463, 517–537  
*Amazona oratrix*, 229–246  
*Amazona pretrei*, 517–537  
*Amazona vinacea*, 517–537  
*Amazonetta brasiliensis*, 517–537  
 American Oystercatcher, 505–516  
*Ammodramus humeralis*, 517–537  
*Anas discors*, 517–537  
*Anas flavirostris*, 465–469  
 Ancestor reconstruction, 553–568  
 Andean Condor, 161–172, 257–265  
 Anderson, D. J., 347–359  
 Andes, 601–614  
 Angulo-Gastélum, U. T., 131–142  
 Annual survival, 421–436  
*Anodorhynchus glaucus*, 517–537  
 Anthropogenic disturbances, 589–599  
*Anthus nattereri*, 517–537  
*Antilophia galeata*, 69–77  
 Apolo, 1–14  
 Apple snail, 281–293  
*Aptenodytes forsteri*, 517–537  
*Aptenodytes patagonicus*, 517–537  
*Ara ararauna*, 329–338  
*Ara chloropterus*, 517–537  
*Ara macao*, 143–153  
*Ara militaris*, 517–537  
 Aramburú, R. M., 111–119, 459–463  
*Aramides cajanea*, 517–537  
*Aramides saracura*, 517–537  
*Aratinga acuticaudata*, 459–463, 517–537  
*Aratinga aurea*, 517–537  
*Aratinga leucophthalmus*, 517–537  
*Aratinga nana*, 229–246  
 Arbeiter, S., 39–57  
 Arboreal roosting, 477–479  
*Ardea cocoi*, 517–537  
 Ardiles, K., 121–130  
*Arenaria interpres*, 517–537  
 Areta, J. I., 219–227  
 Argentina, 281–293, 313–315, 387–404, 453–457, 517–537  
*Arramom aurantiirostris*, 421–436  
 Arriaga-Weiss, S., 229–246  
*Asio flammeus*, 207–217, 517–537  
*Aspidosperma quebrachoblanco*, 111–119  
*Asthenes pyrrholeuca*, 387–404  
 Astore, V., 161–172  
 Astudillo, P. X., 257–265  
*Athene cunicularia*, 517–537  
 Atlantic forest, 195–206  
 Avalos, V. del R., 1–14  
 Avendaño, C., 39–57  
 Avian community composition, 421–436  
 Avian diversity, 387–404  
 Avifauna, 229–246  
 Azure-rumped Tanager, 39–57  
 Bahamonde-Viuneza, D., 471–475  
 Bananaquit, 421–436  
 Band-bellied Owl, 471–475  
 Barranca, 387–404  
 Barrientos, C., 121–130  
*Basileuterus leucoblepharus*, 387–404  
 Bayly, N.-J., 339–345  
 Beadle, D., 623–626  
 Belize, 339–345  
 Beltzer, A. H., 387–404  
 Berkunsky, I., 459–463  
*Biatas nigropectus*, 517–537  
 Bilenca, D., 295–305  
 Biondi, L. M., 453–457  
 Bird communities, 229–246  
 Bird conservation, 229–246  
 Bird-plant interactions, 69–77  
 Black-cheeked Ant-Tanager, 421–436

- Black-faced Ant-Thrush, 421–436  
 Black-hooded Antshrike, 421–436  
 Blake, J. G., 387–404, 437–445  
 Blendinger, P. G., 615–621  
 Blue crowned Parakeet, 459–463  
 Blue-crowned Manakin, 421–436  
 Blue-fronted Parrot, 459–463  
 Bodrati, A., 195–206  
 Bolivia, 155–158  
 Bonaccorso, E., 471–475  
 Brazil, 319–327, 329–338  
 Breeding biology, 121–130, 369–378  
 Breeding success, 131–142  
 Brightsmith, D. A., 143–153, 627–631  
 Brilliant Sunangel, 601–614  
 Broadleaf forest, 39–57  
 Brood parasites, 553–568  
 Brooks, D. M., 267–279  
*Brotogeris versicolorus*, 517–537  
 Brown-hooded Gull, 505–516  
*Bubo virginianus*, 517–537  
*Bubulcus ibis*, 517–537  
 Buchán, E., 39–57  
 Buenos Aires, 295–305  
 Burge, A., 39–57  
*Bursera graveolens*, 207–217  
*Buteo albicaudatus*, 517–537  
*Buteo magnirostris*, 517–537  
*Buteo nitidus*, 517–537  
*Buteo swainsoni*, 517–537  
*Buteo ventralis*, 405–420  
*Buteogallus meridionalis*, 517–537  
*Butorides striata*, 517–537  
 Caatinga, 15–24  
*Cacicus haemorrhoous*, 313–315  
 Cadena, C. D., 59–67  
 Cadena-Ortiz, H., 471–475  
*Cairina moschata*, 229–246, 517–537  
 Cajas National Park, 257–265  
*Camptostoma obsoletum*, 15–24, 387–404  
*Caprimulgus parvulus*, 15–24, 517–537  
*Caprimulgus rufus*, 517–537  
*Capsicum frutescens*, 207–217  
*Caracara plancus*, 517–537  
*Cardiospermum galapageium*, 207–217  
*Cariama cristata*, 517–537  
 Caribbean, 447–451  
 Carmiol's Tanager, 155–158  
 Carrera, J., 459–463  
 Carvajal-Rueda, A., 539–551  
*Caryocar brasiliense*, 329–338  
*Castela galapageia*, 207–217  
 Castillo, M. A., 577–587  
 Castillo-Guerrero, J. A., 131–142  
*Cathartacta antarctica*, 517–537  
*Cathartacta chilensis*, 517–537  
*Cathartacta macormicki*, 517–537  
*Cathartes aura*, 517–537  
*Cathartes burrovianus*, 517–537  
*Catharus ustulatus*, 421–436  
 Cavities, 465–469  
*Cedrus libani*, 111–119  
*Celtis ehrenbergiana*, 111–119  
*Celtis tala*, 111–119  
 Centla Marshes Biosphere Reserve, 229–246  
 Central America, 339–345  
 Cerrado, 319–327, 329–338  
 Cerros del Sira, 267–279  
 Cerulean-capped Manakin, 623–626  
*Charadrius semipalmatus*, 517–537  
*Charitospiza eucosma*, 517–537  
 Charming Hummingbird, 421–436  
 Cherrie's Tanager, 421–436  
 Chestnut-backed Antbird, 421–436  
 Chick age, 361–368  
 Chick growth, 143–153  
 Chile, 247–256, 369–378, 465–469  
 Chiloé Island, 103–110  
*Chionis albus*, 517–537  
*Chiroxiphia pareola*, 437–445  
*Chloephaga rubidiceps*, 517–537  
*Chlorothraupis carmioli*, 155–158  
*Chordeiles pusillus*, 517–537  
*Chroicocephalus cirrocephalus*, 505–516  
*Chroicocephalus maculipennis*, 505–516  
*Chunga burmeisteri*, 517–537  
*Ciconia maguari*, 517–537  
*Cinclus schulzii*, 517–537  
 Circular statistics, 161–172  
*Claravis goedefrida*, 517–537

- Claravis pretiosa*, 517–537  
 Class, A.-M., 89–102  
 Climatic variation, 143–153  
*Cnemotriccus fuscatus*, 517–537  
*Coccyzus melacoryphus*, 15–24, 207–217, 387–404  
*Cochlearius cochlearius*, 517–537  
 Cockle, K. L., 195–206  
 Codesido, M., 295–305  
*Coereba flaveola*, 517–537, 421–436  
 Coffee plantation, 39–57  
*Colaptes campestris*, 517–537  
*Colaptes melanochloros*, 517–537  
 Coleopterans, 471–475  
*Colibri serrirostris*, 517–537  
 Collister, D. M., 421–436  
 Colombia, 539–551  
 Coloniality, 131–142  
 Colony size, 131–142  
 Color spectra, 25–38  
*Columbina cruziana*, 219–227  
*Columbina minuta*, 517–537  
*Columbina picui*, 15–24, 387–404, 517–537  
*Columbina squammata*, 517–537  
*Columbina talpacoti*, 517–537  
 Common Pauraque, 173–185  
 Communication, 219–227  
 Competition, 465–469  
 Conservation, 1–14, 569–576  
 Conservation status, 517–537  
 Constrictor Boa, 459–463  
 Contreras, M. S., 187–194  
 Control methods, 111–119  
 Cooperative breeding, 39–57  
*Coragyps atratus*, 517–537  
*Cordia lutea*, 207–217  
*Cordia sebestena*, 447–451  
*Coryphospiza melanosis*, 517–537  
*Coryphospingus cucullatus*, 387–404, 517–537  
 Costa Rica, 173–185  
 Cowbirds, 553–568  
 Cracid ecology, 267–279  
 Cracidae, 615–621  
*Crax fasciolata*, 517–537  
 Crested Hornero, 459–463  
 Cricetidae, 295–305  
*Croton scouleri*, 207–217  
*Crotaphaga ani*, 207–217, 517–537  
*Crotaphaga major*, 517–537  
*Crypturellus obsoletus*, 517–537  
*Crypturellus parvirostris*, 517–537  
*Crypturellus tataupa*, 517–537  
 Cuban Green Woodpecker, 447–451  
 Cuervo, P., 517–537  
 Cueto, V. R., 483–494  
*Culicivora caudacuta*, 517–537  
*Cupressus sempervirens*, 111–119  
*Cyanerpes cyaneus*, 421–436  
*Cyanocompsa brissonii*, 517–537  
*Cyanocorax caeruleus*, 517–537  
*Cyclarbis gujanensis*, 517–537  
*Cygnus melanoryphus*, 517–537  
 Cytogenetics, 517–537  
*Dacnis cayana*, 517–537  
 Dalsgaard, B., 447–451  
 DAP, 111–119  
*Daption capense*, 517–537  
 David, S., 307–311  
 de Araújo, C. B., 79–88  
 de León Lux, A., 39–57  
 de León Lux, J., 39–57  
 De Martino, E., 161–172  
*Dendrocincla anabatina*, 229–246  
*Dendrocygna bicolor*, 517–537  
*Dendrocygna viduata*, 517–537  
*Dendroica petechia*, 229–246  
*Dendroica petechia aureola*, 207–217  
*Dendroica tigrina*, 229–246  
 Díaz, I. A., 103–110  
 Diet, 79–88, 471–475  
*Digitaria horizontalis*, 207–217  
*Diomedea epomophora*, 517–537  
*Diomedea exulans*, 517–537  
 Distribution model, 1–14  
*Diuca diuca*, 369–378  
 Diuca Finch, 369–378  
 Diversity, 15–24  
 Dot-winged Antwren, 421–436  
*Dryocopus galeatus*, 517–537  
 Durães, R., 437–445

- Durand, A., 623–626  
*Dysithamnus mentalis*, 517–537  
 Eastern Kingbird, 421–436  
 Ecuador, 437–445, 471–475, 601–614  
 Egg mass, 553–568  
 Eggs, 121–130, 155–158, 307–311  
 Eisermann, K., 39–57  
*Elaenia chilensis*, 15–24  
*Elaenia flavogaster*, 15–24  
*Elaenia mesoleuca*, 15–24  
*Elaenia parvirostris*, 517–537  
*Elaenia spectabilis*, 15–24, 387–404  
 Emberizidae, 319–327  
*Emberizoides herbicola*, 517–537  
 Emergent rocks, 589–599  
*Empidonax alnorum*, 339–345, 517–537  
*Empidonax flaviventris*, 339–345  
 Empidonax flycatchers, 339–345  
*Empidonax minimus*, 339–345  
*Empidonax traillii*, 339–345  
*Empidonax virescens*, 339–345  
*Empidonomus varius*, 15–24  
 Endemic avifauna, 267–279  
 Engilis, Jr., A., 247–256  
*Enicognathus leptorhynchus*, 103–110, 465–469  
 Enríquez, P. L., 577–587  
 Escalante, I., 173–185  
 Escobar, O., 229–246  
*Eucalyptus* sp., 111–119  
*Eudromia elegans*, 517–537  
*Eudyptes chrysocome*, 517–537  
*Eudyptes chrysophrys*, 517–537  
*Eudyptes robustus*, 517–537  
*Eudyptes schlegeli*, 517–537  
*Eudyptes sclateri*, 517–537  
*Euscarthmus meloryphus*, 15–24, 387–404  
 Exaptations, 553–568  
 Faegre, S. K. I., 459–463  
*Falco peregrinus*, 517–537  
*Falco sparverius*, 517–537  
 Favero, M., 505–516  
 Feeding behavior, 79–88  
 Feeding habits, 495–504  
 Feeding station, 257–265  
 Feeley, K. J., 347–359  
 Fernández Ajó, A. A., 361–368  
 Ferrari, S. F., 15–24  
 Ferruginous Pygmy-Owl, 577–587  
 Figueroa R., R. A., 121–130, 405–420  
*Fluvicola albiventer*, 387–404  
 Food niche breadth, 295–305  
 Food provisioning, 361–368  
 Foraging behavior, 247–256  
 Foraging strategy, 453–457, 505–516  
 Foraging tactics, 495–504  
*Formicarius analis*, 421–436  
*Forpus xanthopterygius*, 517–537  
 Fraga, R. M., 313–315  
 Fragmentation, 421–436  
*Fregetta tropica*, 517–537  
 Frugivory, 69–77, 207–217, 615–621  
 Furnariidae, 247–256, 477–479  
*Furnarius cristatus*, 459–463  
*Furnarius rufus*, 387–404, 517–537  
 Galapagos, 207–217  
 Gallery forest, 69–77  
*Gallinula chloropus*, 517–537  
 García, G. O., 505–516  
 Gastañaga, M., 267–279  
 Gatto, A., 361–368  
*Geoffroea decorticans*, 111–119  
 Geographic distribution, 483–494  
 Geospizinae, 207–217  
*Geothlypis aequinoctialis*, 387–404  
*Geothlypis poliocephala*, 229–246  
*Geothlypis trichas*, 229–246  
*Geotrygon montana*, 517–537  
*Geranoaetus melanoleucus*, 517–537  
*Geranospiza caerulescens*, 517–537  
 GIS, 161–172  
*Glaucidium ridgwayi*, 577–587  
*Glyptohynchus spirurus*, 421–436  
*Gnorimopsar chopi*, 517–537  
 Gómez, C., 339–345  
 Gómez, Llamas, C. J., 187–194  
 González-Acuña, D., 121–130  
 González-Fischer, C. M., 295–305  
 González-García, F., 229–246  
 Gorla, N., 517–537  
 Graham, C. H., 257–265

- Granivorous birds, 483–494  
 Gressler, D. T., 319–327  
 Grey-hooded Gull, 505–516  
 Growth rate, 143–153  
 Guatemala, 39–57  
*Gubernatrix cristata*, 517–537  
 Guerrero, A. M., 207–217  
*Guira guira*, 517–537, 387–404  
 Gussoni, C. O. A., 495–504  
*Gymnopelia*, 219–227  
*Habia atrimaxillaris*, 421–436  
 Habitat, 187–194  
 Habitat use, 539–551  
*Haematopus palliatus*, 505–516  
*Haplospiza unicolor*, 517–537  
*Harpia harpyja*, 517–537  
*Harpyhaliaetus coronatus*, 517–537  
 Heermann's Gull, 131–142  
*Helianthus splendidus* spec. nov., 601–614  
*Helianthus splendidus pyropus* subsp. nov., 601–614  
*Helianthus splendidus splendidus* subsp. nov., 601–614  
*Helianthus viola*, 601–614  
 Helmeted Curassow, 267–279  
 Helmeted Manakin, 69–77  
*Hemithraupis guira*, 15–24  
 Hennessey, B., 267–279  
 Hidalgo, J. R., 437–445  
 High Andes, 219–227  
*Hirundo rustica*, 517–537  
 Hispaniolan Woodpecker, 447–451  
 Home range, 161–172  
 Horned Curassow, 267–279  
*Hydropsalis torquata*, 517–537  
*Hylopetes perspicillatus*, 421–436  
*Hylophilus decurtatus*, 421–436  
*Hylophilus ochraceiceps*, 421–436  
*Icterus chrysater*, 229–246  
*Icterus pyrrhopterus*, 313–315  
 Incubation behavior, 59–67  
 Incubation period, 553–568  
 Introduced birds, 207–217  
 Introduced plants, 207–217  
 Invasive, 207–217  
 Ippi, S., 103–110  
 Islands, 207–217  
*Jabiru mycteria*, 229–246, 517–537  
 Jácome, N. -L., 161–172  
 Jiménez, J. E., 465–469  
 Kacoliris, F. P., 459–463  
 Karyotype, 517–537  
 Kelt, D. A., 247–256  
 Kirwan, G. M., 623–626  
 Kittlein, M. J., 281–293  
 Kleptoparasitism, 453–457, 505–516  
*Knipolegus cyanirostris*, 517–537  
*Lantana camara*, 207–217  
*Lantana peduncularis*, 207–217  
*Laporteaaestuans*, 207–217  
 Lara, J., 121–130  
*Larus atlanticus*, 517–537  
*Larus dominicanus*, 517–537  
*Larus fuscus*, 517–537  
*Larus heermanni*, 131–142  
*Laterallus melanophaius*, 517–537  
 Latta, S. C., 257–265  
*Legatus leucophaius*, 517–537  
 Lek, 437–445  
*Lepidocolaptes angustirostris*, 459–463  
*Lepidocolaptes falcinellus*, 195–206  
*Lepidothrix coeruleocapilla*, 623–626  
*Lepidothrix coronata*, 421–436, 437–445  
*Leptasthenura aegithaloides*, 247–256  
*Leptodon cayanensis*, 517–537  
*Leptophaps*, 219–227  
*Leptotila conoveri*, 539–551  
*Leptotila rufaxilla*, 517–537  
*Leptotila verreauxi*, 387–404, 517–537  
 Lesser Greenlet, 421–436  
*Lochmias nematura*, 517–537  
 Loera, C. J., 187–194  
 Logistic model, 143–153  
 Loiselle, B. A., 437–445  
 Londoño, G. A., 59–67, 307–311  
 Long-billed Hermit, 421–436  
 Lopez de Casenave, J., 483–494  
 López, G., 39–57  
 Lorenzón, R. E., 387–404  
 Losada-Prado, S., 539–551

- Lotic environments, 589–599  
 Lower Paraná, 387–404  
*Lurocalis semitorquatus*, 517–537  
 MacGregor-Fors, I., 569–576  
*Machaeropterus regulus*, 437–445  
*Machetornis rixosus*, 387–404  
 MacLeod, R., 267–279  
*Macronectes giganteus*, 517–537  
*Manacus aurantiacus*, 421–436  
 Management, 111–119  
 Manakin, 437–445  
 Maness, T. J., 347–359  
 Mapelli, F. J., 281–293  
 Mar Chiquita Lagoon, 453–457  
 Marcondes-Machado, L. O., 79–88  
 Marin, M., 369–378  
 Marini, M. Á., 319–327  
 Mark-recapture, 421–436  
 Marone, L., 483–494  
 Martínez, O., 155–158  
 Matorral, 247–256  
 Medel H., J., 405–420  
*Megacyrle torquata*, 517–537  
*Megarhynchus pitangua*, 15–24, 517–537  
*Megascops guatemalae*, 577–587  
*Melanerpes flavifrons*, 195–206  
 Mellink, E., 131–142  
 Melo, C., 69–77  
 Mena, M., 161–172  
*Merganetta armata*, 589–599  
*Mergus octosetaceus*, 517–537  
 Mermoz, M. E., 553–568  
*Metriopelia*, 219–227  
*Metriopelia aymara*, 219–227  
*Metriopelia ceciliae*, 219–227  
*Metriopelia melanoptera*, 219–227  
*Metriopelia morenoi*, 219–227  
 Mexico, 569–576  
*Miconia robinsoniana*, 207–217  
*Microrhopias quixensis*, 421–436  
 Migration, 15–24  
 Migration routes, 339–345  
 Milesi, F. A., 483–494  
*Mihago chimachima*, 517–537  
*Mihago chimango*, 453–457  
*Haematopus palliatus*, 453–457  
 García, G. O., 453–457  
*Mimus saturninus*, 517–537  
 Minimum convex polygon, 161–172  
*Mionectes oleaginous*, 421–436  
*Molothrus*, 553–568  
*Molothrus bonariensis*, 369–378, 387–404, 517–537  
*Momordica charantia*, 207–217  
 Monitoring, 257–265  
 Monk Parakeet, 111–119  
 Monteleone, D., 219–227  
 Moore, I.-T., 89–102  
 Moreno, L., 121–130  
*Morphnus guianensis*, 517–537  
*Morus capensis*, 517–537  
 Mottled Owl, 577–587  
 Mountain rivers, 589–599  
*Myiarchus ferox*, 517–537  
*Mycteria americana*, 517–537  
*Myiarchus magnirostris*, 207–217  
*Myiarchus tyrannulus*, 15–24  
*Myiodynastes maculatus*, 517–537  
*Myiopagis viridicata*, 15–24  
*Myiophobus fasciatus*, 15–24  
*Myiopsitta monachus*, 111–119, 387–404  
*Myiozetetes similis*, 517–537  
*Myrmeciza exsul*, 421–436  
*Nandayus nenday*, 517–537  
 Narrow-billed Woodcreeper, 459–463  
 Native birds, 207–217  
 Native plants, 207–217  
 Natural history, 39–57, 319–327, 569–576  
 Nazca Booby, 347–359  
 Neotropical Austral migrants, 483–494  
*Nephelomyias ochraceiventris.*, 59–67  
*Nesomimus parvulus*, 207–217  
*Nesotriccus ridgwayi*, 207–217  
 Nest, 121–130, 155–158, 307–310  
 Nest architecture, 623–626  
 Nest attentiveness., 59–67  
 Nest defense, 195–206  
 Nest height, 111–119  
 Nest site, 111–119  
 Nest site selection, 39–57

- Nest success, 39–57  
 Nesting, 313–315, 465–469, 623–626  
 Nesting biology, 59–67  
 Nesting sites, 405–420  
 Nestling asymptotic body mass, 553–568  
 Nestlings, 155–158, 195–206, 459–463  
 Northeastern Brazil, 15–24  
*Notharchus hyperrhynchus*, 229–246  
*Nothofagus* forest, 103–110  
*Nothura maculosa*, 517–537  
*Numenius borealis*, 517–537  
*Nyctibius griseus*, 517–537  
*Nycticorax nycticorax*, 517–537  
*Nyctidromus albicollis*, 173–185, 517–537  
 Oaxaca, 569–576  
*Oceanites oceanicus*, 517–537  
 Ochraceous-breasted Flycatcher, 59–67  
 Ochre-bellied Flycatcher, 421–436  
*Odontophorus capueira*, 517–537  
*Opuntia*, 207–217  
 Orange-billed Sparrow, 421–436  
 Orange-breasted Bunting, 187–194  
 Orange-chinned Parakeet, 421–436  
 Orange-collared Manakin, 421–436  
 Ornithophily, 447–451  
*Ornithodoros canicollis*, 517–537  
 Orthopterans, 471–475  
 Osa Peninsula, 421–436  
*Pachyramphus castaneus*, 517–537  
*Pachyramphus polychoterus*, 15–24  
 Palkachupa, 1–14  
 Pampean region, 295–305  
*Pandion haliaetus*, 229–246, 517–537  
*Panicum dichotomiflorum*, 207–217  
*Parabuteo unicinctus*, 517–537  
 Páramo, 257–265  
 Parental care, 195–206  
*Paroaria coronata*, 387–404, 517–537  
*Paspalum conjugatum*, 207–217  
*Passerina leclancherii*, 187–194  
*Passerina rositae*, 569–576  
*Patagioenas cayennensis*, 517–537  
*Patagioenas picazuro*, 517–537  
*Patagioenas speciosa*, 517–537  
 Patagonia, 361–368  
*Pauxi koepckeae*, 267–279  
*Pauxi pauxi*, 267–279  
*Pauxi unicornis*, 267–279  
*Pauxi unicornis koepckeae*, 267–279  
*Pauxi unicornis unicornis*, 267–279  
 Peña-Foxon, M., 103–110  
*Penelope dabbenei*, 615–621  
*Penelope obscura*, 517–537  
*Penelope superciliaris*, 517–537  
 Peralta, N. A., 59–67  
 Pérez-Sánchez, C. E., 569–576  
 Peru, 143–153, 601–614  
*Phacellodomus ruber*, 387–404  
*Phaethornis superciliosus*, 421–436  
*Phalcoboenus megalopterus*, 517–537  
 Phenology, 187–194  
*Pheucticus aureoventris*, 517–537  
*Phibalura flavirostris*, 1–14  
*Phleocryptes melanops*, 121–130  
*Phoebe tricolor*, 517–537  
*Phoenicoparrus andinus*, 517–537  
*Phoenicopterus chilensis*, 517–537  
*Phrygilus carbonarius*, 483–494  
*Phylloscartes kronei*, 495–504  
 Phylogeny, 25–38  
*Phytotoma rutila*, 387–404  
*Piana cayana*, 517–537  
 Picidae, 447–451  
*Picumnus cirratus*, 517–537  
*Pinus pinea*, 111–119  
*Pionopsitta pileata*, 517–537  
*Pionus maximiliani*, 517–537  
*Pipile jacutinga*, 517–537  
*Pipra erythrocephala*, 437–445  
*Pipra filicauda*, 437–445  
*Pipra mentalis*, 421–436  
*Pipra pipra*, 437–445  
 Pipridae, 69–77, 437–445, 623–626  
*Piprites pileatus*, 517–537  
*Pitangus sulphuratus*, 15–24, 517–537  
 Plain-mantled Tit-spinetail, 247–256  
*Platalea ajaja*, 517–537  
*Platanus hispanica*, 111–119  
*Platyrinchus leucoryphus*, 517–537  
*Platyrinchus mystaceus*, 517–537

- Pluvialis dominica*, 517–537  
*Pluvialis squatarola*, 517–537  
*Podiceps gallardoi*, 517–537  
*Politi*, N., 589–599  
 Pollination, 447–451  
*Pomacea canaliculata*, 281–293  
*Poospiza baeri*, 517–537  
*Poospiza ornata*, 483–494  
 Population, 1–14  
 Population size, 257–265  
*Populus nigra*, 111–119  
*Porphyrio martinica*, 517–537  
*Portulaca oleracea*, 207–217  
*Porzana albicollis*, 517–537  
*Porzana spiloptera*, 517–537  
 Post-fledging parental care, 347–359  
 Prey capture, 247–256  
 Prey selection, 281–293  
*Primolius maracana*, 517–537  
*Procellaria aequinoctialis*, 517–537  
*Procellaria westlandica*, 517–537  
*Procnias nudicollis*, 517–537  
*Progne chalybea*, 517–537  
*Progne modesta*, 517–537  
*Progne tapera*, 387–404  
*Prosopis affinis*, 111–119  
*Prosopis nigra*, 111–119  
*Psidium guajava*, 207–217  
 Psittacidae, 79–88, 103–110, 329–338  
*Pterodroma incerta*, 517–537  
*Pteroglossus bailloni*, 517–537  
*Pteroglossus castanotis*, 517–537  
*Puffinus creatopus*, 517–537  
 Pujol, E. M., 553–568  
*Pulsatrix melanota*, 471–475  
*Pulsatrix perspicillata*, 517–537  
*Pygochelidon cyanoleuca*, 517–537  
*Pygoscelis adeliae*, 517–537  
*Pygoscelis antarcticus*, 517–537  
*Pygoscelis papua*, 517–537  
*Pyriglen a leucoptera*, 517–537  
*Pyrocephalus rubinus*, 517–537  
*Pyrocephalus rubinus nanus*, 207–217  
*Pyrrhura frontalis*, 517–537  
*Pyrrhura molinae*, 517–537  
*Quercus ilex*, 111–119  
 Quero, M., 517–537  
 Rafael Villegas-Patraca, 569–576  
 Ragusa-Netto, J., 329–338  
 Rainfall, 143–153  
*Rallus antarcticus*, 517–537  
*Rallus limicola*, 229–246  
*Ramphastos dicolorus*, 517–537  
*Ramphastos toco*, 517–537  
*Rampuculus bresilius*, 517–537  
*Rampuculus costaricensis*, 421–436  
 Rangel-Salazar, J. L., 577–587  
*Rattus rattus*, 207–217  
 Ravine, 569–576  
 Rechberger, J., 155–158  
 Red-capped Manakin, 421–436  
 Red-faced Guan, 615–621  
 Red-legged Honeycreeper, 421–436  
 Red-rumped Cacique, 313–315  
 Reproduction, 405–420  
 Reproductive synchrony, 89–102  
 Restinga Tyrannulet, 495–504  
*Rhea americana*, 517–537  
*Rhea pennata*, 517–537  
*Rhynchosciurus rufescens*, 517–537  
 Ribeiro, A. S., 15–24  
*Riparia riparia*, 517–537  
 Riparian, 187–194  
 Riparian forest, 387–404  
 Rivas-Fuenzalida, T., 405–420  
 Rivera, L., 589–599  
 Rocha, P. A., 15–24  
 Ronchi-Virgolini, A. L., 387–404  
 Rose-bellied Bunting, 569–576  
 Rost, J., 187–194  
*Rostrhamus sociabilis*, 229–246, 281–293  
 Royal Tern, 131–142  
*Rubus niveus*, 207–217  
 Rufous-tailed Hawk, 405–420  
 Rufous-tailed Hummingbird, 421–436  
 Ruggera, R. A., 459–463, R. A., 615–621  
 Ruiz-Esparza, D. P. B., 15–24  
 Ruiz-Esparza, J., 15–24  
 Ryder, T. B., 437–445  
 Sagario, C. M., 483–494

- Saltator aurantiirostris*, 387–404  
*Saltator caerulescens*, 387–404, 517–537  
*Saltator maximus*, 229–246  
*Saltator similis*, 387–404, 517–537  
*Saltator* spp., 387–404  
 Sandoval, L., 173–185  
 Santiago-Alarcon, D., 229–246  
 Santos, M. C. de O., 495–504  
*Sarcoramphus papa*, 517–537  
 Sardina Aragón, P., 589–599  
*Sarkidiornis melanotos*, 517–537  
 Satellital transmitter, 161–172  
*Scalesia pedunculata*, 207–217  
 Scarlet Macaw, 143–153  
 Schmitz-Ornés, A., 25–38  
 Schuchmann, K.-L., 25–38  
*Sclateria naevia*, 307–311  
*Sclerurus caudacutus*, 477–479  
*Sclerurus guatemalensis*, 477–479  
*Sclerurus mexicanus*, 477–479  
*Sclerurus rufigularis*, 477–479  
*Scutia spicata*, 207–217  
 Seabirds, 131–142  
 Season, 577–587  
 Seasonal floods, 387–404  
 Seasonal movement, 161–172  
 Seasonality, 15–24, 89–102  
 Seed dispersal, 69–77, 207–217, 615–621  
 Seed predation, 329–338  
*Selenidera maculirostris*, 517–537  
*Setophaga ruticilla*, 229–246  
 Sex-specific parental effort, 347–359  
 Sexual size dimorphism, 347–359  
 Shiny Cowbird, 369–378  
 Shirihai, H., 623–626  
 Short-billed Leaffosser, 477–479  
*Sicalis citrina*, 319–327  
*Sicalis flaveola*, 517–537  
 Silva, A.-M., 69–77  
 Silvered Antbird, 307–311  
 Sira Curassow, 267–279  
*Sittasomus griseicapillus*, 195–206  
 Slender-billed Parakeet, 103–110  
 Small mammals, 295–305  
 Snail Kite, 281–293  
*Solanum americanum*, 207–217  
*Solanum cheesmaniae*, 207–217  
 Song variation, 173–185  
 South American Tern, 361–368  
 Southern temperate rain-forest, 405–420  
 Spatial scale, 483–494  
 Spatial variation, 387–404  
 Species richness, 421–436  
 Spectrophotometry, 25–38  
*Spheniscus magellanicus*, 517–537  
*Spizaetus tyrannus*, 517–537  
*Sporophila bouvreuil*, 15–24  
*Sporophila caerulescens*, 517–537  
*Sporophila cinnamomea*, 517–537  
*Sporophila corvina*, 421–436  
*Sporophila fuscicollis*, 517–537  
*Sporophila frontalis*, 517–537  
*Sporophila leucoptera*, 517–537  
*Sporophila lineola*, 517–537  
*Sporophila nigriceps*, 517–537  
*Sporophila palustris*, 517–537  
*Sporophila zelichi*, 517–537  
*Stelgidopterix ruficollis*, 517–537  
*Stephanophorus diadematus*, 517–537  
*Sterna hirundinacea*, 361–368  
*Sterna hirundo*, 517–537  
*Sterna nilotica*, 517–537  
*Sterna paradisea*, 517–537  
 Stomach analysis, 471–475  
 Stopover-behavior, 339–345  
 Streak-chested Antpitta, 421–436  
*Streptoprocne zonaris*, 517–537  
 Strigidae, 577–587  
 Stripe-tailed Yellow-fin, 319–327  
*Strix hubula*, 517–537  
*Strix hylophila*, 517–537  
*Strix squamulata*, 577–587  
*Strix virgata*, 517–537  
*Sturnella defilippii*, 517–537  
*Suiriri suiriri*, 387–404  
*Sula granti*, 347–359  
 Swainson's Thrush, 421–436  
 Swallow-tailed Cotinga, 1–14  
*Synallaxis albescens*, 387–404  
*Synallaxis frontalis*, 387–404

- Syrigma sibilatrix*, 517–537  
*Tachycineta leucorrhoa*, 387–404  
*Tachycineta thalassina*, 229–246  
*Tachyphonus luctuosus*, 421–436  
*Tachyphonus rufus*, 517–537  
 Tail stiffness, 477–479  
*Tangara cabanisi*, 39–57  
*Tangara cayana*, 517–537  
*Tangara preciosa*, 517–537  
*Taoniscus nanus*, 517–537  
 Tawny-crowned Greenlet, 421–436  
 Taxonomy, 25–38, 601–614  
 Temperate rainforest, 103–110  
 Temporal variation, 387–404  
 Territoriality, 89–102, 577–587  
 Teta, P., 295–305  
*Thalassarche chlororhynchos*, 517–537  
*Thalassarche chrysostoma*, 517–537  
*Thalassarche melanophrys*, 517–537  
*Thalasseus maximus*, 131–142  
 Thamnophilidae, 307–311  
*Thamnophilus bridgesi*, 421–436  
*Thamnophilus caerulescens*, 387–404  
*Thamnophilus doliatus*, 517–537  
*Thlypopsis sordida*, 387–404  
 Thraupidae, 155–158  
*Thraupis bonariensis*, 517–537  
*Thraupis palmarum*, 517–537  
*Thraupis sayaca*, 15–24, 517–537  
*Tigrisoma lineatum*, 229–246, 517–537  
*Tinamus solitarius*, 517–537  
 Tinoco, B. A., 257–265  
 Tolima, 539–551  
 Tolima Dove, 539–551  
*Topaza*, 25–38  
*Topaza pella*, 25–38  
*Topaza pella amaruni*, 25–38  
*Topaza pella jiparana*, 25–38  
*Topaza pella microrhyncha*, 25–38  
*Topaza pella pamprepta*, 25–38  
*Topaza pella pyra*, 25–38  
*Topaza pella smaragdula*, 25–38  
*Topaza pyra*, 25–38  
 Tori, W. P., 437–445  
 Torrent duck, 589–599  
*Tournefortia psilosachya*, 207–217  
*Tournefortia pubescens*, 207–217  
 Transects, 577–587  
*Tringa flavipes*, 517–537  
 Trochilidae, 25–38, 601–614  
*Troglodytes aedon*, 517–537  
*Troglodytes cobbi*, 517–537  
 Tropical, 89–102  
 Tropical deciduous forest, 187–194  
*Turdus albicollis*, 517–537  
*Turdus amaurochalinus*, 15–24, 517–537  
*Turdus leucomelas*, 517–537  
*Turdus rufiventris*, 387–404, 517–537  
*Turdus* spp., 387–404  
 Tye, A., 207–217  
 Tyrannidae, 15–24, 495–504  
*Tyrannus melancholicus*, 15–24, 387–404, 517–537  
*Tyrannus savana*, 387–404, 517–537  
*Tyrannus tyrannus*, 421–436  
*Tyrannus verticalis*, 229–246  
*Tyto alba*, 295–305, 517–537  
 Tytonidae, 295–305  
 Van Els, P., 477–479  
 Variable Oriole, 313–315  
 Variable Seedeater, 421–436  
 Vassallo, A. I., 505–516  
 Vázquez-Pérez, J. R., 577–587  
 Vegetation, 187–194  
 Vegetation types, 577–587  
 Vermiculated Screech-Owl, 577–587  
 Vigo, G., 143–153  
 Violet-throated Sunangel, 601–614  
*Vireo olivaceus*, 15–24, 387–404, 517–537  
*Vireo pallens*, 229–246  
 Vocalization, 437–445  
 Vocalization rate, 437–445  
 Vocalizations, 173–185  
 Voices, 219–227  
*Volatinia jacarina*, 15–24, 517–537  
 Volpe, N. L., 111–119  
*Vultur gryphus*, 161–172, 257–265, 517–537  
 Wedge-billed Woodcreeper, 421–436  
 Weller, A.-A., 601–614  
 West Indies, 447–451

- Westbrock, M. A., 347–359  
Wetlands, 121–130  
White, Jr., T. H., 465–469  
White-shouldered Tanager, 421–436  
Whitney, B.-M., 477–479  
Williams, M., 143–153  
Wilson, A. G., 421–436  
Wilson, S., 421–436  
Wren-like Rushbird, 121–130  
*Xanthopsar flavus*, 517–537  
*Xiphidiopicus percussus*, 447–451
- Xiphocolaptes albicollis*, 195–206  
*Xolmis cinereus*, 517–537  
*Xolmis dominicana*, 517–537  
Yellow-faced Parrot, 79–88  
Yorio, P., 361–368  
Yungas, 155–158, 615–621  
*Zenaida auriculata*, 517–537  
*Zenaida galapagoensis*, 207–217  
*Zonotrichia capensis*, 15–24, 89–102, 229–246, 517–537

## REVIEWERS FOR ORNITOLOGÍA NEOTROPICAL, VOLUME 22

The following individuals kindly reviewed one or more (\*) manuscripts for Volume 22: L. dos Anjos\*, P. C. Araujo dos Santos, E. Arbeláez, W. Arendt, J. I. Areta\*, V. del R. Avalos, L. Bala, I. Berkunsky\*, C. Bianchi, J. Blake\*, A. Bodrati, J. Brawn, D. Brooks, R. T. Brumfield, T. Brush, S. Bush, D. Calderon-Franco, R. Caparroz, P. Caplonch, J. Chaves-Campos, C. Cattau, G. Cerón, B. Childress, N. Cleere, J. Collazo, P. Corcuera, J. Cornejo, C. Cornelius, M. del Coro Arizmendi, M. P. Dantas Santos, A. Di Giacomo, M. A. dos Santos Alves, C. Duca, J. C. Eitnhear\*, R. M. Erwin, I. Escalante, G. Fernandez, B. Fessl, R. A. Figueroa R., J. Fitzpatrick, R. M. Fraga\*, A. Gantz, S. Gill, A. Giraudo, L. G. Gómez-Bernal, A. Gómez Laich, O. Hamann, B. Hillcoat, D. Hoffmann\*, M. Isler, O. Jahn, F. Jaksic, R. Johnston, L. Jones, T. Katzner, G. M. Kirwan, S. Koenig, N. Krabbe\*, A. Kusch, S. Lambertucci, B. Loiselle, G. Londoño, L. Lopes, J. Luévano, C. G. Machado, R. Macedo, C. Marantz, G. Marateo, P. R. Martin, M. A. Martinez Morales, D. R. Martinez Ponce, D. McDonald, S. McGehee, I. McGregor-Fors,

R. McNeil\*, E. Mellink\*, C. de Melo, H. Mikola, T. A. Miller, D. Montalti, A. Navarro Siguenza\*, V. Ojeda, E. Oliveira, F. Olmos, E. Palacios, U. Pardiñas, T. Z. Pequeño Saco, M. A. Pizo, J. Polop, D. Porter, P. Pulgarin, J. Ragusa-Netto, M. Raposo\*, J. Rappole\*, K. Renton\*, R. Restall, R. Ribon, L. Rivera, W. D. Robinson\*, K. Roesler, E. Ruelas Inzunza\*, V. Ruiz-Gutierrez, T. B. Ryder, K. Rylander\*, A. Salinas-Melgoza, S. Salvador, L. A. Sánchez-González, L. Sandoval, M. Sazima, A. Schmitz\*, K.-L. Schuchmann\*, T. Schulenberg, A. Simeone, A. Solano-Ugalde, A. L. T. de Souza, J. Taylor, J. M. Thiollay\*, S. Thorn, R. Torres-Avilés, C. A. Valle, T. Valqui, M. F. Vasconcelos\*, P. Vergara, F. Villela\*, A.-A. Weller\*, T. H. White, Jr., J. Wiley\*, A. Yanosky, P. Yorio, T. Zaiden, K. Zyskowski.

Some of the abstracts and/or manuscripts were kindly corrected by J. C. Eitnhear, A. Miglo, and J. Rappole for English, by E. Ruelas I. and A. Schmitz for Spanish, and by L. dos Anjos, M. Raposo, and M. F. Vasconcelos for Portuguese. Page-setting was done by A.-A. Weller.

