

CONTENTS

INTRODUCTION	1
METHODS AND MATERIALS	2
DESCRIPTION OF STUDY AREAS	3
RESULTS	10
Breeding Season Censuses	10
Densities	10
Species richness	14
Diversities	15
Behavior	16
Activity patterns	16
Foraging methods	17
Tree species selection	18
Horizontal tree position	19
Perch selection	20
Stance	22
Foliage Utilization	23
Mean height and use range	24
Foliage profiles	24
Cluster analyses	37
Foliage use index	37
Bulge use	39
Territory Size	40
Energy Requirements	40
Body Weight	44
DISCUSSION	44
Community Composition	44
Species numbers and densities	44
Diversities	46
Bird Species Diversity vs. Vegetational Complexity	46
Resource Partitioning and the Niche	47
Composite cluster analyses	48
Species segregation	48
Foliage Utilization	52
Territory Size	53
Energy Flow	55
Species Substitutions	58
Species Dominance	58
SUMMARY	62
ACKNOWLEDGMENTS	63
LITERATURE CITED	63

TABLES

Table 1.	Composition of trees on all study plots	5
Table 2.	Breeding densities of species and foraging and nesting guilds in 1973	10
Table 3.	Breeding densities of species and foraging and nesting guilds in 1974	11
Table 4.	Breeding densities of species and foraging and nesting guilds in 1975	12
Table 5.	Species richness, diversity, and evenness for the bird communities on all study plots	14
Table 6.	Behavioral responses to habitat alteration	15
Table 7.	Activity pattern alterations by four selected bird species	16
Table 8.	Foraging method alterations by four selected bird species	17
Table 9.	Alterations in tree species selection by five bird species	18
Table 10.	Alterations in horizontal tree positions by four bird species	20
Table 11.	Perch selection alterations by five selected bird species	21
Table 12.	Changes in stance by the Pygmy Nuthatch and White-breasted Nuthatch	21
Table 13.	Mean heights and use ranges for seven selected bird species	22
Table 14.	Foliage use indices for nine bird species	37
Table 15.	Mean territory sizes of nine bird species	38
Table 16.	Mean weight, consuming biomass, and existence energy requirements per individual during the breeding season	39
Table 17.	Standing crop biomass and consuming biomass of the breeding birds of the study areas	40
Table 18.	Participation of individual species in energy flow through the bird community in terms of existence energy in 1973	41
Table 19.	Participation of individual species in energy flow through the bird community in terms of existence energy in 1974	42
Table 20.	Participation of individual species in energy flow through the bird community in terms of existence energy in 1975	43
Table 21.	Relationship between mean territory size, use of the bulge, and fit with the foliage profile	54

FIGURES

Figure 1.	Map of Beaver Creek Watershed, Coconino National Forest, Arizona	4
Figure 2.	Control plot	6
Figure 3.	Silviculturally cut plot	6
Figure 4.	Strip cut plot	8
Figure 5.	Severely thinned plot	8
Figure 6.	Clear cut plot	9
Figure 7.	Mountain Chickadee use of the available foliage volume on the silviculturally cut plot	23
Figure 8.	Mountain Chickadee use of the available foliage volume on the control plot ..	23
Figure 9.	White-breasted Nuthatch use of the available foliage volume on the severely thinned plot	25
Figure 10.	White-breasted Nuthatch use of the available foliage volume on the strip cut plot	25
Figure 11.	White-breasted Nuthatch use of the available foliage volume on the silviculturally cut plot	26
Figure 12.	White-breasted Nuthatch use of the available foliage volume on the control plot	26
Figure 13.	Pygmy Nuthatch use of the available foliage volume on the silviculturally cut plot	27
Figure 14.	Pygmy Nuthatch use of the available foliage volume on the control plot	27
Figure 15.	Western Bluebird use of the available foliage volume on the severely thinned plot	28

Figure 16.	Western Bluebird use of the available foliage volume on the strip cut plot	28
Figure 17.	Western Bluebird use of the available foliage volume on the silviculturally cut plot	29
Figure 18.	Solitary Vireo use of the available foliage volume on the severely thinned plot ..	29
Figure 19.	Solitary Vireo use of the available foliage volume on the strip cut plot	30
Figure 20.	Solitary Vireo use of the available foliage volume on the silviculturally cut plot ..	30
Figure 21.	Solitary Vireo use of the available foliage volume on the control plot	31
Figure 22.	Yellow-rumped Warbler use of the available foliage volume on the silviculturally cut plot	31
Figure 23.	Grace's Warbler use of the available foliage volume on the severely thinned plot	32
Figure 24.	Grace's Warbler use of the available foliage volume on the strip cut plot	32
Figure 25.	Grace's Warbler use of the available foliage volume on the silviculturally cut plot	33
Figure 26.	Grace's Warbler use of the available foliage volume on the control plot	33
Figure 27.	Gray-headed Junco use of the available foliage volume on the silviculturally cut plot	34
Figure 28.	Gray-headed Junco use of the available foliage volume on the control plot	34
Figure 29.	Total bird use of the available foliage volume on the severely thinned plot	35
Figure 30.	Total bird use of the available foliage volume on the strip cut plot	35
Figure 31.	Total bird use of the available foliage volume on the silviculturally cut plot	36
Figure 32.	Total bird use of the available foliage volume on the control plot	36
Figure 33.	Dendrogram resulting from cluster analysis of the Euclidean distance matrix of composite behavior for the breeding birds of the severely thinned plot	47
Figure 34.	Dendrogram resulting from cluster analysis of the Euclidean distance matrix of composite behavior for the breeding birds of the strip cut plot	48
Figure 35.	Dendrogram resulting from cluster analysis of the Euclidean distance matrix of composite behavior for the breeding birds of the silviculturally cut plot	49
Figure 36.	Dendrogram resulting from cluster analysis of the Euclidean distance matrix of composite behavior for the breeding birds of the control plot	49
Figure 37.	Relationship between mean territory size, utilized foliage volume, bulge use, and foliage fit	56
Figure 38.	Relation between the number of behavioral changes and overall presence	59
Figure 39.	Relation between the number of behavioral changes and overall standing crop ...	60



Mountain Chickadee (*Parus gambeli*), by Virgil Scott