

CONTENTS

AUTHOR ADDRESSES ix

ABSTRACT 1

INTRODUCTION 2

STUDY AREA 3

METHODS 5

 Aerial Survey Methodology 5

 Transect Location Design 6

 Survey Timing Design 6

 At-sea Sub-areas 7

 Coastal Sub-areas 8

 Spatial Analysis Methods 8

 Statistical Analyses 8

 Comparisons to Past Density Estimates 9

 Distribution Maps 10

RESULTS 10

SPECIES ACCOUNTS 42

 Gaviidae 42

Common Loon 42

Pacific Loon 42

 Western Grebe and Clark’s Grebe 42

 Procellariidae 47

Black-footed Albatross 47

Laysan Albatross 47

Northern Fulmar 47

Sooty Shearwater and Short-tailed Shearwater 50

Pink-footed Shearwater 51

Black-vented Shearwater 54

Leach’s Storm-Petrel 54

Black Storm-Petrel 54

<i>Ashy Storm-Petrel</i>	58
Brown Pelican	58
Cormorants	61
<i>Double-crested Cormorant</i>	61
<i>Brandt's Cormorant</i>	65
<i>Pelagic Cormorant</i>	65
Surf Scoter and White-winged Scoter	68
Scolopacidae	68
<i>Red-necked Phalarope</i>	73
<i>Red Phalarope</i>	73
Laridae	73
<i>Heermann's Gull</i>	73
<i>Bonaparte's Gull</i>	73
<i>California Gull</i>	78
<i>Western Gull</i>	78
<i>Black-legged Kittiwake</i>	82
<i>Sabine's Gull</i>	82
<i>Caspian Tern</i>	82
Alcidae	86
<i>Common Murre</i>	86
<i>Pigeon Guillemot</i>	87
<i>Xantus's Murrelet</i>	87
<i>Cassin's Auklet</i>	89
<i>Rhinoceros Auklet</i>	92
<i>Tufted Puffin</i>	92
DISCUSSION	92
ACKNOWLEDGMENTS	95
LITERATURE CITED	96

TABLES

TABLE 1a. Densities (birds/km² ± SE) of seabirds within at-sea sub-area S1 (north) during January, May, and September from 1999–2002 12

TABLE 1b. Densities (birds/km² ± SE) of seabirds within at-sea sub-area S2 (west-central) during January, May, and September from 1999–2002. 14

TABLE 1c. Densities (birds/km² ± SE) of seabirds within at-sea sub-area S3 (central) during January, May, and September from 1999–2002 16

TABLE 1d. Densities (birds/km² ± SE) of seabirds within at-sea sub-area S4 (south-east) during January, May, and September from 1999–2002..... 18

TABLE 1e. Densities (birds/km² ± SE) of seabirds within at-sea sub-area S5 (south) during January, May, and September from 1999–2002 20

TABLE 2a. Densities (birds/km² ± SE) of seabirds along all coastlines within the study area during January, May, and September from 1999–2002 22

TABLE 2b. Densities (birds/km² ± SE) of seabirds along mainland coastlines within the study area during January, May, and September from 1999–2002..... 24

TABLE 2c. Densities (birds/km² ± SE) of seabirds along island coastlines within the study area during January, May, and September from 1999–2002 26

TABLE 3a. Densities (birds/km² ± SE) of seabirds along the northern mainland coastline during January, May, and September from 1999–2002 28

TABLE 3b. Densities (birds/km² ± SE) of seabirds along the central mainland coastline during January, May, and September from 1999–2002 30

TABLE 3c. Densities (birds/km² ± SE) of seabirds along the southern mainland coastline during January, May, and September from 1999–2002 32

TABLE 4a. Densities (birds/km² ± SE) of seabirds from coastal transects around the northern Channel Islands’ coastlines in the Southern California Bight during January, May, and September from 1999–2002. Northern Channel Islands include San Miguel, Santa Rosa, Santa Cruz, and Anacapa islands..... 34

TABLE 4b. Densities (birds/km² ± SE) of seabirds from coastal transects around the southern Channel Islands’ coastlines in the Southern California Bight during January, May, and September from 1999–2002. Southern Channel Islands include Santa Barbara, San Nicolas, Santa Catalina, and San Clemente islands 36

TABLE 5. Significance tests based on F-statistics from the GLMM model for analyzing season, sub-area, and season-by-sub-area interaction effects on at-sea densities of seabirds by species. All tests were conducted for the range of months and sub-areas having a positive density estimate. Differences among all months (January, May, and September) and all sub-areas (S1 through S5) were tested, unless otherwise noted. Species types with no test for a season, sub-area, or interaction effect did not have sufficient density information to test that effect. Any effect with F-statistic leading to a P < 0.05 is considered to be statistically significant 38

TABLE 6. Significance tests based on F-statistics from the GLMM model for analyzing season, sub-area, and season-by-sub-area interaction effects on coastal densities of birds by species. All tests were conducted for the range of months and sub-areas having a positive density estimate. Differences among all months (January, May, and September) and all sub-areas (NIC = Northern Island Coastline, SIC = Southern Island Coastline, NMC = Northern Mainland Coastline, CMC = Central Mainland Coastline, and SMC = Southern Mainland

Coastline) were tested, unless otherwise noted. Species types with no test for a season, sub-area, or interaction effect did not have sufficient density information to test that effect. Any effect with F-statistic leading to a $P < 0.05$ is considered to be statistically significant..... 39

TABLE 7a. Significance tests based on Wald’s Z-statistics from the GLM model for analyzing differences in at-sea densities of seabirds between 1975–1983 and 1999–2002, by species and sub-area (S1, S2, and all five sub-areas combined). Species with no test for a sub-area did not have sufficient density information to test period differences in that sub-area. A negative Z-statistic indicates densities were greater from 1975–1983. A positive Z-statistic indicates densities were greater from 1999–2002. Any effect with a $P < 0.05$ is considered to be statistically significant..... 40

TABLE 7b. Significance tests based on Wald’s Z-statistics from the GLM model for analyzing differences in at-sea densities of seabirds between 1975–1983 and 1999–2002, by species and sub-area (S3, S4, and S5). Species with no test for a sub-area did not have sufficient density information to test period differences in that sub-area. A negative Z-statistic indicates densities were greater from 1975–1983. A positive Z-statistic indicates densities were greater from 1999–2002. Any effect with a $P < 0.05$ is considered to be statistically significant..... 41

FIGURES

FIGURE 1. Map of central and southern California showing locations of county boundaries, major cities, coastal points, and islands..... 3

FIGURE 2. Map of central and southern California showing oil lease and platform locations and survey lines flown by Briggs et al. (1987). Oil leases are represented by squares. Platforms are represented by solid circles within lease areas. Lines surveyed in 1975–1978 are represented by solid lines. Lines surveyed in 1980–1983 are represented by dotted lines. 4

FIGURE 3. Map of central and southern California showing locations of core area and non-core area transect lines. Core area transect lines are represented by thicker lines. Non-core area transect lines are represented by thinner lines. The core area was surveyed twice each survey month from 1999–2002..... 6

FIGURE 4. Map of central and southern California showing locations of at-sea and coastal subareas. At-sea sub-areas are numbered 1–5. Coastal sub-area boundaries are denoted by bars. NMC = northern mainland coast. CMC = central mainland coast. SMC = southern mainland coast..... 7

FIGURE 5. All seabird densities (birds/km²) and distribution off southern California from 1999–2002 during January, May, and September..... 11

FIGURE 6. Loon densities (birds/km²) and distribution off southern California from 1999–2002 during January, May, and September 43

FIGURE 7. Unidentified loon densities (birds/km²) and distribution off southern California from 1999–2002 during January, May, and September..... 44

FIGURE 8. Common Loon densities (birds/km²) and distribution off southern California from 1999–2002 during January and September..... 45

FIGURE 9. Pacific Loon densities (birds/km²) and distribution off southern California from 1999–2002 during January and May 46

FIGURE 10. Western Grebe densities (birds/km²) and distribution off southern California from 1999–2002 during January, May, and September..... 48

FIGURE 11. Procellariid densities (birds/km ²) and distribution off southern California from 1999–2002 during January, May, and September	49
FIGURE 12. Black-footed Albatross sightings off southern California during January and May of 2000 and January and September of 2001	50
FIGURE 13. Laysan Albatross sightings off southern California during January of 2000 and 2001.....	51
FIGURE 14. Northern Fulmar densities (birds/km ²) and distribution off southern California from 1999–2002 during January, May, and September	52
FIGURE 15. Sooty Shearwater densities (birds/km ²) and distribution off southern California from 1999–2002 during January, May, and September	53
FIGURE 16. Pink-footed Shearwater densities (birds/km ²) and distribution off southern California from 1999–2002 during January, May, and September	55
FIGURE 17. Black-vented Shearwater densities (birds/km ²) and distribution off southern California from 1999–2002 during January, May, and September	56
FIGURE 18. Leach’s Storm-Petrel densities (birds/km ²) and distribution off southern California from 1999–2002 during January, May, and September	57
FIGURE 19. Black Storm-Petrel densities (birds/km ²) and distribution off southern California from 1999–2002 during January, May, and September	59
FIGURE 20. Ashy Storm-Petrel densities (birds/km ²) and distribution off southern California from 1999–2002 during January, May, and September	60
FIGURE 21. Brown Pelican densities (birds/km ²) and distribution off southern California from 1999–2002 during January, May, and September	62
FIGURE 22. Cormorant densities (birds/km ²) and distribution off southern California from 1999–2002 during January, May, and September	63
FIGURE 23. Unidentified cormorant densities (birds/km ²) and distribution off southern California from 1999–2002 during January, May, and September	64
FIGURE 24. Double-crested Cormorant densities (birds/km ²) and distribution off southern California from 1999–2002 during January, May, and September	66
FIGURE 25. Brandt’s Cormorant densities (birds/km ²) and distribution off southern California from 1999–2002 during January, May, and September	67
FIGURE 26. Pelagic Cormorant densities (birds/km ²) and distribution off southern California from 1999–2002 during January, May, and September	69
FIGURE 27. Surf Scoter densities (birds/km ²) and distribution off southern California from 1999–2002 during January, May, and September	70
FIGURE 28. Phalarope densities (birds/km ²) and distribution off southern California from 1999–2002 during January, May, and September	71
FIGURE 29. Unidentified phalarope densities (birds/km ²) and distribution off southern California from 1999–2002 during January, May, and September	72
FIGURE 30. Red-necked Phalarope densities (birds/km ²) and distribution off southern California from 1999–2002 during January, May, and September	74
FIGURE 31. Red Phalarope densities (birds/km ²) and distribution off southern California from 1999–2002 during January, May, and September	75
FIGURE 32. Larid densities (birds/km ²) and distribution off southern California from 1999–2002 during January, May, and September	76

FIGURE 33. Heermann's Gull densities (birds/km ²) and distribution off southern California from 1999–2002 during January, May, and September	77
FIGURE 34. Bonaparte's Gull densities (birds/km ²) and distribution off southern California from 1999–2002 during January and May	79
FIGURE 35. California Gull densities (birds/km ²) and distribution off southern California from 1999–2002 during January, May, and September	80
FIGURE 36. Western Gull densities (birds/km ²) and distribution off southern California from 1999–2002 during January, May, and September	81
FIGURE 37. Black-legged Kittiwake densities (birds/km ²) and distribution off southern California from 1999–2002 during January and May	83
FIGURE 38. Sabine's Gull densities (birds/km ²) and distribution off southern California from 1999–2002 during May and September	84
FIGURE 39. Caspian Tern densities (birds/km ²) and distribution off southern California from 1999–2002 during January, May, and September	85
FIGURE 40. Alcid densities (birds/km ²) and distribution off southern California from 1999–2002 during January and May	86
FIGURE 41. Common Murre densities (birds/km ²) and distribution off southern California from 1999–2002 during January and May	88
FIGURE 42. Pigeon Guillemot densities (birds/km ²) and distribution off southern California from 1999–2002 during January and May	89
FIGURE 43. Xantus's Murrelet densities (birds/km ²) and distribution off southern California from 1999–2002 during January and May	90
FIGURE 44. Cassin's Auklet densities (birds/km ²) and distribution off southern California from 1999–2002 during January and May	91
FIGURE 45. Rhinoceros Auklet densities (birds/km ²) and distribution off southern California from 1999–2002 during January and May	93