



Mated ♂ Savannah and ♀ Ipswich Sparrows Martinique Beach, Halifax Co., N.S. Photo/I. A. McLaren.

sight records. Extralimital Seaside Sparrows were carefully observed at Hampton, N.H., Sept. 27 (PDeB et al.), and at Brier I., N.S., Oct. 9 (ELM). About 25 Lark Sparrows were reported, only 6 of them north of Massachusetts. Of the 9 Clay-colored Sparrows seen, the more unusual were single birds at Seabrook, N.H., Sept. 18 (DJA, DWF, photographed), one of very few New Hampshire records, and at Harkness Park, New London, Conn., Oct. 21 (GB, fide RCD), one of very few Connecticut records. This fall saw a particularly heavy flight of White-crowned Sparrows, noted at P.E.I. and in New Brunswick, New Hampshire and Massachusetts. The peak in New Brunswick was the first week of October, when the birds were said to be "all over the place" (DSC); in New Hampshire on Oct. 3, 100+ were counted in the Lakes Region (VHH, RCH) and 53 at Littleton (RB), and of many reports in Massachusetts, the high count was 40 at Corn Hill in Truro, Cape Cod, Oct. 11 (CAG, WRP et al.). Finally, an individual with the characteristics of the western race *gambelii* was photographed at Seal I., N.S., Oct. 12 (IAMcL).

SUB-REGIONAL EDITORS, CONTRIBUTORS (both in boldface), OBSERVERS AND OTHER ABBREVIATIONS — **Dennis J. Abbott III, Kenneth P. Able, Peter Alden, Charles R. K. Allen, H. Cook Anderson, Kathleen S. Anderson, R. Ross Anderson, Arthur Argue, Margaret Argue, Wallace Bailey, James Baird, Larry S. Benjamin, Thomas Bickel, Grace Bissell, Robert Bradley, Dorothy W. Briggs, David T. Brown, Lewis A. Buck, Harold M. Card, James E. Cavanagh, Edward L. Chalif, William H. Chisholm, David S. Christie, Don Clark, Nancy Claybourne, Richard Claybourne, Robert V. Clem, Eric H. Cooke, Hamilton Coolidge, Davis H. Crompton, Louise D. Daley, Severyn S. Dana, George G. Daniels, Paul DeBenedictis, Paul J. Desjardins, Robert C. Dewire, Benjamin K. Doane, Phyllis R. Dobson, Paul K. Donahue, R. Kenneth Edwards, William S. Emerson, Ruth P. Emery, Richard L. Fern, Davis W. Finch, Richard A. Forster, Sylvia J.**

**Fullerton, Constance D. Gallagher, J. Roswell Gallagher, W. Earl Godfrey, Carl A. Goodrich III, Barbara Grace, Patrick W. Grace, Madeleine Grumbach, Richard C. Hebert, Vera H. Hebert, Oliver H. Hewitt, Sibley Higginbotham, Deborah V. Howard, Richard R. Howie, Sarah P. Ingalls, H. Lawrence Jodrey, Cecil L. Johnson, Alice D. Jones, Vincent W. Kaselis, Allan R. Keith, Hugh E. Kingery, Everett Knapp, Douglas L. Kraus, Adelma E. Lajoie, Christopher W. Leahy, Jean-Paul Lebel, Paul Lehman, H. Willa MacCoubrey, Sara MacLean, Mary Majka, Mabel W. McIntosh, Ian A. McLaren, Paul S. Miliotis, Hazel Miller, Eric L. Mills, Nettie Moore, Ian C. T. Nisbet, Helen W. Parker, Philip Parsons, Peter A. Pearce, Wayne R. Petersen, Elizabeth W. Phinney, Leon G. Phinney, Richard L. Plunkett, Roger Pocklington, William C. Russell, Richard M. Sargent, Eloise A. Saunders, Robert Scully (RSe), Robert W. Smart, Sidney Smith, Gerald L. Soucy, Wayne T. Stobo, Robert Stymeist (RS), Helen Tetrault, Edward V. Thompson, William C. Townsend, Dallas Wait, Arthur Wang, Richard Webster, Berna B. Weissman, Daniel A. Welsh, James G. Wilson, Charles Wood, John C. Yrizarry, Joseph Zeranski; Audubon Society of New Hampshire, Manomet (Mass.) Bird Observatory, Saint John (N.B.) Naturalists' Club, University of New Hampshire, v.o., various observers — DAVIS W. FINCH, Knoll Creek Farm, Route 376, Hopewell Junction, N.Y. 12533.**

#### HUDSON-ST. LAWRENCE REGION /Ned R. Boyajian

It was another notably mild fall, with southern areas experiencing considerable heavy rains but northern areas remaining fairly dry until the first heavy snows in mid-November. Hurricane Doria, which brushed the coast in late August, produced some interesting results, and there were a few good flights. But owing

to infrequent cold fronts, many locations, especially along the coast, reported an uneventful season, at least in terms of marked flights. On the other hand, the mild season resulted in a great number of late lingerers. The most outstanding are shown in Table I. Note especially the number of vireos.



**DORIA** — The storm produced 2 **Sooty Terns** at Great Gull I. (RP) and 1 each at Riis Park, and Shinnecock, L.I., and Manasquan Inlet, N.J. (fide, PB). It was probably responsible also for a total of 4-5 **Leach's Petrels** off Long I. from Great Gull to Riis Park (RP, L.S.N.Y. et al.). Although the Leach's is doubtless a regular migrant offshore, the comparatively few Regional records have mostly followed major storms. Although perhaps not a result of Doria, southern landbirds seemed notably more frequent than is usual in Fall, among them Prothonotary Warbler, 1, Great Gull I., Sept. 8 (RP) and 1, Staten I., Sept. 28 (HF); there are only about 15 previous fall records for the New York City area; Yellow-throated Warbler, 1, Brooklyn, Sept. 4 (ES); only 4-5 previous fall records for the area; Summer Tanager, 1, Tobay Beach, L.I., Sept. 9 (AW); Blue Grosbeak, 9 for the season on Long I. and Staten I., the last one the late date of Oct. 29 at Upper Brookville L.I. (BC). Naturally, with breeding species it cannot be determined whether a given bird is a stray or a migrant. For example, a total of 8 Worm-eating Warblers on Long I. in early September seemed more than might normally be expected there.

**MAJOR FLIGHTS** — A marked early movement of landbirds was noted in coastal Westchester Co., N.Y. Aug. 22, with vireos and warblers predominant (RA). There was a flight on Oct. 7-9, apparently manifest only in the south, mostly on the coast, and less noticeable inland, where, as is often the case, it only became evident a day or two after hitting the coast. Notable sightings included a flicker with red shafting at New Haven (NP), 25 Downy Woodpeckers at Far Rockaway, L. I. (JB, DF & HH), 12,000 Com. Scoters (DF)

150 Brown Creepers (research station) and 35 White-crowned Sparrows (DF) at Fire I. L.I., and in 3 hours at Alpine, N.J., 64 Sparrow Hawks and 462 Am. Goldfinches (NB). A second on Oct. 16-17 was Region-wide, and notable for widespread counts of 100 to 2000-3000 Canada Geese and a good movement of "coastal" species in the Hudson Valley. For example, season highs of Double-crested Cormorant, scoters and Red-breasted Mergansers and a Caspian Tern at Alpine (NB) and very carefully identified at Poughkeepsie, the second Dutchess Co. record of **Black-legged Kittiwake** (JK & MK, fide EP & OW). Other finds included, on Oct. 16, 42 late Broad-winged Hawks at N. Adams, Mass. (BBE); Oct. 17, a Black-billed Magpie at W. Orange, N.J. (RR) and 25 White-crowned Sparrows at Baxter Creek, Bronx Co., N.Y. (RK). A third flight developed on Oct. 30-31. This was an unusual period, highlighted by a widespread flight of Snow Geese, two species in Quebec well beyond their normal ranges and on extraordinarily late dates, and some dislocation of coastal species inland. At Alpine, N.J., Snow Geese, mostly in high-flying flocks of 50-200, totaled 704 birds (NB), other flocks were reported from locations from Winhall, Vt. to Long I. On Oct. 31 a **White-eyed Vireo** was very carefully identified at Deschenes, Que. (DB & JM, fide RF). Its appearance was attributed to several successive days of s.w. winds, a phenomenon which may also explain the presence on Nov. 6 of a **Rough-winged Swallow** at Beauharnois (JS & TT, fide MM). Odd inland occurrences included 2 Red-throated Loons, 242 Brant and 3 Dunlin at Alpine (NB), 37 Gadwall and a Laughing Gull at Point View Res., N.J. (RRo). Other notable finds were 37 Hooded Mergansers at Rye, N.Y. (GD) and 35 Old Squaws and a Lark Bunting at Baxter Creek (RK, DR & NS).

**OTHER SEASONAL TRENDS** — Certain Raptors were more numerous than usual, but as is always the case, it is difficult to say to what extent this might have been owing to favorable circumstances of observation and to what extent real increases. For instance, counts from Raccoon Ridge (HD et al.), N.J. Nov. 4 of 1022 Red-tailed Hawks, about twice the previous New York City area record, followed by 500 each Nov. 5 & 6, was, in the observers' opinion, owing entirely to the virtual absence of the species through October. Similar conditions may in part account for some of the following. **Golden Eagle**. No less than 38 fully satisfactory reports. Even with the widest allowance for duplication, e.g., a bird seen in Quebec might conceivably be the same as one reported southward a few days later, the total is still at least 20-25. Furthermore, several were from locations, e.g., inner coastal plain of N.J., or the s. Bronx, where the species is casual. **Osprey**. Many, both inland and coastally. Some daily counts were: 80, Hudson R. Valley, Dutchess Co. N.Y., to Alpine, N.J., Sept. 19 (R.T.W.B.C., F.N.A.). NB doubtless with much duplication, plus 17 more to the westward at Mt. Peter, N.Y. (H.A.S.); 69 at East Haven, Conn., Sept. 30, and 62 on Oct. 3 (NC). Some season totals were: Hook Mt., N.Y., 180 (F.N.A.); Alpine, 106 (NB); Watertown, Conn., 61 (NC) and Pittsfield, Mass., 39 (LD & RG). **Peregrine Falcon**. Total of 65, with many from locales where

unusually recorded, e.g., Topsham, Vt. Oct. 2 (A & EM).

Snowy Owls appeared as expected, reaching the Montreal area by mid-October, where, by season's end, 10 had been sighted. Southward, there was 1 at Poughkeepsie, N.Y., Nov. 1 (MK) and 1 in the Bronx Nov. 7 (L.S.N.Y.) with the first widespread flight around Nov. 23, bringing about 12 by the end of the season.

Winter finches were more in evidence than last year, although to date not attaining the magnitude of 1969. A summary will appear in a later report.

**LOON, GREBE, CORMORANTS, EGRET, IBIS** — Red-throated Loons seemed back to normal after last year's influx, although 1 at Mississquoi Bay, Nov. 14, was rather unusual (JS, fide MM). The annual Eared Grebe at Jamaica Bay Wildlife Refuge, L.I., (J.B.W.R.) was seen Oct. 22 but apparently did not linger (DF). A Great Cormorant at Great Gull I., Sept. 10, was quite early for the New York City area (N.Y.C.A.) (RP). One was also seen Oct. 9 at Cote Ste. Catherine, Que., where it is always unusual (MG; ND, fide MM). An imm. Double-crested Cormorant at Onondaga L., July 5-9 and an ad. at Fox Ridge, N.Y. July 25, were the first midsummer records for the Oneida L. Basin in recent years (fide MR & CS). A Com. Egret was seen at Hyde Park, N.Y., where unusual, Aug. 19 (DF) and there was 1 at Milford Pt. as late as Nov. 24 (CH). Two Glossy Ibises were seen at Stratford, Conn., Sept. 18 (NP) and 1 lingered at J. B. W. R. until Nov. 21 (DPo).

**WATERFOWL** — Whistling Swans were even more numerous than last year. High was 19 at Sagaponack Pond, L.I., Nov. 2 (fide PB) and groups of 2-10 were seen in most coastal, and scattered inland, locales, including 1, Oct. 24-31 at Deschenes, Que., where it is quite unusual (BM). On Nov. 24, 35 Brant were seen at S. Windsor, Conn. (PD) and on Nov. 9, one at Pittsfield, Mass. (RG) both unusual finds for the locations. Snow Geese were widespread and numerous all season, including 1 as early as Oct. 7 at Far Rockaway (JB et al.). A Blue Goose, always a good find along the coast, was seen at Woodbridge, Conn., Oct. 28 (NP). Gadwall and Shoveler continued to increase. The former was recorded for the first time at Gale Meadows, Vt., where 1 was seen Nov. 2 (WN). Elsewhere, 30 on Dec. 4 at C. Ste. Catherine, where some had been present since Oct. 30, was a good local count (MM). Peaks of 60 Shovelers at N. Arlington on Sept. 20 (RK) and 153 at N. Bergen, N.J., Oct. 28 (NB) were also high. Successful breeding by many free-flying pairs just across the Hudson R. in Bronx Park this past summer (fide, RR) may account for these. European Widgeons were about normal; a total of 6 were reported from s. Conn. to coastal N.J. At Onota L., Pittsfield, Mass., 102 Wood Ducks were seen in early October, a good local count (RG). A Redhead, always a rarity in that area, was also seen there Oct. 14 (RG). On Oct. 15, 50 Oldsquaw and 12 Surf Scoters were found on Plymouth L., Vt.; both new species for the Ottauquechee Valley (RFa). A **Harlequin Duck** at Hull, Que., Nov. 20 was a new species for that area (RPo & RF); another at Fire I., Sept. 25 (MCa) was an early record for N.Y.C.A.

by more than a month. Both eiders were abundant. Two Commons were seen as early as Oct. 10 at Shinnecock (CM, fide PB) and a few scattered Kings arrived by late September. By late November, flocks of 20-60 Commons, with a high count of 200 at Ditch Plains, L.I., Nov. 13 (PB) and 2-10 Kings were frequent on e. Long I., with lesser numbers westward. The Ruddy Duck, always rare in Quebec, was seen twice: 1, Deschenes, Nov. 7 (RF & RPi) and 1, Delson, Nov. 21 (MM & TT).

**RAPTORS** — About 20 Goshawks were reported, scattered from Quebec to n. N.J. and n.w. Pa. The majority of Broad-winged Hawks passed through in three days. *Sept. 19:* Mt. Peter, 2213 (H.A.S.); two locations in Berkshire Co., Mass., 1725 and 2000 with no duplication (RG); two locations, Dutchess Co., N.Y., 2000 and 6000 (R.T.W.B.C.). *Sept. 20:* Hook Mt. 1445, (F.N.A.). *Sept. 21:* Mt. Peter, 6658 (H.A.S.) Watertown, N.Y., 10,917 (NC et al.). An ad. ♂ **Harris' Hawk**, apparently the first for the Northeast, was discovered at Rye Ridge, Westchester Co., Nov. 7 (GK, fide RA) seen several times thereafter, to late Dec. The bird showed no traces of recent captivity (feather wear, jesses, etc.), but captive Harris' Hawks are presently held in the Region. While the record must remain hypothetical, there are numerous other occurrences of weaker-flying s.w. species in fall in the Region (Sage Thrasher, Green-tailed Towhee, Black-throated Sparrow, etc.) which no one has ever suspected of being escapes. The preceding period had been marked by strong s.w. winds along a front from Texas to New England; local temperatures were in the 60s. A remarkable find was a dark phase **Swainson's Hawk**, very carefully identified at New Vernon, N.J., Oct. 23 by an experienced observer familiar with the species (AK). There are 2 previous records for the N.Y.C.A. About 19 Bald Eagles were reported, 50 per cent immatures. Despite the many **Gyrfalcons** in the Northeast this Fall, only 1 was sighted in the Region, a white phase bird well observed at New Haven, Oct. 19 (NP). At Winhall, where it is rare at any time, a Pigeon Hawk was seen on the late date of Nov. 7 (WN).

**PHEASANT, RAIL, SHOREBIRDS** — The previously reported decrease of Ring-necked Pheasants in c. N.Y. since the cessation of stocking was noted also in the Schenectady area (RY). A King Rail at Mill Neck, L.I., Nov. 10 (EF) was a good find for the location. A count of 45 Am. Golden Plovers at S. Windsor, Conn., Oct. 4 was good for the area (PD) as were 80 at N. Arlington, N.J., Sept. 25 (RK). At Suffolk Co. Airport, L.I., 21 Upland Plovers in mid-July (E & LS) may have been early migrants but hopefully they indicated a successful local breeding season. A **Willet** at Onondaga L., July 5-6, was the eighth record for the Oneida L. Basin (CS). A **Knot** at N. Arlington, Aug. 22 (RK) was apparently the first inland record for the N.Y.C.A. Baird's and Buff-breasted Sandpipers were unusually widespread. About 20 of each were reported from Quebec to n. N.J., many from unusual locales. For example, Baird's, 2 at Pittsfield Aug. 29-30, 1 lingering until Sept. 20 (RG; BH); 1, S. Windsor, Sept. 5 (PD et al.); 3 Madison, Conn., Oct. 3 (NP). Buff-breasted, 3, So. Windsor, Aug. 29-Sept. 5 (PD & MF) and N. Arlington, Sept. 6 (RK). The

Sanderling, very rare away from the coast, was reported inland at N. Arlington, a total of 3, Sept. 6-12 (RK) and more unusually, at Martin's Creek, Pa., where there were 3 on July 30 (DE). Quite unusual for Berkshire Co. was a N. Phalarope at N. Adams, Sept. 14-15 (BBE). This species was also recorded at Ile du Moine, 6 on Aug. 21 (MM) and twice in coastal Conn. in late September (JG; NP et al.). Even more unusual for the latter area were single Red Phalaropes at New Haven, Sept. 18-20 (NP et al.) and Litchfield, Oct. 9 (MH, GL et al.). At J.B.W.R., a Ruff was seen Sept. 6 (RPI, fide PB).

**JAEGERS, GULLS, TERNS, ALCIDS** — Jaegers were much in evidence along the Long I. coast from late August through mid-October. High for Pomarine was 17 at Cox's Ledge, Sept. 11 (L.S.N.Y.) and for Parasitic, 35 on Oct. at Jones Beach (DF). A **Lesser Black-backed Gull** which had been frequenting the Ontario shore obligingly appeared on the Quebec side of the Ottawa R. at Wynchwood, Nov. 28 to mark the first provincial sight record (DB, fide RF). Another was seen at Southampton, L.I. on the early date of Sept. 25 (MC). There were 2, possibly 3, "Thayer" type gulls at Beauharnois, Nov. 21-27 (MM). Fifteen Ring-billed Gulls at Onota L., Oct. 18, was notably more than the usual 2-3 per season recorded in Berkshire Co. (RG). A total of about 15 Black-headed and 10 Little Gulls were reported, mostly from their usual haunts, i.e., New Haven Harbor, Lower Hudson-Upper New York Bay, etc. A **Sabine's Gull**, always a great rarity in the Region, was seen at Southampton, L.I., Oct. 30 (CM, fide PB). At Shinnecock Inlet, 20 Forster's Terns were seen on the rather late date of Nov. 20 (PB). They were reported to be scarce on the Conn. coast, where 3 were seen at New Haven, Sept. 20, along with a good count of 6000 Com. Terns (DF). Several of the latter lingered into November, including 1 on Nov. 24 at Milford Pt. (CH). Least Terns were present in the N. Arlington area from late July to late August, with a high of 80 on July 27 (RK). Though sometimes cited as very unusual inland, this bird is actually an annual migrant through the Overpeck-Hackensack N.J., marsh system. Alcids appeared inshore on e. Long I. frequently in November. For example, 20-25 Dovekies and a Thick-billed Murre were seen at Shinnecock Inlet Nov. 20 (E & LS), several sightings of Razorbill, mostly at Montauk (fide GT) and rarest of all, **Common Puffin**, present at Fire I. the last week of the month (BD, fide PB). West and southward, sightings decreased, although there was a scattering of Dovekies and 1-2 Razorbills down to the s. end of the Region.

**DOVE, PARAKEET, OWL, HUMMINGBIRD, WOODPECKERS** — Evidence of the late season were flocks of 60-100 Mourning Doves in the Montreal area into late October (MM). The Monk Parakeet nests at Lordship, Conn. were deserted but new construction was noted at the sites in Valley Stream, L.I., Rockaway Beach and Staten I. (HC; DF; et al.). New sites were reported from Yonkers, N.Y., (DPI, fide RR) and Edgewater, N.J. (NB & WD) and single birds from many southern locations. Banders at Fire I. captured 23 Saw-whet Owls Nov. 1-24 (DFo et al., fide RPa). On Sept. 18, 31 Ruby-throated Hummingbirds

were banded at Vischer Ferry, N.Y. (RY). Migrant Red-headed Woodpeckers seemed scarce. Only 2 were seen at Alpine where the season total is usually 5-6 (NB) and only a scattered few noted elsewhere. Three-toed woodpeckers were much in evidence. During the summer a pair of Black-backed with 3 young (BBu) and a single (FS) were found in Herkimer Co., N.Y. From mid-October on there were 8 in the Montreal area (MM) and 5-6 others in the north. A single **North-ern Three-toed Woodpecker** was seen at Five Ponds, N.Y., July 18, the fourth record for the oneida L. Basin (FS, fide MR). There was also 1 at Montreal, Oct. 10 (WF, fide MM) and 1 at Lucerne, Que., Oct. 30 (RF et al.).

**SWALLOWS TO SHRIKES** — Good counts of swallows included 9000 Barns in 5 hours at Gilgo, L.I., Aug. 12 (RPa) and a remarkable concentration of 490 Purple Martins at E. Brunswick, N.J. Sept. 12 following torrential rains (GC). A Com. Raven, very rare in the N.Y.C.A., especially away from the ridges, was seen at Baxter Creek, Nov. 6 (DR, fide PB). In addition to the N.J. Black-billed Magpie, 1 at Tobay, Sept. 25 and 1 at Babylon, L.I., Oct. 6, were deemed by the observer (PB) to be legitimately wild. Another at Tobay, Oct. 14-16 (mob) may have been one of the above, but perhaps was a third. A Gray Jay, always rare in the Oneida L. Basin, was seen in the Five Ponds, N.Y. area, July 11 & 18. (FS). In the north, chickadees were on the move. New bandings in the Ottawaueechee Valley were ten times last years totals with the same net-hours, reaching 50 a week at peak. (O.V.E.A.C.). In the south, the influx was not so evident, e.g., 10-15 per morning at Alpine vs. 6-8 normally (NB). Boreal were noted Chickadees daily at O.V.E.A.C. and frequently at Winhall (WN) from Oct. 15 through Nov. 8. Southward, there was 1 on Staten I., Nov. 5 (E & LM) and 1 at Brookhaven, Nov. 8 (DP). Red-breasted Nuthatches, very scarce last year, began moving in mid-summer and by end of season had been seen in numbers in most areas. Golden-crowned Kinglets and Cedar Waxwings were also up from last year, the former appearing in record numbers in some areas. For example, 400 at Fire I., Oct. 7 (TD); season average of 15 per hour at Alpine, including 67 in 1½ hrs., Oct. 17 (NB). Cedar Waxwings averaged from 1-20 a day at Topsham (AE&M) to 100-160 a day at Alpine (NB) with counts of 537 at New Haven, Sept. 25 (NP) and 200 in Berkshire Co., early November (RG). Two Loggerhead Shrikes, rarities in Berkshire Co., were at Pittsfield, Aug. 23-24 (GS).

**WARBLERS, ICTERIDS, FINCHES** — There was an Orange-crowned Warblers on the rather early date of Sept. 22 at Milford Pt. (JG) and a rather high total of 11 others from various locations mostly in October. Quite remarkable were 2 reports of **Audubon's Warbler**, a bird unknown in the Region prior to 1969. One was banded and photographed at Tobay, Nov. 6 (AL) and another, different bird, photographed at Cedar Beach, L.I., Nov. 21 (SS, AW & CW). A Mourning Warbler, very rare in fall on e. Long I., was picked up alive at Hampton Bays, Sept. 19 (KM & E & LS). A flight of 23,000 blackbirds, 65 per cent Com. Grackle, was seen at Alpine, Nov. 6 (NB & RR). A Yellow-headed Blackbird, an annual species

**TABLE I**  
**Notable Late Dates, 1971**

Species	Date	Locality	Obs.
Little Blue Heron	11/16	L.I.	PB
Am. Woodcock	12/3	sw. Que.	DL
Roseate Tern	11/2 *	L.I.	PB
Yellow-billed Cuckoo	11/3	n. N.J.	RK
Ruby-thr. Hummingbird	9/26	c. Vt.	A&EM
	10/12	c. N.Y.	RY
Gt. Crested Flycatcher	11/4	L.I.	DF
Rough-winged Swallow	9/18 *	n. N.J.	NB
White-eyed Vireo	10/31	L.I.	DF
Yellow-throated Vireo	11/3 *	L.I.	MS
	11/12*	L.I.	BB
Solitary Vireo	11/6	c. N.Y.	FG&MV
	11/14	L.I.	AL
Philadelphia Vireo	10/16	n. N.J.	NB&RR
Blue-winged Warbler	11/2 *	L.I.	ED
	11/7 *	L.I.	ED
Warbling Vireo	10/2	c. N.Y.	EB
Blue-winged Warbler	10/3	c. N.Y.	EB
Tennessee Warbler	11/4	L.I.	DFO
Nashville Warbler	11/18	L.I.	RPa
Cape May Warbler	11/6 *	c. N.Y.	MV
Black-thr. Green Warbler	11/14	L.I.	ML
Blackburnian Warbler	10/31*	L.I.	G&MK
Chestnut-sided Warbler	10/17	n. N.J.	NB
Pine Warbler	9/18	c. N.Y.	JK&MK
Ovenbird	10/2	c. N.Y.	FG
N. Waterthrush	10/23	s. N.Y.	RK
Connecticut Warbler	10/9	s. Vt.	WN
Hooded Warbler	9/23	c. N.Y.	CT
Am. Redstart	10/30	n. N.J.	GT
	11/12	c. N.Y.	CT
	11/25*	c. N.Y.	CT
Bobolink	11/14	c. N.Y.	T&WS
Red-winged Blackbird, 10	11/12*	c. Vt.	A&EM
Rusty Blackbird	11/29*	c. Vt.	A&EM
Scarlet Tanager	10/22	c. N.Y.	MK
Indigo Bunting	11/4	L.I.	DF
	11/7	n. N.J.	NB&RR
Field Sparrow	11/1	sw. Que.	RF
Lincoln's Sparrow	11/3	n. N.J.	RK

All sightings are single individuals unless otherwise indicated. \*denotes - exceeds previous local late date for migrant.

in recent years, was seen at Fire I., Sept. 25 (TD & HHo). Dickcissels had one of their better years. Most areas reported 2 or 3, highs were 9 at New Haven, Sept. 25 (NP) and 8 at Fire I., Oct. 8 (DF). At Riis Park, an Ipswich Sparrow was found on the early date of Oct. 6 (DF). About 5-6 Lark Sparrows were reported on the coast, and more unusually, 1 inland at E. Brunswick, Aug. 28 (GC). A **Green-tailed Towhee** at the Haeni feeder, Scarsdale, Oct. 19 and subsequently (RA, et al.) was the third N.Y.C.A. record, first for New York State. A imm. Slate-colored Junco was seen at Litchfield, Conn., where it is not known to breed, on the early date of Aug. 25 (MH). There has been a slight increase in nesting in other southern areas recently. At N. Arlington, a Fox Sparrow was seen Oct. 11 (RK) one of the earliest dates for the N.Y.C.A.

**CONTRIBUTORS** — Robert Arbib, B. Baumann, (BBe) Bruce Beehler, Ned Boyajian, Dan Brunton, Paul Buckley, Morgan Bulkeley, John Bull, (BBU) B.

Burt, Enid Butler, (MCA) Matthews Cashman, Howard Cleaves, Robert Clermont, Gary Cohen, Mark Collie, Barbara Conolly, Neil Currie, Ed Daley, Gus Daniels, Harry Darrow, Normand David, Tom Davis, Paul Desjardins, Robert Dewire, William Downin, Laura Drummond, Bob Dunnigan, David Edwards, (RFa) Richard Farrar, Davis Finch, Howard Fischer, (DFo) Darrel Ford, Roger Foxall, Mary French, W. Frey, Elsie Fruson, Fyke Nature Association, J. Roswell Gallagher, Florence Germond, Richard Goodrich, Michele Gosselin, Michael Harwood, Bartlett Hendricks, Highlands Audubon Society, Charles Hills, Helen Hirschbein, (HHo) Howard Honig, Alice Jones, (RK) Richard Kane, George Kappus, R. Kearney, Allan Keith, Jim Key, Mary Key, Georges and Michel Kleinbaum, Don LaFontaine, Anthony Lauro, Paul Lehman, Milliard Lindauer, Linnaean Society of New York, Gordon Loery, Alison and Elizabeth Macdonald, B. MacTavish, J. McCraig, Mabel McIntosh, Christopher McKeever, Edward and Laura Mills, Karl Mueller, William Norse, Ottauqueechee Valley Environmental Analysis Center, Roger Pasquier, (RPa) Robert Paxton, Elinor Pink, (RPi) Ron Pittaway, (RPl) Richard Plunkett, (RPo) Richard Poulin, (DPo) Don Powers, Noble Proctor, Dennis Puleston, Dan Rafferty, Ralph T. Waterman Bird Club, Richard Rogers, (R Ro) Richard Rosche, Marge Rusk, (RR) Richard Ryan, Eric and Lorna Salzman, Fritz Scheider, Seymour Schiff, Neil Schoenfeld, Gary Shampang, Martin Sohmer, Chris Spies, J. Steeves, Timothy Sterrett, Trixie and William Strauss, Esther Swayer, Czecher Terhune, Stiles Thomas, Terry Thormin, Greg Toffic, (GTu) Guy Tudor, Marion Van Wagner, Cornelius Ward, Otis Waterman, Alvin Wollin, Robert Yunic, Guy Zern, — many observers (mob), NED. R. BOYAJIAN, 9 Reading Terrace, Fair Lawn, N.J. 07410.

**MIDDLE ATLANTIC COAST REGION**  
**/F. R. Scott and David A. Cutler**

The fall weather featured a September and October that were far warmer than usual, with Philadelphia, Pa. having deviations from normal for these months of +4.9° and +7.8°F., respectively. Rainfall was also excessive. Two hurricanes affected the Region rather severely. *Doria* passed northward along the coast dur-

