

- American Naturalist, Nov., Dec., 1884, Jan., 1885.
 Bulletin Essex Institute, XV, Nos. 7-9, XVI, Nos. 4-6.
 Canadian Science Monthly, Sept., 1883.
 Journal Cincinnati Soc. Nat. Hist., VII, Nos. 1-3.
 Naturalist, The, A Journ. of Nat. Hist. for the North of England, Nos. 112-113, Nov., Dec., 1884.
 Ornithologist and Oölogist, Aug.-Dec., 1884.
 Proceedings Acad. Nat. Sci. Philadelphia, 1884, Pt. II.
 Random Notes on Natural History, I, Nos. 10-12, 1884.
 Science Record, II, Nos. 11, 12.
 Zeitschrift für die gesammte Ornithologie, I, Heft 3.
 Zoölogist. Oct., Nov., Dec., 1884.

GENERAL NOTES.

Albino Robins (*Turdus migratorius*).—My friend James M. LeMoine, Esq., informs me that he has just added to his fine collection at Spencer Grange, Quebec, a pure white Robin, said to be the first seen in that vicinity. I have lately examined an albino of this species recently captured near St. John. The only colored feathers to be seen on the bird are three single ones forming spots on the breast.—MONTAGUE CHAMBERLAIN, *St. John, N. B.*

Nest and Eggs of the Golden-winged Warbler (*Helminthophila chrysoptera*).—This bird selects a semi-swampy situation, overgrown with bushes, in which to nest. It commences to build the last of May or first of June. The nest is placed on the ground, and is supported laterally by three or four bushes situated from four to five inches apart. It is composed externally of dry leaves placed edgewise, and well lined with fine bark fibres, interspersed with a few coarse hairs. The nest when finished measures, inside, three and one-half to four inches in depth, and one and one half to two inches in diameter; the sides are nearly parallel, only slightly contracted above. It is well concealed by the overhanging bushes and leaves.

The eggs (four or five in number) measure: the largest, $.52 \times .70$; smallest $.46 \times .62$. The average is within a fraction of $.50 \times .68$; they are white, sparingly sprinkled and blotched with light reddish brown, more on the greater end.

I have found but two nests, one, which had young, July 17; the other I discovered while the birds were building. When secured, June 10, it contained five eggs, one of which was a Cowbird's.

Mr. B. F. Goss of Pewaukee has, in his magnificent collection, thirteen nests, containing between fifty and sixty eggs—all taken by one collector