Journal Cincinnati Soc. Nat. Ilist., XIV, Nos. 1, 2, 1891. Observer, The, II, Nos. 6-8, 1891.

Observer, The, II, Nos. 0-0, 1391.

Ornithologist and Botanist, I, Nos. 7, 8, July, Aug., 1891.

Ornithologisches Jahrbuch, II, pt. 4, July, 1891.

Ornithologist and Oölogist, July, August, 1891.

Ottawa Naturalist, V, Nos. 4, 5, 1891.

Proceedings of Linnæan Society of New York, Abstract of, 1891.

Wisconsin Naturalist, I, Nos. 11, 12, June, July, 1891.

Zoologist, July-Sept. 1891.

GENERAL NOTES.

The Yellow-legs (*Totanus flavipes*) Breeding in Ontario County, N. Y.—A pair of Yellow-legs (*Totanus flavipes*) remained this season and bred in a wet pasture in company with Killdeers (Ægialitis vocifera) and Spotted Sandpipers (Actitis macularia). I do not know of any former record of their breeding in the County. The owner of the pasture, who is somewhat of a sportsman, and who first called my attention to the birds, would not permit their being disturbed, as he was anxious that they should make it a regular breeding ground. I repeatedly saw the birds at close quarters and fully identified them. During the latter part of June I noticed four young accompanying the parents who permitted of much closer approach than formerly. The young were about half grown and I endeavored to catch one of them but they eluded pursuit and finally escaped into a swamp close at hand.—B. S. BOWDISH, Phelps, N. Υ .

The Turkey Vulture (*Cathartes aura*) in Ontario County, N. Y.— July 3 Charles Donelly, a young sportsman of this place, shot a Turkey Vulture (*Cathartes aura*) which was one of a flock of eight, seen in the vicinity of a marshy wood. On the 5th the bird was presented to me, and although in poor condition—being badly shot—was preserved as a good specimen. The bird is a female. The flock remained in the vicinity the entire afternoon, but the next morning were gone, and none have been seen since. This is, so far as I know, the first record of the bird being taken in the County.—B. S. BOWDISH, *Phelps, N. Y*.

An Abnormal Specimen of Coccyzus maynardi.—A peculiar example of what is probably *Coccyzus maynardi* has been lately received from Inagua, Bahamas. The bird has the general appearance and marking of *C. maynardi*, but the head, neck, and underparts are plumbeous gray, and the back and upper tail-coverts ash-gray. The quills are pale brown and the tail-feathers black, which, with the exception of the two central feathers are tipped with white. The bird, a female, was taken near Northwest Point, Inagua, May 28, 1891.—CHARLES B. CORY, Boston, Mass.