

Slide and the other on the Table Land at an altitude of a few hundred feet lower than the top of the highest peak (5,215 feet). Unfortunately I was unable to obtain a specimen, but a familiarity with the song of this bird acquired in the White Mountains and during the migrations, leaves no doubt whatever in my own mind of the identification. This subspecies has never been reported from Maine, I believe, though it is included in the 'hypothetical' list in Mr. Ora W. Knight's recent list of Maine birds. The only other birds noted on the mountain which I did not also find in the lowlands about there were *Dendroica striata*, which were common along the Slide, and *Dendragapus canadensis*, one female of which I observed on the Slide. Ktaadn affords but little cover for birds, the upper three thousand feet being for the most part very steep and rocky, giving no chance for trees. The trees along the Slide are almost entirely deciduous, and no coniferous woods were to be seen at any height except those in the great South Basin on the northeast side of the mountain, about 2300 feet below the highest summit or about 3000 feet above sea-level. It is quite possible that these Basin woods may have contained some more northern forms, but I was unable to visit them. Some one should go there in the breeding season. The Basin is best visited from the east side. The scrub fir on the Table Land harbored *Turdus aliciae bicknelli* and *Zonotrichia albicollis*, and doubtless *Junco hyemalis* too, but it is too low to make very good cover.—FRANCIS H. ALLEN, *West Roxbury, Mass.*

**Two Species new to the List of Birds found in West Virginia.**—CHUCK-WILL'S-WIDOW (*Antrostomus carolinensis*), ♂ adult. Picked up in a grove, apparently benumbed by cold, by Mr. John H. Crawford, near Lewisburg, Greenbrier Co., W. Va., on April 23, 1897. Now in Mr. Crawford's possession.

SWAINSON'S HAWK (*Buteo swainsoni*), ♂ adult. Shot by Mr. M. M. Collins four miles north of White Sulphur Springs, W. Va., September 16, 1897. This hawk is now being mounted for Mr. M. M. Collins of Covington, Va.

I believe that, heretofore, Nashville, Tenn., has been about the farthest north, in the interior, from which *Antrostomus carolinensis* has been recorded.—THADDEUS SURBER, *White Sulphur Springs, W. Va.*

**LAKE MICHIGAN NOTES.**—*Larus glaucus*. GLAUCOUS GULL.—While walking along the beach of Lake Michigan, east of Millers, Indiana, August 8, 1897, in company with Mr. J. G. Parker, Jr., and Mr. Fred Hilgard, I had the good fortune to take a fine female of this species. It is in the pure white plumage of the young of the second year; it is immaculate, with the exception of a few feathers on the wing-coverts, which are of a pale brownish gray. I believe this is the first record of capture for the Calumet Region and Indiana.