

tioned there was an almost continuous succession of flocks containing from three or four to as many as one hundred and fifty or two hundred birds in a flock. Their notes, quite unlike those of their red cousins, remind one more of the twittering notes which the Pine Finch utters when feeding. Two or three flocks of Red Crossbills were also noted.

November 21, I again visited the same locality and several hundred 'White-wings' were seen, but very few compared to the numbers seen on the previous day, and all were flying very high. On both days they appeared to be very restless, and when called down would alight in the tops of the cedars. After picking for a moment or two as if in search of food and finding none they would at once resume their journey westward across the harbor. Sixteen specimens were examined and all were in good flesh; their stomachs contained only a little sand and traces of vegetable matter. All of those seen appeared to be adults except one female taken, whose skull indicated immaturity. The red males appeared to be the more numerous.

November 22 again found me in the same locality, but only a single 'White-wing' was seen, this was in company with a flock of Goldfinches. On several occasions from this time on until the 20th of February, when I left Millers Place, single birds or a small flock would be seen, but I think very few spent the winter on the island.

The Red Crossbills (*Loxia curvirostra minor*) were also more numerous than usual during the past winter, and I think a few remained and bred on the island. May 6, two small flocks were noticed among the pitch pines south of Millers Place, one flock of six containing five red males and one female. The following day I procured a female in worn plumage, whose ovaries indicated that they had recently been in an active state. — ARTHUR H. HELME, *Brooklyn, N. Y.*

The Lark Finch and Baird's Bunting on Long Island, N. Y. — An adult male Lark Finch, *Chondestes grammacus* was taken at Millers Place, Long Island, Nov. 27, 1899. It was first noticed feeding in a clump of rag-weeds in company with a small mixed flock of Song, Fox and Tree Sparrows.

At Montauk Point, L. I., Nov. 13, 1899, I secured a small Sparrow that proves to be *Ammodramus bairdi*. The sex I was unable to determine, as it was too badly injured by shot. Dr. Jonathan Dwight, Jr., who has examined the specimen, writes me that it is "chiefly in juvenal plumage passing to first winter." So far as I am aware this is the first instance of its occurrence that has been reported from the Atlantic coast. — ARTHUR H. HELME, *Brooklyn, N. Y.*

Bachman's Sparrow in Virginia. — In May, 1897, I took a pair of Bachman's Sparrows (*Peuceea aestivalis bachmanii*) on Blackwater Creek, Campbell County, Va., together with their nest and eggs. This was the