

locality and were passing down the line of willows along an extensive canal. A strange faint note arising from out of the meadows beyond arrested our attention, the author of which we could not discover. It struck me that this was the object of our search although I had never met with the species before. This supposition proved correct, as a few minutes later Taverner was able to flush a small bird, which he secured, proving it to be a female Henslow's Sparrow. We found a small colony of these birds in this portion of the meadows—about twelve pairs as far as we could judge. The two females taken demonstrated that they were breeding by the bare abdomen and the tough thickened skin of the belly. In the limited amount of time at our disposal no nests were found; indeed, it was a difficult matter to locate the birds themselves, much less the nests. In the vast expanse of meadow the search was in vain. By careful concentration we could detect the *se-slick* notes of the birds arising from out of the medley caused by the Meadowlark, Bobolink, and Red-winged Blackbird, but to an unaccustomed ear the notes would easily be passed over or ignored.

The birds favored perching on the tops of the various weed stalks, golden-rod, asters, etc., that invested certain portions of the field. Once flushed the birds would fly in a peculiar wavering manner a short distance, drop suddenly into the tangled grass, and were with extreme difficulty put up again. Mouse-like they would worm their way through the matted grass and were shortly in a different direction from the place one expected to find them. Towards sunset the birds were more active and the notes would arise from various points. I did not hear them after sunset.

We visited this colony again on July 23 and 24 and August 21. On the latter date the birds had moved up to higher ground where a tangle of golden-rod and other weeds made a good cover. On September 25 thorough search failed to reveal even a single bird. On October 2, in Wayne County, we were passing through a neglected piece of meadow when I flushed a small sparrow which, by its peculiar flight I realized was an *Ammodramus* of some species. This bird quickly dropped into the grass and we searched for an hour before Taverner flushed it again in a distant portion of the field and secured it. This bird proved to be a male Henslow's Sparrow—a much more brightly plumed bird than the June specimens; the dark streaks of the sides showed up very plainly against the ochreous wash on the breast.

I think that *A. henslowii* will prove to be a much more abundant species in this section of Michigan than is supposed, though very locally distributed.—BRADSHAW H. SWALES, *Detroit, Mich.*

**The Northern Parula Warbler in Southern Michigan.**—I am enabled by the courtesy of Mr. J. Claire Wood of Detroit to record the first specimen of the Northern Parula Warbler (*Compsothlypis americana usneæ*) taken in Wayne County, Michigan. On September 25, 1904, Mr. Wood

met with several of these birds and secured an adult male which is now in his collection. I have expected to find this species here for some time, as Mr. Taverner found it at Port Huron, St. Clair County, to the north, and it has been taken in Monroe and Washenaw Counties, southeast of here. — BRADSHAW H. SWALES, *Detroit, Mich.*

**Bachman's Warbler in Leon County, Florida.** — In the October, 1904, number of 'The Auk' Mr. R. W. Williams, Jr., in a list of the birds of Leon County, Florida, says concerning Bachman's Warbler: "Only one record. I took this specimen on August 4, 1900." On March 22, 1904, while in company with Mr. Morgan Hebard of Thomasville, Georgia, I collected a male specimen of this species in a black gum swamp in the extreme northeastern section of Leon County, about four miles distant from the Georgia line. The individual taken was in company with several others which appeared to be the same species, but as the identity was not known until the specimen was in hand, no others were secured. — JAMES A. G. REHN, *Acad. Nat. Sciences, Philadelphia, Pa.*

**The First Hooded Warbler Taken in Maine.** — On the 9th of September, 1904, Mr. Samuel T. Dana, of Portland, informed me that he had seen a Hooded Warbler (*Wilsonia mitrata*) at Falmouth. As this bird had never, to my knowledge, been recorded in Maine, I asked him to secure a specimen, which he did on the 10th of September, 1904. The bird was brought to a Portland taxidermist the same day and mounted. It is now in the possession of Mr. Dana. It is a male bird, an adult, and is in perfect plumage. It was taken in the trees near a dwelling, and had been there several days, in company with other warblers. The capture of this warbler and the establishing of a new record has attracted considerable comment among local ornithologists. — W. H. BROWNSON, *Portland, Maine.*

**Breeding of the Hudsonian Chickadee (*Parus hudsonicus*) at Dover, Me.** — There has been so little said or written in relation to the breeding of this species that the record of a nest with young discovered by the writer the present season may be of some value to the working ornithologists.

During a 12 years' residence at Dover, Piscataquis County, I have occasionally during my rambles met this species, but the meetings have usually occurred during the late fall or winter seasons, and have been so infrequent as to merit a special record in my notes. Accordingly it was indeed a surprise to discover a pair engaged in the act of rearing a brood of young this season. The date was June 21. I had spent the morning botanizing in a place locally known as Sangerville bog, located due west from Dover village, the nearest portion of the bog being about a mile distant. The boundary line between the towns of Dover and Sangerville passes directly through the morass, a portion lying in either town, but the 'find' was located on the Dover side.