

met with several of these birds and secured an adult male which is now in his collection. I have expected to find this species here for some time, as Mr. Taverner found it at Port Huron, St. Clair County, to the north, and it has been taken in Monroe and Washenaw Counties, southeast of here. — BRADSHAW H. SWALES, *Detroit, Mich.*

Bachman's Warbler in Leon County, Florida. — In the October, 1904, number of 'The Auk' Mr. R. W. Williams, Jr., in a list of the birds of Leon County, Florida, says concerning Bachman's Warbler: "Only one record. I took this specimen on August 4, 1900." On March 22, 1904, while in company with Mr. Morgan Hebard of Thomasville, Georgia, I collected a male specimen of this species in a black gum swamp in the extreme northeastern section of Leon County, about four miles distant from the Georgia line. The individual taken was in company with several others which appeared to be the same species, but as the identity was not known until the specimen was in hand, no others were secured. — JAMES A. G. REHN, *Acad. Nat. Sciences, Philadelphia, Pa.*

The First Hooded Warbler Taken in Maine. — On the 9th of September, 1904, Mr. Samuel T. Dana, of Portland, informed me that he had seen a Hooded Warbler (*Wilsonia mitrata*) at Falmouth. As this bird had never, to my knowledge, been recorded in Maine, I asked him to secure a specimen, which he did on the 10th of September, 1904. The bird was brought to a Portland taxidermist the same day and mounted. It is now in the possession of Mr. Dana. It is a male bird, an adult, and is in perfect plumage. It was taken in the trees near a dwelling, and had been there several days, in company with other warblers. The capture of this warbler and the establishing of a new record has attracted considerable comment among local ornithologists. — W. H. BROWNSON, *Portland, Maine.*

Breeding of the Hudsonian Chickadee (*Parus hudsonicus*) at Dover, Me. — There has been so little said or written in relation to the breeding of this species that the record of a nest with young discovered by the writer the present season may be of some value to the working ornithologists.

During a 12 years' residence at Dover, Piscataquis County, I have occasionally during my rambles met this species, but the meetings have usually occurred during the late fall or winter seasons, and have been so infrequent as to merit a special record in my notes. Accordingly it was indeed a surprise to discover a pair engaged in the act of rearing a brood of young this season. The date was June 21. I had spent the morning botanizing in a place locally known as Sangerville bog, located due west from Dover village, the nearest portion of the bog being about a mile distant. The boundary line between the towns of Dover and Sangerville passes directly through the morass, a portion lying in either town, but the 'find' was located on the Dover side.