

Coming from so distinguished an entomologist, these conclusions carry much weight, and they are well worth the attention of certain economic entomologists of the United States, who have expressed very different views of the relative importance of birds and parasitic insects.— W. L. M.

**Bryant, on Relation of Birds to an Insect Outbreak in California.**<sup>1</sup>—

This valuable economic paper deals with an outbreak of a butterfly, *Eugonia californica*, which swarmed over portions of California and the larvæ of which defoliated the Snow-brush and Buck-brush, two species of *Ceanothus*. Mr. Bryant's investigations show that five species of birds fed upon the butterflies. Brewer's Blackbird (*Euphagus cyanocephalus*) being the most important and consuming 95 per cent of all the butterflies eaten by birds. Butterflies seem to be rarely eaten by birds under normal conditions and the change of food in this instance is interesting as illustrating how valuable a bird not usually of economic importance may become under extraordinary conditions. The great benefit entailed in the destruction of female butterflies before or during ovipositing as compared with the destruction of larvæ is also pointed out by the author and he estimates that of one Brewer's Blackbird destroyed 100,000 butterflies in a month and his observations seem to support him. If one third of these were females, the destruction of eggs would amount to 336,000! — W. S.

**Economic Ornithology in Recent Entomological Publications.**—

The following reviews relate exclusively to publications of the U. S. Bureau of Entomology, hence the name of that office is not repeated in the reference. The first article,<sup>2</sup> in point of date of issue, which we desire to note deals with the bill-bug (*Sphenophorus callosus*). This species, which does great damage to corn in many states, is commonly known in the Carolinas as the "curlew-bug." This appellation refers to a point in common between the bird and the beetle — a long curved beak. One bird enemy of the curlew-bug, the Nighthawk, is mentioned on the authority of the Biological Survey. The finding of not only this species, but of several other *Sphenophorus*, in stomachs of Nighthawks, has a bearing on a debated point, *i. e.*, whether these beetles fly. There is no doubt that most if not all of those eaten by the Nighthawk are taken on the wing.

The false wireworms of the genus *Eleodes*, family Tenebrionidæ, are said<sup>3</sup> to do more damage to newly planted grain in the northwestern states than any insect pests other than the true wireworms of the family Elateridæ. The Sage Hen, the Burrowing Owl, and Butcherbird are said to feed upon them and Brewer's Blackbirds often follow the plow to pick up the larvæ and pupæ. Western Bluebirds were seen to feed on larvæ which had

<sup>1</sup> The Relation of Birds to an Insect Outbreak in Northern California during the Spring and Summer of 1911. By Harold C. Bryant. The Condor, Vol. XIII, Nov., 1911, pp. 195-208.

<sup>2</sup> Webster, F. M. The so-called "curlew-bug." Bull. 95, Pt. IV. April 10, 1912.

<sup>3</sup> Hyslop, J. A. Bull. 95, Pt. V, April 22, 1912.

been driven to the surface by a heavy rain. From observations and experiments upon domesticated and other confined birds it was learned that chickens, ducks, and the Reeves and Silver Pheasants will eat the *Eleodes*, and that turkeys and the Golden, and Lady Amherst Pheasants refuse them. The author remarks however, that the latter birds were "quite annoyed by our presence, and might have eaten the beetles had they not been frightened." The account of the natural enemies includes also a list of 13 species of birds which the Biological Survey has found to feed upon adult *Eleodes*.

In his account of 'Two destructive Texas ants,'<sup>1</sup> Mr. W. D. Hunter inserts a list of the known bird enemies of the agricultural ant (*Pogonomyrmex barbatus molefaciens*). This list also is taken from Biological Survey records (with the exception of one name), and includes 8 species of birds.

The 'Preliminary report on the alfalfa weevil,'<sup>2</sup> gives a list of the vertebrate enemies of this new, but important pest, based on the work of Mr. E. R. Kalmbach of the Biological Survey. This represents the results of one season's field work on the relations of birds to this weevil. Thirty-one species are mentioned. A noteworthy point brought out by this investigation is that English Sparrows are among the most effective enemies of the pest, practically rearing their young on a weevil diet. Mr. Kalmbach writes that this season these birds are fully equalling last year's performance.—W. L. M.

**The Food of Birds in Scotland.**—Miss Laura Florence publishes in the Transactions of the Highland Agricultural Society<sup>3</sup> detailed results of the examinations of the alimentary tracts of 616 birds representing 74 species. Since the contents of the intestines as well as of the stomachs were studied, the author is able to present data on the imperfection of digestion in some cases and to indicate possibilities in the distribution of seeds and insect eggs. No general conclusions are drawn, and the results for each species are only very briefly summarized. The detailed information is worthy of record, however, and will no doubt be very welcome to all Britons interested in economic ornithology.

The identity of several of the birds with United States species and the recurrence of many familiar seed and insect names, make an American feel much at home while reading the paper.—W. L. M.

**Scott and Sharpe on the Birds of Patagonia.**—The third installment of the report on the birds of the Princeton University Expedition to Patagonia<sup>4</sup> is presented with the same wealth of illustrations and breadth

<sup>1</sup> Circular 148, April 26, 1912.

<sup>2</sup> Bull. 112, May 14, 1912.

<sup>3</sup> Fifth Ser., Vol. XXIV, 1912, pp. 180-219.

<sup>4</sup> Reports of The Princeton University Expeditions to Patagonia, 1896-1899, J. B. Hatcher in charge. Edited by William B. Scott. Volume II—Ornithology, Part III, Charadriidæ-Anatidæ. By William Earl Dodge Scott associated with R. Bowdler Sharpe. Princeton, N. J.—Stuttgart. Pp. 345-504, text figs. 175-252, pl. 1.