

Additional Notes to the 'Birds of Gallatin County, Montana'.—

These notes form an appendix to my paper published in 'The Auk,' January, 1911, pp. 26-49. The numbers are those of my list.

37. **Gallinago delicata.** WILSON'S SNIPE.— This species appears to be fairly common and regular as a winter resident in warm swamps near Bozeman. I observed seven birds on December 26, 1911.

43. **Catoptrophorus semipalmatus inornatus.** WESTERN WILLET.— Mr. W. L. Thomas of Belgrade found two nests of this species during the summer of 1911.

120. **Hesperiphona vespertina montana.** WESTERN EVENING GROSBEAK.— I observed a small flock in Bozeman on January 2, 1911. The species is evidently occasionally a winter resident, though much commoner as a migrant.

126. **Leucosticte tephrocotis littoralis.** HEPBURN'S ROSY FINCH.— I observed this subspecies in company with *L. t. tephrocotis* in Bridger Cañon, March 17, 1911. Large flocks were seen and about 10 percent. of the birds were of this form.

170. **Vermivora celata lutescens.** LUTESCENT WARBLER.— I observed a male of this species in song, in an aspen grove near Bozeman, July 4, 1910. It probably breeds in the region and the breeding bird is probably *V. c. celata*, but it was neither secured nor observed closely.

199. **Cyanocephalus cyanocephalus.** PIÑON JAY.— A flock of about 40 birds was observed flying over the western part of Bozeman on Sept. 11, 1911. This adds a new species to the county list, and to my knowledge, is the most westerly record for this species in Montana.—ARETAS A. SAUNDERS.

Two new records for Washington State.— The following birds appear to be new to the State list.

Ceryle alcyon caurina. NORTHWESTERN BELTED KINGFISHER.— A specimen taken September 17, 1896, was recently identified for me by Mr. Oberholser as this new form. It was collected by me on Puget Sound, in the vicinity of Tacoma, Wash., which I believe extends their range this much southward. As I have no other specimens, some further collecting must be done in order to prove this to be the resident form.

Zonotrichia albicollis. WHITE-THROATED SPARROW.— At Sherlock, in Thurston County, on October 13, 1912, I was so fortunate as to collect a female of this Sparrow. My bird was with a very large company of Nuttall's Sparrows (*Zonotrichia leucophrys nuttalli*), which had begun to arrive the previous evening, but a careful search failed to reveal any more *albicollis*. An interesting feature of this migration, possibly of some significance, is that the local breeding *nuttalli* had long since left for the south, and the migration in question was a considerable surprise to me. I believe my bird forms a new record for the state, if not for the Pacific coast north of Oregon.— J. HOOPER BOWLES, Tacoma, Wash.