

CHANGES IN THE A. O. U. CHECK-LIST OF NORTH
AMERICAN BIRDS PROPOSED SINCE THE
PUBLICATION OF THE SIXTEENTH
SUPPLEMENT.

THE changes in the A. O. U. Check-List which have appeared from time to time in the several Supplements have originated in the publications of various ornithologists throughout the world, and have been considered by the Committee on Classification and Nomenclature, and accepted or rejected as the case may be.

The cases that come before the Committee are of two kinds:

- (a.) Nomenclatural — cases which are easily settled by correspondence between the members of the Subcommittee on Nomenclature, all of these being governed by a definite Code of rules.
- (b.) Ornithological — much more difficult cases involving the study of series of specimens and resting largely upon the personal opinion of the members, which may vary widely.

It was formerly the custom for the Committee to meet in Washington annually in April to dispose of accumulated cases. Of late years however, this has proven unsatisfactory as neither time nor material was available for the proper consideration of many proposed new forms based upon very slight differences. The following plan has therefore been adopted by the Committee in the consideration of ornithological cases.

To publish an annual list of proposed additions and alterations inviting study and criticism of them by ornithologists, whether members of the committee or not. In succeeding lists attention will be called to any publication endorsing or opposing changes or additions previously proposed, and it is hoped that in this way, the Committee may secure valuable assistance to guide them in reaching a decision, when a new edition of the Check-List is to be published, or at such time as it may be considered desirable to issue a Supplement.

In order to put this plan into operation the Acting Chairman

submits the following summary of changes and additions proposed since the publication of the 1910 edition of the Check-List and not considered in the Sixteenth Supplement,¹ exclusive of purely nomenclatural cases which, as explained above, are disposed of by the Subcommittee on Nomenclature. The list is probably incomplete but omissions can be added in subsequent publications.—WITMER STONE, *Acting Chairman Com. Classification and Nomenclature.*

I. ADDITIONS TO THE CHECK-LIST.

(a) Forms described as new, or rejected forms re-established.

Larus thayeri sp. nov. Ellesmere Land. Brooks, Bull. M. C. Z., LIX, No. 5, p. 373.

Puffinus puffinus bermudæ subsp. nov. Bermuda. Nichols and Mowbray, Auk, 1916, p. 195.

Astrelata cahow sp. nov. Bermuda. Nichols and Mowbray, Auk, 1916, p. 194.

Histrionicus histrionicus pacificus subsp. nov. Pacific coast. Brooks, Bull. M. C. Z., LIX, No. 5, p. 393.

Oidemia deglandi dixoni subsp. nov. Alaska. Brooks, Bull. M. C. Z., LIX, No. 5, p. 393.

Ardea herodias treganzai Court.

“ “ **sanctilucæ** Thayer and Bangs.

“ “ **hyperoncha** subsp. nov. Pacific coast, Cal.

“ “ **oligista** subsp. nov. Santa Barbara Isl.

These four races recognized by Oberholser. Proc. U. S. Nat. Mus., 43, pp. 531–559.

Colinus virginianus taylori subsp. nov. Yuma Co., Col. Lincoln, Proc. Biol. Soc. Wash., May 27, 1915, p. 103.

Lophortyx gambelli sanus subsp. nov. Olathe, Colo. Mearns, Proc. Biol. Soc. Wash., XXVII, p. 113.

Dendragapus obscurus flemingi subsp. nov. Yukon Terr. Taverner, Auk, 1914, p. 385.

Bonasa umbellus thayeri subsp. nov. Digby, N. S. Bangs, Auk, 1912, p. 378.

Bonasa umbellus yukonensis subsp. nov. Yukon Terr. Grinnell, Condor, 1916, p. 166.

Lagopus leucurus altipetens Osgood, revived by Riley, Canad. Alpine Jour., 1912, p. 60.

Lagopus rupestris sanfordi subsp. nov. Tanaga Isl., Aleutian Chain. Bent, Smithson. Misc. Collns. 56, No. 30, p. 1.

Zenaidura macroura caurina subsp. nov. Oregon. Ridgway, Bds. N. and Mid. Amer., VII, p. 348.

¹ ‘The Auk,’ 1912, pp. 380–387.

Buteo lineatus texanus subsp. nov. Corpus Christi, Texas. Bishop, Auk, 1912, p. 232.

Otus asio quercurinus subsp. nov. So. California. Grinnell, Auk, 1915, p. 60.

Otus asio hasbrouckii subsp. nov. Palo Pinto Co., Tex. Ridgway, Bds. N. and Mid. Amer., VI, p. 694.

Otus asio brewsteri subsp. nov. Salem, Oregon. Ridgway, Bds. N. and Mid. Amer., VI, p. 700.

Bubo virginianus neochanes subsp. nov. Newfoundland. Oberholser, Proc. Biol. Soc. Wash. XXVII, p. 46.

Bubo virginianus occidentalis Stone.

" " **icelus** (Oberholser).

" " **lagophonus** (Oberholser).

These three races revived by Ridgway, Birds of N. and Mid. Amer., VI.

Strix occidentalis lucida Nelson with which **S. o. huachucæ** Swarth is regarded as identical. Oberholser, Proc. U. S. N. M., 49, pp. 251-257.

Cryptoglaux acadica brooksi subsp. nov. Queen Charlotte Islands. Fleming, Auk, 1916, p. 422.

Glaucidium gnoma vigilante subsp. nov. Mountains of S. California and southern Sierra Nevada. Grinnell, Auk, 1913, p. 224.

Glaucidium gnoma swarthyi subsp. nov. Vancouver Island. Grinnell, Auk, 1913, p. 224.

Glaucidium gnoma grinnelli subsp. nov. Humboldt Bay, Cal. Ridgway, Birds N. and Mid. Amer., VI, p. 791.

Micropallas whitneyi sanfordi subsp. nov. Lower California. Ridgway, Birds N. and Mid. Amer., VI, p. 809.

Micropallas whitneyi idoneus subsp. nov. Hidalgo, Texas. Ridgway, Birds N. and Mid. Amer., VI, p. 810.

Conuropsis carolinensis interior subsp. nov. Nebraska. Bangs, Proc. N. E. Zoöl. Club, IV, p. 93. [Applicability of *ludovicianus* of Gmelin for this race must also be considered.]

Dryobates pubescens microleucus subsp. nov. Newfoundland. Oberholser, Proc. Biol. Soc. Wash., XXVII, p. 43.

Dryobates pubescens glacialis Grinnell.

" **villosum leucothorectes** Oberholser.

" " **orius** Oberholser.

" " **icastus** Oberholser.

" " **sitkensis** Swarth.

" **scalaris eremicus** Oberholser.

" " **cactophilus** Oberholser.

Colaptes auratus borealis Ridgway.

Xenopicus albolarvatus gravirostris Grinnell.

Phloeotomus pileatus floridanus Ridgway.

" " **picinus** Bangs.

All the above 11, subspecies rejected or held in abeyance by the Committee are recognized by Ridgway, Birds of N. and Mid. Amer., VI.

- Chordeiles virginianus howelli**, subsp. nov. Texas. Oberholser, Bull. 86, U. S. Nat. Mus., p. 57.
- Chordeiles acutipennis inferior** subsp. nov. Lower California. Oberholser, Bull. 86, U. S. Nat. Mus., p. 109.
- Mr. Ridgway (Birds N. and Mid. Amer., VI) endorses these two new races of Nighthawks.
- Setochalcis** gen. nov. for the Whip-poor-will as distinct from the Chuck-will's-widow; and,
- Chordeilidae** fam. nov. for the Night Hawks. Oberholser, Bull. 86, U. S. Nat. Mus.
- Corvus brachyrhynchos paulus**, subsp. nov. Alabama. Howell, Proc. Biol. Soc. Wash., XXVI, p. 199.
- Perisoreus canadensis sanfordi** subsp. nov. Newfoundland. Oberholser, Proc. Biol. Soc. Wash., XXVII, p. 49.
- Agelaius phoeniceus aciculatus** subsp. nov. Kern Co., Cal. Mailliard, Condor, XVII, p. 13.
- Agelaius phoeniceus nevadensis** subsp. nov. Nevada. Grinnell, Proc. Biol. Soc. Wash., XXVI, p. 107.
- Pinicola enucleator eschatosus** subsp. nov. Newfoundland. Oberholser, Proc. Biol. Soc. Wash., XXVII, p. 5.
- Loxia curvirostra percna** subsp. nov. Newfoundland. Bent, Smithson. Miscel. Collns., 60, No. 15, p. 1.
- Carduelis carduelis bermudiana** subsp. nov. Bermuda. Kennedy, Bull. Brit. Orn. Club, CXCI, p. 33.
- Melospiza melodia beata**, subsp. nov. Enterprise, Fla. Bangs, Proc. N. E. Zoöl. Club, IV, p. 85.
- Melospiza melodia acadica** subsp. nov. Nova Scotia. Thayer and Bangs, Proc. N. E. Zoöl. Club, V, p. 67.
- Pipilo maculata falcifer** McGregor.
 " " **curtatus** Grinnell.
 " " **falcinellus** subsp. nov. Interior California. The first two rejected races reestablished and a new one described. Swarth, Condor, XV, No. 5, p. 168.
- Pipilo erythrophthalmus canaster** subsp. nov. Alabama. Howell, Proc. Biol. Soc. Wash., XXVI, p. 202.
- Dendroica coronata hooveri** McGregor.
- Geothlypis trichas scirpicola** Grinnell.
- Hirundo erythrogaster palmeri**.
 Rejected races revived, Swarth, Univ. Cal. Publ. Zoöl. 10, No. 1, pp. 65-71.
- Toxostoma redivivum sonomae** subsp. nov., Sacramento Valley, etc., Grinnell, Distributional List of Birds of Cal.
- Nannus hiemalis semidiensis** subsp. nov. Semidi Isl., Alaska. Brooks, Bull. Mus. Comp. Zoöl., LIX, No. 5, p. 400.
- Salpinctes guadaloupensis proximus** subsp. nov. San Martin Isl., Lower California. Swarth, Condor, XVI, No. 5.
- Leucosticte tephrocotis dawsoni**, subsp. nov. Sierra Nevada. Grinnell, Condor, 1913, p. 77.

(b.) Species which have been found within the limits of the Check-List.

Marila ferina (Linn.).

Marila fuligula (Linn.).

Cryptoglaux funerea funerea (Linn.).

Coccothraustes c. japonicus Temm. & Schl.

All four taken in the Pribilof Islands. Evermann, Auk, 1913, p. 15.

Dendrocygna viduata (Linn.), in New Jersey, G. B. Grinnell, Auk, 1913, p. 110.

Petrochelidon fulva pallida Nelson, in Texas, Bishop, Auk, 1910, p. 459.

Clangula clangula clangula (Linn.).

Fringilla montifringilla, Linn. Pribilof Islands, Hanna, Auk, 1916, p. 402.

II. CHANGES IN NAMES.

(a) Due to the subdivision of species or subspecies and the restriction of the "typical" race to an extralimital range.

Gelochelidon nilotica becomes **G. n. aranea** Wils. Mathews, Birds of Austral., II, pt. 3, p. 329, confirms Coues (Proc. Acad. Nat. Sci. Phila., 1862, p. 536) that the North American form is distinct from the European.

Sterna caspia becomes **Hydroprogne tschegreva imperator** (Coues). Mathews, *do.*, p. 337. Again confirms Coues that the American bird is distinct. [Changes in generic and specific names are considered among nomenclature cases.]

Sterna anætheta becomes **Melanosterna anæthetus recognita** subsp. nov. Mathews, *do.*, pt. 4, p. 397. The American bird proves "different throughout" when compared with the typical form from the Philippines [changes in generic and specific names considered elsewhere].

Puffinus cuneatus becomes **P. pacificus allenii** subsp. nov. Mathews, *do.*, pt. 1, p. 83. *P. cuneatus* of the Marshall Islands is a subspecies of *P. pacificus* (Kermadec Isl.) but the American bird (from San Benedicto Isl.) is different from either.

Puffinus griseus is divided into **P. g. stricklandi** (Atlantic coast) and **P. g. chilensis** (Pacific). Mathews, *do.*, p. 96. True *griseus* is restricted to New Zealand and Australian seas.

Fregetta grallaria becomes **F. leucogaster** Gould. Mathews, *do.*, pt. 2, p. 39. These two species prove different upon examining the types. The latter is the bird referred to in the Check-List.

Phœbetria palpebrata = **P. p. auduboni** subsp. nov. Nichols and Murphy, Auk, 1914, p. 531.

Dafila acuta becomes **D. a. americana** Reich. Thayer and Bangs, Auk, 1916, p. 45. Confirm Reichenow, Orn. Monatsb. IX, p. 17, 1901, that the American and European birds are distinct. [The applicability of the name *tzitzioha* Vieill. for this race is also involved].

Aegialitis hiaticula becomes **A. h. major** Seebold. P. R. Lowe, Ibis, 1914, p. 395.

Squatarola squatarola becomes **S. s. cynosure** subsp. nov. Thayer and Bangs (Proc. N. E. Zoöl. Club, V, April 4, 1914) claim that the American bird differs from the European. Todd (Ann. Carnegie Mus., X, p. 146) fails to find constant differences.

Gallinula galeata becomes **G. chloropus cachinnans** subsp. nov. Bangs, Proc. N. E. Zoöl. Club, V, pp. 93-99. The North American bird is considered separable from that of South America (true *galeata*) and both regarded as subspecies of *G. chloropus* of Europe.

Callipepla squamata squamata becomes **C. s. pallida** Brewst. Bangs, Proc. N. E. Zoöl. Club, IV, p. 99.

Melanerpes formicivorus formicivorus becomes **Balanosphyrus f. aculeata** Mearns. Ridgway, Birds N. and Mid. Amer., VI, p. 105. True *formicivorus* restricted to Mexico. [Generic name considered elsewhere.]

Dryobates scalaris bairdi becomes **D. s. symplectus** Oberholser. Ridgway, Birds N. and Mid. Amer., VI, p. 257. Endorses Oberholser in restricting true *bairdi* to Mexico.

Passer domesticus becomes **P. hostilis** Kleinschmidt, Falco, XI, No. 2, Dec. 1915. The English bird, from which our American birds are probably descended, is separated from the continental *P. domesticus*.

(b.) Species which are changed to subspecies.

Thalassogeran culminatus becomes **T. chrysostoma culminatus** Mathews, Birds Austral., II.

Oceanodroma kaedingi becomes **O. leucorhoa kaedingi** Willett, Auk, 1915, p. 301.

Somateria dresseri becomes **S. mollissima dresseri** Townsend, Auk, 1916, p. 291.

Aluco pratincola becomes **Tyto alba pratincola** Hartert, Vög. pal. Fauna, Heft VIII.

Otus xantusi becomes **O. asio xantusi** Ridgway, Birds N. and Mid. Amer., VI.

Atthis morcomi becomes **A. heloisa morcomi**.

Amizilis cerviniventris chalconota becomes **A. yucatanensis chalconota**, Ridgway, Birds N. and Mid. Amer., V.

Corvus caurinus becomes **C. brachyrhynchos caurinus**. Swarth, Univ. Cal. Publ. Zoöl. 10, No. 1, cf. also Rhoads, Auk.

(c) Subgenera raised to Genera or New Genera proposed.

Melanosterna
Phœbastria
Thalasseus
Hydroprogne
Onychoprion
Sternula

} Mathews, Birds Austral., II. The first two are not even recognized as subgenera in the Check-List.

Oreopeleia	Ridgway, Birds N. and Mid. Amer., VII.
Chlorocenas	
Patagioenas	
Hierofalco , B. O. U., List of British Birds.	
Cerchneis , Chapman, Bull. A. M. N. H., XXXIV, p. 363. Cory, Field Mus. N. H., I, No. 9, p. 303.	
Megaceryle	Miller.
Chloroceryle	Ridgway, Birds N. and Mid. Amer., VI.
Balanosphyra	Ridgway, Birds N. and Mid. Amer., VI.
Phrenopicus	
Trogonurus	Ridgway, Birds N. and Mid. Amer., V.
Nephœcetes	
Corthylio , Miller, Auk, 1915, p. 234.	

III. FORMS WHICH IT IS PROPOSED TO REJECT.

Ixobrychus neoxenus , a melanistic phase of <i>I. exilis</i> . Bangs, Auk, 1915, p. 483.
Ardea herodias wardi .
Ardea wurdemanni , both of these are phases of <i>A. h. occidentalis</i> : Bangs, Auk, 1915, p. 484.
Falco rusticulus rusticulus
Falco r. gyrfalco Hartert considers all the American Gyrfalcons referable to <i>F. r. candicans</i> (N. A. and Greenland) and <i>F. r. obsoletus</i> (Labrador). Novit. Zool. XXII, pp. 167-185.
Strix occidentalis caurina = <i>S. o. occidentalis</i> . Oberholser, Proc. U. S. N. M., 49, p. 251.
Otus flammeolus idahoensis = <i>O. f. flammeolus</i> . Ridgway, Birds N. and Mid. Amer., VI.
Cryptoglaux acadicus scotæus = <i>C. acadicus</i> . Ridgway, Birds N. and Mid. Amer., VI.
Coccyzus minor minor = <i>C. m. maynardi</i> , so far as North American records are concerned. Ridgway, Birds N. and Mid. Amer., VII.
Phalaenoptilus nuttalli nitidus = <i>P. n. nuttalli</i> . Ridgway, Birds N. and Mid. Amer., VI.
Uranomitra is united with <i>Amizilis</i> . Ridgway, Birds N. and Mid. Amer., V.
Melospiza melodia morphna = <i>M. m. rufina</i> . Swarth, Univ. Cal. Publ. Zoöl., 10, No. 1, p. 61.
Dendroica cærulescens cairnsi = <i>D. cærulescens</i> . Eifrig, Auk, 1915, p. 108.
Geothlypis trichas arizela = <i>G. t. occidentalis</i> . Grinnell, Publ. Zoöl. Univ. Cal., 12, No. 4, p. 202.