

it with *Emberiza* Brisson, but an examination of its type and comparison with typical species of *Emberiza* shows that it is well differentiated as a generic group. It differs from *Emberiza* Brisson (type, by tautonymy, *Emberiza citrinella* Linnæus) as follows; bill slenderer, more compressed, more sharply pointed, thus less conical; basal two-thirds of culmen straight or even somewhat concave, instead of convex; maxillar and mandibular tomia vertically not so strongly concave, thus not giving the closed commissure the somewhat open appearance that it has in typical species of *Emberiza*; palatal surface of maxilla lacking the peculiar rounded protuberances of *Emberiza*; mandible more rounded (less squarish) basally; gonys very long, its length much more than the height of the bill at base (instead of about equal to that dimension), and not strongly ascending, the gonydeal angle therefore not so prominent; tertials and tail much shorter.

The species to be included in this genus are at least the three originally indicated by Cabanis, the last one of which is North American by reason of its accidental occurrence on Kiska Island in the Aleutian Islands, Alaska. These are:

Hypocentor aureolus (Pallas).

Hypocentor fucatus (Pallas).

Hypocentor rusticus (Pallas).

HARRY C. OBERHOLSER, *Washington, D. C.*

A Correction Involving Some Juncos.—An error that may be explained as due to oversight, inadvertence, plain stupidity or all three combined, crept into my paper on the Juncos (Bull. Am. Mus. Nat. Hist. XXXVIII, 1918, p. 296) and Mr. Todd has called my attention to it. In placing *insularis* under *mearnsi* as a race, I quite forgot that the former name has many years priority. Therefore the Pink-sided Juncos should stand as follows:—

Junco insularis mearnsi

Junco insularis insularis

Junco insularis townsendi

JONATHAN DWIGHT, *M. D. New York City.*

An Additional Record of *Ammodramus savannarum bimaculatus* in Eastern Washington.—Although the breeding range of the Western Grasshopper Sparrow is stated by the Check List (A. O. U. Check-List of North American Birds, 1910, p. 257) to embrace "Transition and Austral zones from southeastern British Columbia, northwestern Montana, and southern Minnesota south to southern California and southern Texas," it appears that only one actual record of occurrence in eastern Washington has been published to date. Dr. Lee R. Dice took two adult males in breeding plumage in a wheat field in the Touchet Valley, near Prescott, Walla Walla County, on June 16, 1908 (Auk, Vol. XXVII, 1910, p. 217).

On May 29, 1918, a bird which I am practically certain was of this species was encountered in a grassy swale not far from Pullman, Whitman County. When first sighted it was perched on a grassy tussock near the bottom of the swale. When flushed it flew to a grass clump some distance up a gentle hill slope, disappearing from view in the usual slinking fashion. Too much reliance cannot, of course, be placed on this record, since the bird was not secured.

On June 13 I noted the song of a Grasshopper Sparrow in a grain field near Six Mile Ranch, six miles south of Sprague, just over the line in Adams County. The bird was pursued for some time before it was finally taken. Its actions were as usually described, the bird characteristically dropping behind a grass tussock, ledge of earth or pile of brush, and then, with bill low, body in crouching position, and tail drooping, sneaking off through the grassy vegetation, refusing to flush until one was too close to shoot.

The bird is now No. 262090, U. S. National Museum, Biological Survey Collection. It is a male in much worn plumage.

These experiences during the past field season indicate that the Grasshopper Sparrow is probably more common in eastern Washington than has previously been supposed.—WALTER P. TAYLOR, *Biological Survey, Washington, D. C.*

The Dickcissel in New Hampshire.—At Concord, New Hampshire, on October 13, 1918 I shot a male Dickcissel (*Spiza americana*) in immature plumage. It was alone at the moment, in birches at the edge of woods that bordered extensive fields of corn and stubble, the autumnal resort of sparrows of several kinds, which were then swarming there among the weeds. The only records of the bird from north and east of Massachusetts with which I am acquainted are as follows:

Maine, September 29, 1884. C. W. Townsend (Auk, 1885, p. 106).

Maine, October 10, 1888. A. H. Norton (Auk, 1893, p. 302).

Nova Scotia, September 13, 1902. J. Dwight, Jr. (Auk, 1903, p. 440).

FRANCIS BEACH WHITE, *Concord, N. H.*

Early Nesting of the Loggerhead Shrike *Lanius ludovicianus ludovicianus* at Savannah, Ga.—I am indebted to Mr. Gilbert R. Rossignol, of Savannah, Ga., for the privilege of announcing the taking by him on February 15, 1919, at Savannah, of a nest and five eggs of the Loggerhead Shrike. Mr. Rossignol first discovered the birds building the nest in a live oak tree, among a cluster of vertical shoots, on January 16. The eggs were all fresh and the nest was approximately twenty feet from the ground.

In the vicinity of Charleston, S. C., the earliest dates upon which I have found eggs were on March 24, 1916, six eggs almost hatched, and March 13, 1917, five fresh eggs, both nests being found in the same live oak tree and doubtless belonging to the same pair of birds.—ARTHUR T. WAYNE, *Mt. Pleasant, S. C.*