winter a flock of thirty spent several days about my feeding shelf. One bird flew down a chimney and was killed at Rockport, Illinois, just thirty miles south and was sent to me for identification. At the Illinois State Normal School at Macomb, about sixty miles northeast a solitary live bird was captured. Evidently the Starling is now established in western Illinois.—T. E. Musselman, Quincy, Illinois.

European Starling in upper Peninsula of Michigan.—On September 2, 3 and 7, 1927, a flock of 79 European Starlings was observed feeding in stubble fields with Crows near Sterlingville, which is in Chippewa County, about twenty-four miles south of Sault Ste. Marie, Michigan. When first seen on September 2 we passed within thirty feet of part of the flock as they sat on a fence, so that there was no uncertainty in identification. These Starlings were in the usual autumn plumage with dark bills, brownish backs and speckled breasts.

On the 7th the Starlings were among the hordes of migrating Savannah Sparrows. Several Sharp-shinned Hawks and one Cooper's Hawk were about, apparently preying upon the Sparrows and the bulk of the Starlings were seen to surround the Cooper's Hawk and chase it into a forest. While in the air the fifty or sixty Starlings surrounded the Hawk in a globular flock, perhaps fifteen feet in diameter, wheeling with the Hawk until the timber was reached.

A single Starling was seen by Mr. Walker at Mackinaw City, the northernmost point of the Lower Peninsula, on August 17.—Wm. G. Fargo, *Jackson, Michigan* and Charles F. Walker, *Columbus, Ohio*.

Starling Notes from Lower Michigan.—It is surprising how rapidly the European Starling has increased over Lower Michigan during the past three years.

During the year 1925 we located one pair of these birds breeding in Oakland County. The following year, 1926, we located six nests of this species one in Genessee County, two in Livingston County and three in Oakland County. That same fall together with Mr. Josselyn VanTyne of Ann Arbor we observed great flocks using a wooded area near the Huron River just east and a little North of Ann Arbor for a roost, the birds were mixed with Cowbirds, Bronzed Grackles and Red-winged Blackbirds.

During the year 1927 we found many nests in a great many different counties throughout the lower half of the Lower Peninsula. One of the pairs that we had under observation, raised three broods of young. Others raised two. The birds used bird boxes, hollow trees and Woodpecker holes and similar places, many nested in the crevices of out buildings and one pair used the sky-light of an office building.

On June 3 a small flock was seen in Lake County, near Baldwin; June 10 and 11, in Charlevoix county; July 10, in Cheboygan county; July 30, in Grand Traverse county. The birds were also observed in Roscommon.