

from the big Egret. The two occipital plumes were rather conspicuous on one of the birds.

Another Great White Heron was seen December 28, 1931, between Long Key and Marathon.

Aramus pictus pictus. LIMPKIN.—A bunch of seven was seen along the Kissimmee River near the highway from Okeechobee to Lake Childs, December 19, 1931. They were feeding along the open, somewhat muddy banks of the river and kept in a rather close flock. The head and neck were jerked with each step and at the same time a soft cluck was given.

Phaeopus hudsonicus. HUDSONIAN CURLEW.—Two birds definitely of this species were seen December 3, 1931, with a mixed flock of shore-birds on a small sand-flat in Sarasota Bay, near Whitfield Estates. The Curlew were seen under favorable conditions with 8 x glasses and Willets, Black-bellied Plover and Dowitchers were present for comparison. The decurved bill, crown stripes and size (larger than a Willet) were all noted.

Gelocheidon nilotica aranea. GULL-BILLED TERN.—On December 7, 1931, two of these birds were seen in Sarasota Bay, near Whitfield Estates. Howell, in his new 'Florida Bird Life,' p. 261, gives two other winter records of this species, one in December, 1886, for Hillsborough County, and the other January, 1926, in Wakulla County.—PHILIP A. DUMONT, Berkeley, California.

Notes from Logan, Utah.—In a previous note (Auk, Oct. 1931, p. 611) by the writer, *Junco annectens* Baird was reported as a new record for Utah. Only the male was determined. The skin of the female has since been determined by Dr. Oberholser as *Junco mearnsi* Ridgw. The hybrid species, *J. annectens* (now dropped from the 'Check-List') was therefore mated with *J. mearnsi* known to breed in Utah but seldom reported. I took another female *J. mearnsi* in the mountains east of Logan, Utah, in July 1931.

Two specimens of *Dendroica townsendi* originally reported elsewhere (Univ. Ut. Bul. vol. 21 no. 8) as a new Utah record (from the Henry Mts.) were taken in Logan, September 27, 1930.

One *Vireo solitarius cassini* was also collected the same day from among the many migrants present in the boxelders by my house.

An immature *Lanius ludovicianus gambeli* was collected July 23, 1931, near Promontory, Boxelder Co., Utah.—J. S. STANFORD, U. S. A. C., Logan, Utah.

Some Emendations to the Ranges of the New Check-List.—An attempt to add to, or detract from, the consummation of the work of the Committee which produced the new 'Check-List' would seem to smack of presumption, but it is not with any such intention that these words are written. The 'Check-List' is a human production and none of humanity's works are perfect. That some inaccuracies occur in this revolutionary publication is only natural, and in order to make the record as accurate as possible it would seem desirable for those who have unpublished data of