

May 13, 1932. Eighteen Starlings were seen at Port Daniel, on the southern side of the Gaspé Peninsula. They were not in a flock, but were scattered and seemed to be well established. One pair was seen copulating. Undoubtedly they were nesting at this place.

May 19, 1932. Fifteen Starlings were seen at Murray Bay, Charlevoix County, Quebec, on the north shore of the St. Lawrence River, about eighty miles east of Quebec City. Apparently this species was well established and was nesting at this village.

September 24, 1932. A flock of eighteen Starlings was seen at Pownal, Queens County, Prince Edward Island.

May 9, 1933. Three Starlings were seen together, feeding on the ground among short grass, on Grindstone Island, Magdalen Islands, Quebec, near the central part of the Gulf of St. Lawrence. Half an hour later two of these birds were seen perched together on a telephone wire. One of the two was singing at this time, and it is probable that these two birds were a mated pair.

June 13, 1933. A flock of five Starlings was seen about sunset in conifers at the border of a grassy clearing at Betchewun, Saguenay County, Quebec, on the north shore of the Gulf of St. Lawrence. They seemed very shy and wild. Apparently they had roosted for the night, but when I approached them they flew to another tree close at hand, then, after a pause, flew away out of sight.

June 17, 1933. A Starling was seen at Natashquan, Saguenay County, Quebec. Natashquan is on the north shore of the Gulf of St. Lawrence at a point north of the eastern end of Anticosti Island and is about sixty miles east of Betchewun.—HARRISON F. LEWIS, *National Parks of Canada, Ottawa, Canada.*

Does Bachman's Warbler Winter in Florida?—Recently while looking over the Alden H. Hadley collection of birds at Earlham College, Richmond, Indiana, I examined a Bachman's Warbler (*Vermivora bachmani*). The label is as follows:—"No. 3790. *Helminthophila bachmani*, male, Melbourne, Fla., Jan. 27, 1898."

Mr. Arthur H. Howell in his 'Florida Bird Life,' p. 389, says: "Bachman's Warbler is a very early migrant, the earliest record in the United States being that of a male collected by A. H. Helme at Lukens, February 27, 1909." It seems that the bird first above mentioned is either the earliest migratory record or that the bird at least occasionally winters in Florida.—AMOS W. BUTLER, 52 Downey Ave., Indianapolis, Indiana.

The Calaveras Warbler in Montana.—Apparently the only published record of the occurrence of the Calaveras Warbler (*Vermivora ruficapilla ridgwayi*) in Montana is that of a specimen taken by Silloway at Flathead Lake in 1912 (Saunders, Condor, XVIII, p. 86). In Lincoln County, which occupies the extreme northwestern corner of the state, this species occurs as a very rare summer resident, being found, during