Record of Snow Bunting (Plectrophenax n. nivalis) in North Carolina.—Following the verbal report of a Snow Bunting in eastern North Carolina, the following letter from Miss Grace Wheat, of New York State, is submitted: "The Snowflake was first seen by me near the Love Cottage at White Lake, N. C. (Bladen County), on January 23, 1934. It was feeding with several Fox Sparrows and many White-throated Sparrows under the grape vines in the pullet pen. I saw the bird three days in succession near the buildings, and during three different cold spells after that the bird was seen feeding with Sparrows. Never but one seen. There is no question as to the identity of the Snowflake, as I have seen flocks of them in the dooryard of my home in central New York state."

I have no hesitancy in presenting the foregoing letter as constituting an acceptable record.—H. H. Brimley, *Raleigh*, *N. C.* 

Notes on Some Rare Birds in Southwestern Maine.—Colinus virginianus virginianus. Bob-White.—After many years of absence this bird appeared in Berwick in 1932; a flock was seen within half a mile of our house that season. In 1933 one was heard near the house, and it was repeatedly heard in Wells the same season.

Zenaidura macroura carolinensis. Mourning Dove.—Though of regular occurrence in southern Maine, its occurrence is rare enouth to be of interest. I saw one at Berwick August 17, 1933; flushed another in Wells, August 10, 1934, and on June 6, 1935, saw a pair feeding about two miles from this place in Berwick. I saw one July 26, 1928, at Mount Desert Island in Hancock County.

Telmatodytes palustris dissaeptus. Long-billed Marsh Wren.—Many years ago this bird appeared in our marsh in Berwick, bordering the Salmon Falls River, where I found one of its nests. I have not seen it here since that season.

Cistothorus stellaris. Short-billed Marsh Wren.—This evasive little Wren is now known to be a summer resident in various places in Maine, eastward to the Penobscot River valley and northward somewhat beyond Bangor; but few of these stations have been recorded. July 12, 1933, near Belgrade Station in Kennebec County, in a marshy place I saw closely and heard the outpouring of song of one of these birds. As the botanizing party of which I was a member was moving on, I had not time to look for the nest which I suspected to be near. Several others saw the bird at the same time.

Corthylio calendula calendula. Ruby-crowned Kinglet.—On July 9, 1932, I closely observed a young Ruby-crowned Kinglet in Berwick, the first I have ever seen here in summer.

Dendroica discolor discolor. Prairie Warblers.—July 24, 1921, I saw a pair of Prairie Warblers feeding young, out of the nest, in a scrub oak barren in Sanford, York County, Maine, and again in the same area July 6, 1922, I saw them. No opportunity to visit that section again occurred until the summer of 1932, when I found that an extensive and very destructive fire had consumed all vegetation in the region. Though this bird has not been reported from Maine its occurrence in New Hampshire, some forty-five miles distant, has long been reported.

Anmodramus savannarum australis. Grasshopper Sparrow.—For over forty-five years, to my certain knowledge, at least two pairs of these birds have nested annually on our farm in Berwick. So far this year, (1935), I have seen only one singing male.—Anne E. Perkins, M.D., Berwick, Maine.

Bird Records New or Uncommon to Maryland.—While engaged in field work in Worcester County on the coast of Maryland, May 22, 1935, the writers observed a flock of four Brown Pelicans (*Pelecanus o. occidentalis*) on Sinepuxent Beach some