TURA BARNÉS, JR., Division of Fisheries and Wildlife Conservation, Department of Agriculture and Commerce, Mayaguez, Puerto Rico.

Hooded Merganser in St. Croix, Virgin Islands, U. S. A.—This is an exciting record for Lophodytes cucullatus and one that could have been verified only by the taking of a specimen which I collected from among three birds seen on December 18, 1944. The birds were rather unsuspicious of my presence in the little pond at Rust-op-twist, and quietly swam ahead of me into the mangroves at one point only to emerge soon afterwards at a distance no greater than twenty-five yards on my right. The specimen is a female in winter plumage. It appeared that all three birds were identical in plumage and, therefore, probably were of the same sex. No other species of waterfowl were seen on the pond.—Harry A. Beatty, Christiansted, St. Croix, Virgin Islands.

Sight record of the Western Grebe on Long Island.—While taking a Christmas census for Bird-Lore on December 24, 1934, with Messrs. Grier and Robert Ralston, I noticed a large grebe near the bluffs at Old Field Point, Port Jefferson Harbor, Long Island, New York. Careful study with binoculars revealed that it was a Western Grebe (Aechmophorus occidentalis). The characteristic markings and posture, as well as the distinctive shape of the bill, were clearly evident. Unfortunately, the bird had flown by the time my companions joined me. Because of the rarity of this species on the Atlantic coast, this observation was omitted from the census published in Bird-Lore. Recently, however, a number of records of the occurrence of this grebe in the east have been published, and it seems proper now to add this observation to them. Cruickshank (Birds of New York: 53, 1942) records a previous observation of the Western Grebe on Long Island, at Long Beach, on May 21, 1916.—Fred Mallery Packard, Lt. (j. g.), USNR.

Branta c. hutchinsi on the Atlantic coast.—Taverner, in his account of the Canada Geese (National Museum of Canada, Ann. Rep. for 1929: 30-40, 1 pl., 4 figs., 4 tables, 1931), made sufficient allowance for stragglers of Branta canadensis hutchinsi occurring on the Atlantic Coast, although the main southward flight appears to be through the northern Great Plains and the Mississippi Valley to the Gulf Coast. The A. O. U. Check-List summary (4th ed.: 38, 1931): "Casual on the Atlantic Coast (Maryland and North Carolina)" is, however, too restrictive. Considering the general ranges of the subspecies, it is a fair presumption that any diminutive Canada goose occurring on the Atlantic Coast is hutchinsi. Definite proof to the contrary would be required.

The following records accumulated in the study of bird names indicate more frequent occurrence on that seaboard than appears to be realized. Moreover, the existence of popular names for these small geese probably in most cases indicates repeated occurrence.

Locality	Reference	Local names
New Glasgow, Nova Scotia	J. B. Saunders*	grey-bellied goose
	(*a collaborator of the Biological	mud goose
	Survey; others marked the same)	small goose
Cape Island, Nova Scotia	Bonnycastle Dale	little Canada
	(Rod & Gun in Canada, 25(4):	
	237, 1923)	
Port Joli, Nova Scotia	Bonnycastle Dale	southern goose
	(Rod & Gun in Canada, 25(4):	
	237, 1923)	