

MEASUREMENTS OF *Lanius souzae*, IN MILLIMETERS

Locality or area	Number of specimens	Wing	Tail	Remarks
<i>L. s. souzae</i>				
Angola	12 males	84.0-90.0	81.0-90.0	Includes measurements of two males in the Berlin Museum, kindly provided by E. Stresemann, and those given by Chapin
	13 females	81.0-88.0	76.0-89.0	
Leopoldville, Belgian Congo	1 female	81.0	80.5	In Berlin Museum, as above
Kasai	1 male	81.0	78.0	
<i>L. s. burigi</i>				
N. W. Tanganyika Territory and E. Ruanda	1 male	86.0	89.0	As per Chapin
	1 female	81.0 (+ 3)	79.0	
Uvinza	1 male	83.0	80.0	
	1 female	83.0	81.0	
Mumbwa	1 male	87.5	81.0	
Elisabethville	4 females	80.0-83.0	75.0-83.0	
Nyasaland and Furancungo	7 males	79.0-84.0	79.0-86.0	Includes measurements of three males and one in Transvaal Museum, kindly provided by M. J. Toerien
	6 females	80.0-82.0	80.0-82.0	
Ugano, southern Tanganyika Territory	2 males	79.0-80.0	69.0-71.0	See Ann. Naturh. Mus. Wien, 51: 244, 320, 1941. Kindly measured by M. Sassi.
	3 females	78.0-80.0	68.0-80.0	

A juvenile from Kasai, in the British Museum, has the whole upperside, including the crown and nape, vermiculated on a rich rufous brown, as on lower back of adults of *L. s. souzae*. The underside is similarly vermiculated, though the throat and center of the abdomen are less heavily so.—C. W. BENSON, c/o *The Secretariat, Zomba, Nyasaland*.

***Vireo solitarius* in Honduras.**—A number of bird skins, taken by the well known collector, C. F. Underwood, were recently obtained by the Academy of Natural Sciences of Philadelphia from the Republic of Honduras. Included is an example of one of the Central American forms of the Solitary Vireo from La Flor, Archaga, approximately 12 miles north of Tegucigalpa. The specimen, a male, was collected on June 8, 1937, and has the following measurements: wing (maximum measurement) 65.5 mm.; tail, 45 mm.; exposed culmen, 9.5 mm. The bird is immature, which doubtless accounts for its small size. It cannot be identified subspecifically but is probably referable to *montanus* van Rossem, of Guatemala and El Salvador, rather than to *notius* Van Tyne, of British Honduras.—JAMES BOND, *Academy of Natural Sciences, Philadelphia, Pennsylvania*.