

- SALVIN, O. 1888. Critical notes on the Procellariidae. *Ibis*, 6, 3rd Ser.: 351-360.
- SAUNDERS, H., AND O. SALVIN. 1896. Catalogue of the Gaviae and Tubinares in the collection of the British Museum, 25: 1-475.
- SHARPE, R. B., AND W. R. OGILVIE-GRANT. 1898. Catalogue of the Plataleae, Herodiones, Steganopodes, Pygopodes, Alcae, and Impennes in the collection of the British Museum, 26: 1-687.
- SEEBOHM, H. 1890. On the birds of the Bonin Islands. *Ibis*, 2, 6th Ser.: 95-108.
- SEEBOHM, H. 1891. On the birds of the Volcano Islands. *Ibis*, 3, 6th Ser.: 189-192.
- SNOW, H. J. 1897. Notes on the Kuril Islands. Royal Geographical Soc. London, John Murray.
- SNOW, H. J. 1910. In forbidden seas. London, Edward Arnold.
- U. S. DEPARTMENT OF THE INTERIOR. 1957a. National intelligence survey gazetteer: West Pacific islands. Washington, D. C., Office of Geography, Central Intelligence Agency.
- U. S. DEPARTMENT OF THE INTERIOR. 1957b. National intelligence survey gazetteer: South Pacific islands. Washington, D. C., Office of Geography, Central Intelligence Agency.
- U. S. NATIONAL ARCHIVES. 1923. Rec. Group 37, U. S. Hydrographic Office, corresp., Dec. 1923, 277432/251084.
- U. S. NAVY DEPARTMENT. 1923. Notice to mariners. Hydrographic Office, 52: 1415.
- U. S. NAVY DEPARTMENT. 1943. Gazetteer (No. 4) Hawaiian Islands. Hydrographic Office, Misc. No. 10,884. Washington, D. C., Government Printing Office.

CHARLES A. ELY, *Zoology Department, Fort Hays Kansas State College, Hays, Kansas 67601*, and A. BINION AMERSON, JR., *Pacific Ocean Biological Survey Program, Smithsonian Institution, Washington, D. C. 20560*.

Inland records and first specimen of Black-legged Kittiwake from Mississippi.—On 25 January 1969 I noted an immature Black-legged Kittiwake (*Rissa tridactyla*) among a flock of 125 Bonaparte's Gulls (*Larus philadelphia*) which were resting on the waters of "Lower Lake," a basin lying below the dam and outlet channel of Sardis Lake, Panola County, in northern Mississippi. The bird was seen again on 26 January and 7 and 10 February frequenting the basin area and outlet channel and sharing the latter as feeding ground with Bonaparte's Gulls and Ring-billed Gulls (*Larus delawarensis*). Y. J. McGaha and I collected it on the last date and found it to be a female in first winter plumage weighing 317 g. This specimen, now No. 3565 in the Vaiden Collection of the University of Mississippi Department of Biology, is the first taken in Mississippi.

The first two sightings of the Black-legged Kittiwake in Mississippi were recorded from the Gulf Coast area. A single immature, possibly the same individual in each case, was reported offshore in Mississippi Sound on 28 October 1961 and 11 March 1962 by two groups of observers (*Audubon Field Notes*, 16: 46, 338, 1962). My only previous observations for Mississippi was another inland record of two immatures on 4 December 1967 (*A.F.N.* 22: 51, 1968) that were found on Grenada Lake, another federal flood control reservoir 42 miles south of Sardis Lake. No certain sightings were obtained in later efforts to find and collect the birds on 9 and 20 December 1967 and 17 January 1968.—W. MARVIN DAVIS, *Department of Pharmacology, University of Mississippi, University, Mississippi 38677*.