

INDEX TO VOLUME 106

COMPILED BY K. ELIZABETH K. PACALA

This index is primarily one to new information. Previously published information referred to in this volume has not been indexed unless extensive comparisons or reinterpretations are involved. For some species, especially ones not in the AOU check-list, only scientific names have been entered, and, in some instances, entries under scientific names are more extensive than those under vernacular names. New generic, specific, and subspecific names are printed in **boldface** type.

- Accipiter cooperii*, 451, 452
A. gentilis, 227-230, 452
A. nisus, 225-231
A. striatus, 451, 452
Acrocephalus scirpaceus, 501-502
Aechmophorus clarkii, 467-470
A. occidentalis, 467-470
Aegolius acadicus, 732-733
Afik, Daniel, see Ward, David, et al., and David Ward,
Incubation of dead eggs, 726-728
Agelaius phoeniceus, 107-115, 358-365, 577-582
Ainley, David G., see Boekelheide, Robert J., and —
Alberta, 259-266, 268, 439-445, 635-643, 645-650
Amadon, Dean, In memoriam: Frank B. Smith, 466;
review by, 174
Amazilia saucerottae, 203-207
American Ornithologists' Union, William Brewster
Memorial Award, 351; Elliot Coues Award, 352;
Thirty-seventh supplement to the American Ornithologists' Union Check-list of North American Birds, 532-538; Proceedings of the One-Hundred-Sixth Stated Meeting, 1AA-28AA
Ampelion sclateri, 422-431
A. rubrocristata, 422-431
A. rufaxilla, 422-431
A. stresemanni, 422-431
Ampeloides tschudii, 426, 428
Anatomy and morphology of *Ampelion rubrocristata*, 425-431; *A. rufaxilla*, 425-431; *A. sclateri*, 425-431; *A. stresemanni*, 425-431; *Ampeloides tschudii*, 426, 428; *Carpornis cucullatus*, 426, 428; *Charadrii*, 154-155; *Cinclocerthia ruficauda*, 253; *Conioptilon mcilhennyi*, 426, 428; *Coturnix japonica*, 282, 283, 287; *Geospiza difficilis*, 145-147; *G. fuliginosa*, 145-147; *Lipaugus subalaris*, 426, 428; *Macronectes giganteus*, 303, 309, 310-311; *Oceanodroma leucorhoa*, 80-84; *Phoenicircus nigricollis*, 426, 428; *Phytotoma raimondii*, 425-431; *P. rara*, 425-431; *P. rutila*, 425-431; *Pygoscelis papula*, 303, 309, 310-311; *Ramphocinclus brachyurus*, 254; *Selasphorus rufus*, 724-725; *Scolopaci*, 154-155; *Xipholena punicea*, 426, 428
Anderson, Derek J., see Kikkawa, Jiro, and —
Anderson, Ted R., Determinate vs. indeterminate laying in the House Sparrow, 730-732
Andrew, P., see Collar, N. J., and —
Andrle, Robert F., and Janet R. Carroll (Eds.), The atlas of breeding birds in New York state (rev.), 349
Anhinga anhinga, 383-388
A. novaehollandiae, 383-388
A. rufa, 383-388
Antarctica, 303-312
Anthony, Robert G., see Eberhardt, Lester E., et al.
Anthropoides paradisea, 595-601, 603-613, 616-618
A. virgo, 595-601, 603-613, 616-618
Anthus hodgsoni, 728-729
Aphelocoma coerulescens coerulescens, 742
A. ultramarina, 124-128
Aquila chrysaetos, 520-521, 521-522, 523-524
Aramus guarauna, 603-609, 612, 616-618
Arenaria interpres, 474-476
Arizona, 49-62, 480-482
Arnold, Todd W., Commentary: Sex ratios of fledgling Golden Eagles and jackrabbit densities, 521-522
Atkinson-Willes, G. L., see Owen, Myrfyn, et al.
Atwood, Jonathan L., Speciation and geographic variation in Black-tailed Gnatcatchers (rev.), 347-349
Auklet, Cassin's, 495-497
Austad, Steven N., review by, 165-166
Balda, Russell P., In memoriam: Klaus Immelmann, 723
Balearica pavonina, 595-601, 603-613, 616-618
B. regulorum, 595-601, 603-613, 616-618
Baumel, Julian J., Functional morphology of the tail apparatus of the pigeon (*Columba livia*) (rev.), 763
Bech, Claus, see Klaassen, Marcel, et al.
Bee-eater, European, 375-381
Behavior, activity patterns, 21, 30-31, 54-57, 94-100, 218-224, 227, 342, 343, 353-356, 495-497, 549, 550-551, 552, 556, 557, 661-662; age-related, 218-223, 375-381, 390-399, 590, 591, 592; agonistic, 1-6, 211, 365, 392, 471-472, 647, 648-649; anting, 163-164; antipredator, 152, 225, 226, 230-231, 497, 504, 507, 659-664, 738-741, 742, 747-749; breeding, 1-6, 56-57, 58, 59, 60, 61-62, 86-91, 107-115, 177-184, 209-216, 292-300, 332-335, 336-338, 358-365, 375-381, 389-400, 418-420, 433-437, 480-482, 483-485, 503-507, 549-557, 568-575, 584-592, 619-624, 625-632, 635-643, 700-704, 732-733, 734-737, 738-741; copulatory, 280, 281, 283; courtship, 5, 76, 298, 471; dispersal, 72-77, 589-590, 738-741; distraction, 742; dominance, 2, 127, 314, 316, 336-338, 490-491, 561; effect of environment on, 221-222, 223-224, 234-235, 236, 237, 389-400, 625-

- 632; effect of wing tags on, 619–624; feeding, 42–47, 67, 94–100, 118–123, 141–143, 150–153, 193, 198, 199, 218–224, 225–231, 253–255, 353–356, 418–420, 442–444, 467–470, 504–505, 506–507, 549, 550–551, 552, 556, 557, 629, 632, 700–704; feeding aggregation, 150–153, 225–227, 230–231, 406–407; flight, 141–143, 148–150, 223; food-begging, 506; foraging (see Behavior, feeding) homing, 148–150; incubation, 26–32, 433–437, 439–445, 477, 503–504, 726–727; individual distance, 72–77, 402–408; interspecific, 211–212, 213, 225–231, 339–340, 358–365, 645–650; maintenance, 221, 222–223, 504; mate replacement, 1–6, 292, 293, 294–295, 298–300, 378, 561, 622, 637, 640–641; mating preference, 86–91, 154–155, 155–156, 378, 391, 395, 418–420, 619, 621, 622, 623–624; movements in home range, 402–409; nest helper (see Communal breeding); nesting, 193–201, 209–216, 332–335, 391–393, 568–575, 625–632; nest-site tenacity, 102, 104, 391–392; pair, 1–6, 86–91, 378, 391, 392, 398, 399, 619, 621, 622, 623–624; parental care, 1–6, 18–24, 26–32, 114, 292–300, 360–361, 363, 365, 379, 380, 483–485, 490–491, 504–507, 629, 635–643, 659–644, 700–704, 732, 733; roosting, 151, 342, 343–344; sexual differences in, 72–77, 150–153, 292, 294–296, 299–300, 390, 468, 469, 470, 473, 483–485, 499–500, 659–664; siblicide, 520–521, 522, 523–524; site fidelity, 102, 104, 149, 377, 379, 380, 390, 397–398, 399, 489, 584–592, 623; social, 124–128, 332–335, 560–566; status signaling, 328; territorial, 3, 4, 57–58, 59, 60, 61–62, 102, 104, 107–115, 405, 407, 408, 560–566, 645–650, 733; thermoregulatory, 342, 343–344; vocal, 26, 32, 49–62, 114, 124–128, 227, 332–335, 339–340, 471–473, 498–500, 504–505, 505–506, 645–650, 659–662, 728–729, 748
- Behavior of Accipiter gentilis**, 227–228, 229–231; *A. nisus*, 225–231; *Aechmophorus clarkii*, 467–470; *A. occidentalis*, 467–470; *Aegolius acadicus*, 732–733; *Age-laius phoeniceus*, 107–115; *Aphelocoma ultramarina*, 124–128; *Bombycilla cedrorum*, 64–70; *Bonasa umbellus*, 72–77; *Branta canadensis moffitti*, 218–224, 439–445; *Bucephala albeola*, 568–575; *Burhinus oedicnemus*, 726, 727; *Buteo buteo*, 227–228, 229–231; *Calidris canutus*, 94–100; *C. himantopus*, 94–100; *C. mauri*, 94–100; *C. minutilla*, 94–100; *C. pusilla*, 94–100; *Casmerodus albus*, 625–632; *Catoptrophorus semipalmatus*, 94–100; *Charadrius semipalmatus*, 94–100; *C. wilsonia*, 94–100; *Circus cyaneus*, 227–228, 229–231, 353–356; *Columbina passerina*, 742; *Coturnix japonica*, 280, 281, 283; *Dendroica graciae*, 49–62; *Dolichonyx oryzivorus*, 584–592; *Egretta caerulea*, 625–632; *E. rufescens*, 94–100; *E. garzetta*, 292–300; *E. thula*, 94–100; *E. tricolor*, 94–100; *Eudocimus albus*, 625–632; *Eurypyga helias*, 503–507; *Falco columbarius*, 227–228, 229–231; *F. tinnunculus*, 549–557; *Fringilla coelebs*, 225–231; *F. montifringilla*, 225–231; *Himantopus mexicanus*, 94–100; *Hylocichla mustelina*, 402–409; *Icterus galbula*, 735–737; *Jacana jacana*, 1–6; *Lanius excubitor*, 418–420; *Larus californicus*, 102, 104; *L. delawarensis*, 619–624; *Leptotila rufaxilla*, 339–340; *L. verreauxi*, 339–340; *L. wellsi*, 330–340; *Limnodromus griseus*, 94–100; *Loxia leucoptera*, 483–485; *Merops apiaster*, 375–381; *Mimus polyglottos*, 659–664; *Molothrus ater*, 358–365; *Numenius phaeopus*, 94–100; *Nycticorax nycticorax*, 94–100; *N. violaceus*, 94–100; *Nymphicus hollandicus*, 86–91; *Oceanodroma leucorhoa*, 148–150, 332–335, 498–500; *O. monachus*, 471–473; *Oenanthe oenanthe*, 18–24; *Oxyura jamaicensis*, 42–47; *Parabuteo unicinctus*, 480–482; *Parus atricapillus*, 645–650; *P. gambeli*, 645–650; *P. major*, 490–491; *Passer domesticus*, 150–153; *Passerina cyanea*, 747–749; *Pelecanus erythrorhynchos*, 26–32; *Phalacrocorax penicillatus*, 389–400; *Philomachus pugnax*, 336–338; *Phoenicopterus ruber*, 94–100; *Phoeniculus purpureus*, 342, 343; *Pica pica*, 635–643; *Pluvialis squatarola*, 94–100; *Ptychoramphus aleuticus*, 495–497; *Sitta europaea*, 560–566; *Spizella arborea*, 118–123; *Synthliboramphus antiquus*, 433–437; *Tringa flavipes*, 94–100; *T. melanoleuca*, 94–100; *Upupa epops*, 726–727; *Vireo solitarius*, 477–479.
- Beletsky, Les D., Gordon H. Orian, and John C. Wingfield**, Relationships of steroid hormones and polygyny to territorial status, breeding experience, reproductive success in male Red-winged Blackbirds, 107–117
- Belgium**, 560–566
- Benkman, Craig W.**, Breeding opportunities, foraging rates, and parental care in White-winged Crossbills, 483–485
- Bevier, Louis R.**, review by, 349
- Binford, Laurence C.**, A distributional survey of the birds of the Mexican state of Oaxaca (rev.), 760–761
- Birkhead, T. R.**, see Clayton, N. S., and ———
- Blackbird, Red-winged**, 107–115, 358–365, 577–582; Yellow-headed, 577–582
- Blockstein, David E., and J. W. Hardy**, The Grenada Dove (*Leptotila wellsi*) is a distinct species, 339–340
- Bluebird, Mountain**, 451, 452, 459; Western, 452
- Boag, David A.**, review by, 756–757; see Murphy, Andrew J., and ———
- Bobolink**, 577–582, 584–592
- Bobwhite, Northern**, 492–494
- Boekelheide, Robert J., and David G. Ainley**, Age, resource availability, and breeding effort in Brandt's Cormorant, 389–401
- Boles, Walter E.**, Correct authorship of Pomatomidae, 341
- Bolivia**, 34–39
- Bollinger, Eric K., and Thomas A. Gavin**, The effects of site quality on breeding-site fidelity in Bobolinks, 584–594
- Bombycilla cedrorum**, 64–70
- Bonasa umbellus**, 72–77, 79
- Books**, Gregory G., see Eberhardt, Lester E., et al.
- Bortolotti, Gary R.**, Commentary: Sex ratios of fledgling Golden Eagles, 420–421

- Brambling, 225-231
- Branta canadensis moffitti*, 218-224, 439-445
- Breeding of (see also Behavior, breeding; Breeding season; Breeding success; Clutch size; Eggs; Incubation and incubation period; Nest and nest site) *Aegolius acadicus*, 732-733; *Agelaius phoeniceus*, 107-115; *Bucephala albeola*, 568-575; *Calcarus lapponicus*, 738-741; *Casmerodus albus*, 625-632; *Cephus columba*, 193-201; *Chen caerulescens caerulescens*, 177-184; *Coturnix japonica*, 279-283, 286-289; *Dolichonyx oryzivorus*, 584-592; *Egretta caerulea*, 625-632; *E. garzetta*, 292-300; *E. thula*, 625-632; *E. tricolor*, 625-632; *Empidonax minimus*, 269-276; *Eudocimus albus*, 625-643; *Eurypyga helias*, 503-507; *Falco tinnunculus*, 549-557; *Jacana jacana*, 1-6; *Lanius excubitor*, 418-420; *Larus delawarensis*, 619-624; *Loxia leucoptera*, 483-485; *Melanerpes carolinus*, 209-216; *M. erythrocephalus*, 209-216; *Merops apiaster*, 375-381; *Molothrus ater*, 358-365; *Nymphicus hollandicus*, 86-91; *Oceanodroma leuconota*, 332-335; *Parabuteo unicinctus*, 480-482; *Phalacrocorax penicillatus*, 389-400; *Pica pica*, 635-643; *Sturnus vulgaris*, 128-134, 209-216; *Synthliboramphus antiquus*, 433-437, 653-658; *Uria lomvia*, 433-437; *Vireo solitarius*, 476-479
- Breeding season of *Egretta garzetta*, 297; *Lanius excubitor*, 419-420; *Melanerpes carolinus*, 210-211, 212; *M. erythrocephalus*, 211, 213; *Phalacrocorax penicillatus*, 391, 392; *Sturnus vulgaris*, 210-211; *Vireo solitarius*, 477
- Breeding success, 86-91, 107, 111-113, 115, 132, 185, 187, 189, 211, 214, 215, 272-273, 274-275, 281-283, 335, 360, 361-362, 363, 364-365, 378, 379, 380-381, 393-395, 418-420, 435-436, 477-479, 517-518, 518-519, 568-575, 584-592, 622, 623, 625-632, 635-643, 703-704, 740
- Breeding success of *Aegolius acadicus*, 732, 733; *Agelaius phoeniceus*, 107, 111-113, 115, 363; *Bucephala albeola*, 568-575; *Casmerodus albus*, 625-632; *Coturnix japonica*, 281-283; *Dendroica petechia*, 363; *Dolichonyx oryzivorus*, 584-592; *Egretta caerulea*, 625-632; *E. thula*, 625-632; *E. tricolor*, 625-632; *Empidonax minimus*, 272-273, 274-275; *Eudocimus albus*, 625-632; *Haematopus palliatus*, 185, 187, 189; *Lanius excubitor*, 418-420; *Larus delawarensis*, 622, 623; *Melanerpes carolinus*, 214; *M. erythrocephalus*, 211, 215; *Merops apiaster*, 378, 379, 380-381; *Molothrus ater*, 360, 361-362, 364-365; *Nymphicus hollandicus*, 86-91; *Oceanodroma leuconota*, 335; *Phalacrocorax penicillatus*, 393-395; *Pica pica*, 635-643; *Picoides borealis*, 517-518, 518-519; *Sturnus vulgaris*, 132; *Synthliboramphus antiquus*, 435-436; *Vireo solitarius*, 477-479
- Breitwisch, Randall, Natasha Gottlieb, and Julia Zaias, Behavioral differences in nest visits between male and female Northern Mockingbirds, 659-665; and Joann Hudak, Sex differences in risk-taking behavior in foraging flocks of House Sparrows, 150-153
- Briggs, Kenneth T., W. Breck Tyler, David B. Lewis, and David R. Carlson, Bird communities at sea off California: 1975 to 1983 (rev.), 170-171
- Brink, Karen L., see Roby, Daniel D., et al.
- Briskie, James V., and Spencer G. Sealy, Determination of clutch size in the Least Flycatcher, 269-278
- British Columbia, 193-201, 279-283, 286-289, 433-437, 568-575, 653-658, 733
- Brolga, 595-601, 603-613, 616-618
- Brown, Jerram L., and Eric G. Horvath, Geographic variation of group size, ontogeny, rattle calls, and body size in *Aphelocoma ultramarina*, 124-128; see Ito, Yosiaki, et al.
- Browning, M. Ralph, The correct citation and spelling of *Ptiliogonys* and type locality of *Ptiliogonys cinereus*, 743-746
- Brush, Alan H., reviews by 350, 530-531, 763-764; see Hudon, Jocelyn, et al.
- Bubo bubo*, 539-546
- Bucephala albeola*, 568-575
- Bufflehead, 568-575
- Bugeranus carunculatus*, 595-601, 603-613, 616-618
- Bunting, Indigo, 747-749
- Burger, Alan E., review by, 171-172
- Burger, Joanna, see Lauro, Brook, and —
- Burhinus oedicnemus*, 726, 727
- Buteo buteo*, 227-230, 539-546
- Buteo jamaicensis*, 451, 452
- Butler, Ronald G., see Place, Allen R., et al.
- Buttron, Deborah P., see Nuechterlein, Gary L., and —
- Butyev, V. T., see Ilyichev, V. D., et al.
- Buzzard, Common, 227-230, 539-546
- Calidris lapponicus*, 738-741
- Calder, William A., III, and Elinor G. Jones, Implications of recapture data for migration of the Rufous Hummingbird (*Selasphorus rufus*) in the Rocky Mountains, 488-489
- Calidris alba*, 326-330, 474-476
- C. canutus*, 94-100, 326-330
- C. himantopus*, 94-100, 326-330
- C. mauri*, 94-100
- C. minutilla*, 94-100
- C. pusilla*, 94-100, 474-476
- California, 86-91, 102-105, 326-330, 353-356, 389-400, 447-459, 495-497
- Calls (see Vocalizations)
- Calonectris diomedea*, 411-416, 417
- Cannings, Richard J., see Marks, Jeffrey S., et al.
- Capparella, A. P., see Hudon, Jocelyn, et al.
- Cardinal, Northern, 324-325
- Cardinalis cardinalis*, 324-325
- Carey, Cynthia, see Ligon, J. David, et al.
- Carlson, David R., see Briggs, Kenneth T., et al.
- Carpodacus cassini*, 451-452
- C. purpureus*, 452

- Carpornis cucullatus*, 426, 428
 Carr, David E., see Hill, Geoffrey E., and ———
 Carroll, Janet R., see Andrle, Robert F., and ———
 Carter, Harry R., review by, 170–171
Casmerodus albus, 625–632
 Castro, Gonzalo, and J. P. Myers, Flight range estimates for shorebirds, 474–476
Catoptrophorus semipalmatus, 94–100, 474–476
Cathartes melambrotus, 539–546
C. guttatus, 452
 Census methods, 130–131, 134, 367–372, 480–482
Cephus columba, 193–201
Certhia americana, 451–452
 Chaffinch, 225–231
Charadrius, 154–155, 155–156
Charadrius semipalmatus, 94–100
C. wilsonia, 94–100
Chen caerulescens caerulescens, 177–184
 Cheng, Kimberly M., Andrew R. Hickman, and Cathleen R. Nichols, Role of the proctodeal gland foam of male Japanese Quail in natural copulations, 279–285; Roy F. McIntyre, and Andrew R. Hickman, Proctodeal gland foam enhances competitive fertilization in domestic Japanese Quail, 286–291
 Chickadee, Black-capped, 259–266, 268, 645–650, 652; Mountain, 259–266, 268, 451, 452, 459, 645–650, 652
Chlorostilbon canivetii, 203–207
Chordeiles minor, 451, 452, 539–546
Cinclocerthia gutturalis, 249–256, 258
C. ruficauda, 249–256, 258
Circus cyaneus, 225, 227–230, 353–356
 Clayton, N. S., and T. R. Birkhead, Commentary: Consistency in the scientific name of the Zebra Finch, 750
 Clench, Mary H., and J. William Hardy, In Memoriam: Oliver L. Austin Jr., 706–709
 Clutch size of *Aegialius phoeniceus*, 362, 363; *Branta canadensis moffitti*, 441–442; *Bucephala albeola*, 569–572, 574; *Dendroica petechia*, 362–363; *Empidonax minimus*, 269–276; *Falco tinnunculus*, 550, 552–553, 557; *Melanerpes carolinus*, 213; *M. erythrocephalus*, 214–215; *Merops apiaster*, 378–379, 380; *Passer domesticus*, 730–731; *Phalacrocorax penicillatus*, 394, 395, 396, 397, 398; *Pica pica*, 635–643; *Upupa epops*, 726; *Vireo solitarius*, 477
Coccothraustes affinis, 728–729
C. icterioides, 728–729
C. vespertinus, 451, 452
 Cockatiel, 86–91
Colaptes auratus, 452
Colinus virginianus, 492–494
 Collar, N. J., and P. Andrew, Birds to watch (rev.), 526–527
 Collopy, Michael W., see Edwards, Thomas C., Jr., and ———; see Frederick, Peter C., and ———
Columba fasciata, 452
 Colombia, 34–39
 Colorado, 476–479, 488–489
Columbina passerina, 742
 Colwell, Mark A., review by, 762–763
 Commentaries, Aerial agility and the evolution of reversed sexual dimorphism (RSD) in shorebirds, see Mueller, Helmut C.; Response: Evolution of sexual size dimorphism, see Jehl, Joseph R., Jr., and Bertram G. Murray Jr.; The study of geographic variation, see Zink, Robert M.; Response to R. M. Zink, see Rising, J. D.; Response to P. R. Ehrlich, D. S. Dobkin, and D. Wheye, see Potter, Eloise F.; Cavity roosting, philopatry, and cooperative breeding in the Green Woodhoopoe may not reflect a physiological trait, see Ward, David, et al.; Response to Ward et al., see Ligon, J. David, et al.; A new classification of the living birds of the world, see Mayr, Ernst; Response to E. Mayr, see Sibley, Charles G.; see Monroe, Burt L., Jr.; Statistical inference from color-banding data, see Hill, Geoffrey E., and David E. Carr; Response to Hill and Carr, see Hagan, John M., and J. Michael Reed; Sex ratios of fledgling Golden Eagles, see Bortolotti, Gary R.; Sex ratios of fledgling Golden Eagles and jackrabbit densities, see Arnold, Todd W.; Response to G. R. Bortolotti and T. W. Arnold, see Edwards, Thomas C., Jr., and Michael W. Collopy; Consistency in the scientific name of the Zebra Finch, see Clayton, N. S., and T. R. Birkhead
 Communal breeding, 700–704
 Communal breeding in *Cephus columba*, 193–201; *Merops apiaster*, 378, 381; *Parabuteo unicinctus*, 480–482
Conioptilon mcilhennyi, 426, 428
 Connecticut, 367–372
Contopus borealis, 451–452
 Cooke, Fred, see Quinn, Thomas W., et al.
 Cooperative breeding (see Communal breeding)
 Cormorant, Brandt's, 389–400
Corvus caurinus, 197–198
C. corax, 452
 Costa Rica, 203–207, 503–507
Coturnix japonica, 279–283, 286–289
 Cowbird, Brown-headed, 358–365, 452, 476–479, 577–582, 734–737
 Crane, Black-crowned, 595–601, 603–613, 616–618; Common, 595–601, 603–613, 616–618; Demoiselle, 595–601, 603–613, 616–618; Gray Crowned, 595–601, 603–613, 616–618; Hooded, 595–601, 603–613, 616–618; Red-crowned, 595–601, 603–613, 616–618; Sandhill, 595–601, 603–613, 616–618; Sarus, 595–601, 603–613, 616–618; Siberian, 595–601, 603–613, 616–618; Stanley, 595–601, 603–613, 616–618; Wattled, 595–601, 603–613, 616–618; Whistled, 595–601, 603–613, 616–618; Whooping, 595–601, 603–613, 616–618
 Creeper, Brown, 451–452
 Crossbill, White-winged, 483–485
 Crow, Northwestern, 197–198

- Croxall, J. P. (Ed.), Seabirds: feeding ecology and role in marine ecosystems (rev.), 171–172
- Cruz, Alexander, see Marvil, Rebecca E., and ——
- Cruz, Justine B., review by, 761
- Crypturellus undulatus*, 539–546
- Culicicapa ceylonensis*, 728–729
- Curlew, Stone, 726–727
- Cuthill, Innes C., see Lohrer, Fred E., et al.
- Cyanocitta stelleri*, 451–452
- Cygnus buccinator*, 539–546
- Daan, Serge, see Meijer, Theo, et al.
- Davies, J. Chris, see Quinn, Thomas W., et al.
- Dawson, James W., and R. William Mannan, A comparison of two methods of estimating breeding group size in Harris' Hawks, 480–483
- Demong, Natalie J., see Emlen, Stephen T., et al.
- Dendragapus obscurus*, 451, 452
- Dendroica coronata*, 451–452
- D. graciae*, 49–62
- D. occidentalis*, 451–452
- D. petechia*, 358–365, 451, 452, 456
- D. striata*, 8–15, 17
- de Queiroz, Kevin, see Warheit, Kenneth I., et al.
- de Santo, Toni, see Pennycuick, C. J., and ——
- Dicrurus leucophaeus*, 728–729
- Diseases and parasites of *Larus californicus*, 102–103, 104
- Distribution, spatial, 193–201
- Distribution of *Dendroica striata*, 9–11
- Dittmann, Donna L., Robert M. Zink, and John A. Gerwin, Evolutionary genetics of phalaropes, 326–331
- Dolichonyx oryzivorus*, 577–582, 584–592
- Doremus, John H., see Marks, Jeffrey S., et al.
- Dove, Gray-fronted, 339–340; Grenada, 339–340
- Dowitcher, Long-billed, 326–330; Short-billed, 94–100
- Dryocopus pileatus*, 452
- Duck, Ruddy, 42–47
- Dufy, Alfred M., Jr., An effective anti-estrogen for feral birds, 485–487
- Duke, G. E., A. R. Place, and B. Jones, Gastric emptying and gastrointestinal motility in Leach's Storm-Petrel chicks (*Oceanodroma leucorhoa*), 80–85
- Dunn, Peter O., and Susan J. Hannon, Evidence for obligate male parental care in Black-billed Magpies, 635–644
- Eagle, Golden, 520–521, 521–522, 523–524
- Eagle-Owl, Great, 539–546
- Eberhardt, Lester E., Gregory G. Books, Robert G. Anthony, and William H. Rickard, Activity budgets of Canada Geese during brood rearing, 218–224
- Ecology, bird species richness, 447–459; breeding communities, 447–459, 625–632; competition, 146–147, 209–216, 254, 255; dispersal, 75, 76–77; habitat partitioning, 259, 264–266, 467–470; habitat selection, 75, 76–77, 185–191, 193–201, 218, 222, 224, 256, 402, 406–407, 408, 409, 447–459, 562–563, 565–566; predator-prey relationships, 355–356, 521–522, 523, 632; territoriality, 102, 103–105, 259–266, 562–563, 565–566
- Ecology of *Accipiter cooperii*, 451, 452; *A. gentilis*, 452; *A. striatus*, 451, 452; *Aechmophorus clarkii*, 467–470; *A. occidentalis*, 467–470; *Bombycilla cedrorum*, 70; *Bonasa umbellus*, 75, 76–77; *Branta canadensis moffitti*, 218, 222–224; *Buteo jamaicensis*, 451, 452; *Calidris canutus*, 94–100; *C. himantopus*, 94–100; *C. minutilla*, 94–100; *C. pusilla*, 94–100; *Carpodacus cassini*, 451–452; *C. purpureus*, 452; *Casmerodus albus*, 625–632; *Cathartes guttatus*, 452; *Catoptrophorus semipalmatus*, 94–100; *Certhia americana*, 451–452; *Chadradius semipalmatus*, 94–100; *C. wilsonia*, 94–100; *Chordeiles minor*, 451, 452; *Cinclocerthia ruficauda*, 254; *Circus cyaneus*, 355–356; *Coccothraustes vespertinus*, 451, 452; *Colaptes auratus*, 452; *Columba fasciata*, 452; *Contopus borealis*, 451–452; *Corvus corax*, 452; *Cyanocitta stelleri*, 451–452; *Dendragapus obscurus*, 451, 452; *Dendroica coronata*, 451–452; *D. occidentalis*, 451–452; *D. petechia*, 451, 452, 456; *Dolichonyx oryzivorus*, 584–592; *Dryocopus pileatus*, 452; *Egretta caerulea*, 625–632; *E. rufescens*, 94–100; *E. thula*, 94–100, 625–632; *E. tricolor*, 94–100, 625–632; *Empidonax hammondi*, 451, 452; *E. oberholseri*, 451, 452; *Eudocimus albus*, 625–632; *Falco sparverius*, 451, 452; *Geospiza difficilis*, 146–147; *G. fuliginosa*, 146–147; *Haematopus palliatus*, 185–191; *Himantopus mexicanus*, 94–100; *Hylocichla mustelina*, 402–409; *H. ustulata*, 451, 452; *Juncos hyemalis*, 451, 452; *Larus californicus*, 102, 103–105; *Limnodromus griseus*, 94–100; *Melanerpes carolinus*, 209–216; *M. erythrocephalus*, 209–216; *Molothrus ater*, 452; *Myadestes townsendi*, 452; *Numenius phaeopus*, 94–100; *Nycticorax nycticorax*, 94–100; *N. violaceus*, 94–100; *Opornis tolmiei*, 451, 452, 456; *Oreortyx pictus*, 452; *Parus atricapillus*, 259–266; *P. gambeli*, 259–266, 451, 452, 459; *Passerella iliaca*, 451, 452, 456; *Phalacrocorax penicillatus*, 395–396, 397–400; *Pheucticus melanocephalus*, 452; *Phoenicopterus ruber*, 94–100; *Picoides albolarvatus*, 451, 452, 459; *P. villosus*, 451, 452, 459; *Pinicola enucleator*, 452; *Pipilo chlorurus*, 451, 452, 456; *P. erythrorththalmus*, 451, 452, 456; *Piranga ludoviciana*, 451–452; *Ramphocinclus brachyurus*, 254; *Regulus satrapa*, 451–452; *Sialia currucoides*, 451, 452, 459; *S. mexicana*, 452; *Sitta canadensis*, 452; *S. europaea*, 562–563, 565–566; *Sphyrapicus ruber*, 452; *S. thyroideus*, 452; *Spizella passerina*, 452; *Stellula calliope*, 452; *Strix occidentalis*, 451, 452; *Sturnus vulgaris*, 209–216; *Tringa flavipes*, 94–100; *Tringa melanoleuca*, 94–100; *Troglodytes troglodytes*, 451, 452; *Turdus migratorius*, 452; *Vermivora ruficapilla*, 452; *Vireo gilvus*, 451–452; *V. solitarius*, 452; *Wilsonia pusilla*, 452
- Ectoparasitism (see Diseases and parasites)
- Edwards, Thomas C., Jr., and Michael W. Collopy, Commentary: Response to G. R. Bortolotti and T. W. Arnold, 523–524

- Eggs, composition, 549, 553, 577–582; density, specific gravity, and volume, 577–582; laying, 88, 89, 286–289, 297, 359–360, 361, 364, 365, 391, 476–479, 549, 552–553, 554–557, 638, 730–731; temperature, 26–32
- Eggs, descriptions and measurements, of *Agelaius phoeniceus*, 578–582; *Dolichonyx oryzivorus*, 578–582; *Falco tinnunculus*, 552, 553, 554; *Molothrus ater*, 577–582, 734–737; *Pelecanus erythrorhynchos*, 26–32; *Xanthocephalus xanthocephalus*, 578–582
- Egret, Great, 625–532; Little, 292–300; Reddish, 94–100; Snowy, 94–100, 625–632
- Egretta caerulea*, 625–632
- E. garzetta*, 292–300
- E. rufescens*, 94–100
- E. thula*, 94–100, 625–632
- E. tricolor*, 94–100, 625–632
- Ellison, Walter G., review by, 761–762
- Ellsworth, Darrell L., John L. Roseberry, and W. D. Klimstra, Genetic structure and gene flow in the Northern Bobwhite, 492–495
- Emlen, Douglas J., see Emlen, Stephen T., et al.
- Emlen, Stephen T., Natalie J. Demong, and Douglas J. Emlen, Experimental induction of infanticide in female Wattled Jacanas, 1–7
- Emms, S. K., and N. A. M. Verbeek, Significance of the pattern of nest distribution in the Pigeon Guillemot (*Cephus columba*), 193–202
- Empidonax hammondi*, 451, 452
- E. minimus*, 269–276
- E. oberholseri*, 451, 452
- Endangered species, 91, 334–335, 339–340, 367–372
- England, 336–338
- Environment, influence on *Branta canadensis moffitti*, 221–222, 223–224; *Calidris canutus*, 94–100; *C. himantopus*, 94–100; *C. mauri*, 94–100; *C. minutilla*, 94–100; *C. pusilla*, 94–100; *Casmerodus albus*, 625–632; *Catoptrophorus semipalmatus*, 94–100; *Cephus columba*, 199–200; *Charadrius semipalmatus*, 94–100; *C. wilsonia*, 94–100; *Egretta caerulea*, 625–632; *E. rufescens*, 94–100; *E. thula*, 94–100, 625–632; *E. tricolor*, 94–100, 625–632; *Eudocimus albus*, 625–632; *Himantopus mexicanus*, 94–100; *Limnodromus griseus*, 94–100; *Numenius phaeopus*, 94–100; *Nycticorax nycticorax*, 94–100; *N. violaceus*, 94–100; *Passerella iliaca schistacea*, 161–162; *Phalacrocorax penicillatus*, 389–400; *Philonotus pugnax*, 338; *Phoenicopterus ruber*, 94–100; *Picoides pubescens*, 315, 318–319; *Pluvialis squatarola*, 94–100; *Spizella arborea*, 118–123; *Sterna paradisaea*, 246–247; *Tringa flavipes*, 94–100; *T. melanoleuca*, 94–100
- Eriothacus cyaneurus*, 728–729
- Eudocimus albus*, 141–143, 625–632
- Eurypyga helias*, 503–507
- Evans, Roger M., Egg temperatures and parental behavior during the transition from incubation to brooding in the American White Pelican, 26–33
- Evolution of communal breeding, 200–201; genetic variation, 157–159, 160–163; reversed sexual dimorphism, 154–155, 155–156
- Evolution of *Cinclocerthia gutturalis*, 255; *C. ruficauda*, 255; *Geospiza difficilis*, 147; *G. fuliginosa*, 147; *Passerella iliaca schistacea*, 157–159, 160–163; *Phalaropus fulicaria*, 326, 329–330; *P. lobatus*, 326, 329–330; *P. tricolor*, 326, 329–330; *Ramphocinclus brachyurus*, 255
- Extra-pair fertilization, 128–134
- Falco columbarius*, 225, 227–230
- Falco sparverius*, 451, 452
- Falco tinnunculus*, 539–546, 549–557
- Falcon, Barred Forest, 407, 408
- Farley, Greg H., see Ligon, J. David, et al.
- Feathers (see Molts and Plumages)
- Ficedula hypoleuca*, 501–502
- Finch, Cassin's, 451–452; Purple, 452; Zebra, 750
- Flamingo, Greater, 94–100
- Fledging, success of (see Breeding success)
- Flegg, Jim, see Hosking, Eric, et al.
- Flicker, Northern, 452
- Florida, 625–632, 659–664, 742
- Flycatcher, Dusky, 451, 452; Hammond's 451, 452; Least, 269–276; Olive-sided, 451–452; Pied, 501–502; Spotted, 501–502
- Fogden, Michael P. L., see Lyon, Bruce E., and
-
- Food and feeding habits of (see also Behavior, feeding) *Accipiter gentilis*, 225, 227–230; *A. nisus*, 225, 227–230; *Aechmophorus clarkii*, 467–470; *A. occidentalis*, 467–470; *Aegolius acadicus*, 732, 733; *Amaurornis saucerottei*, 203–207; *Bombycilla cedrorum*, 64–70; *Branta canadensis moffitti*, 218–224, 442–444; *Buteo buteo*, 225, 227–230; *Calidris canutus*, 94–100; *C. himantopus*, 94–100; *C. mauri*, 94–100; *C. minutilla*, 94–100; *C. pusilla*, 94–100; *Cardinalis cardinalis*, 325; *Catoptrophorus semipalmatus*, 94–100; *Charadrius semipalmatus*, 94–100; *Chlorostilbon canivetii*, 203–207; *Cinclocerthia ruficauda*, 253; *Circus cyaneus*, 225, 227–230, 353–356; *Egretta rufescens*, 94–100; *E. thula*, 94–100; *E. tricolor*, 94–100; *Empidonax minimus*, 270, 272, 275; *Eudocimus albus*, 141–143, 629, 632; *Eurypyga helias*, 504–505, 506–507; *Falco columbarius*, 225, 227–230; *F. tinnunculus*, 549, 550–551, 552, 556, 557; *Fringilla coelebs*, 225–231; *F. montifringilla*, 225–231; *Geospiza difficilis*, 146–147; *G. fuliginosa*, 146–147; *Himantopus mexicanus*, 94–100; *Junco hyemalis*, 325; *Lanius excubitor*, 418–420; *Limnodromus griseus*, 94–100; *Mimus polyglottos*, 662–663; *Numenius phaeopus*, 94–100; *Nycticorax nycticorax*, 94–100; *N. violaceus*, 94–100; *Oxyura jamaicensis*, 42–47; *Passer domesticus*, 150–153; *Phoenicopterus ruber*, 94–100; *Pluvialis squatarola*, 94–100; *Ramphocinclus brachyurus*, 254–255; *Spizella arborea*, 118–123, 325; *Sturnus vulgaris*, 675–684; *Tringa flavipes*, 94–100; *T. melanoleuca*, 94–100; *Turdus migratorius*, 675–684; *Zonotrichia querula*, 325

- Foraging (see Food and feeding habits; Behavior, feeding)
- Ford, Hugh A., see Lohrer, Fred E., et al.
- France, 375-381
- Frederick, Peter C., and Michael W. Collopy, Nesting success of five ciconiiform species in relation to water conditions in the Florida Everglades, 625-634
- Freethy, Ron, Auks: an ornithologist's guide (rev.), 525-526
- Fregata aquila*, 383-388
- F. ariel*, 383-388
- F. magnificens*, 383-388
- F. minor*, 383-388
- Fringilla coelebs*, 225-231
- F. montifringilla*, 225-231
- Fry, C. Hilary, Stuart Keith, and Emil K. Urban (Eds.), *The birds of Africa. Vol. 3, Parrots to woodpeckers* (rev.), 752-754
- Fujioka, Masahiro, Mate and nestling desertion in colonial Little Egrets, 292-302
- Fuller, Errol, Extinct birds (rev.), 346-347
- Gaines, Sarah, Comparisons of patch-use models for wintering American Tree Sparrows, 118-123
- Galápagos, 144-147
- Gallus gallus*, 539-546
- Garrido, Orlando H., see Reynard, George B., and
- Garrulax lineatus*, 728-729
- Gaston, Anthony J., and Ian L. Jones, The relative importance of stress and programmed anorexia in determining mass loss by incubating Ancient Murrelets, 653-658; and David W. Powell, Natural incubation, egg neglect, and hatchability in the Ancient Murrelet, 433-438
- Gauthier, Gilles, The effect of experience and timing on reproductive performance in Buffleheads, 568-576
- Gavin, Thomas A., see Bollinger, Eric, K., and
- Genetic variation in (see Genetics, genetic variation)
- Genetics, 129, 131-133, 177-184, 326-330, 336, 337-338, 411-416, 417, 423-425, 426, 429-431, 492-494, 508-512, 513-515, 515-516, 539-546, 595-601, 603-613, 616-618; DNA-DNA hybridization, 508-512, 513-515, 515-516, 595, 598, 599, 600, 603-613, 616-618; gene flow, 157-159, 160-162, 411-416, 492-494; genetic variation, 131, 157-159, 160-163, 326-330, 411-416, 492-494
- Genetics of *Ampelion rubrocristata*, 423-425, 426, 429-431; *A. rufaxilla*, 423-425, 426, 429-431; *A. sclateri*, 423-425, 426, 429-431; *A. strelzmanni*, 423-425, 426, 429-431; *Anthropoides paradisea*, 595-601, 603-613, 616-618; *A. virgo*, 595-601, 603-613, 616-618; *Aramus guarauna*, 603-609, 612, 616-618; *Balearica pavonina*, 595-601, 603-613, 616-618; *B. regulorum*, 595-601, 603-613, 616-618; *Bubo bubo*, 539-546; *Bugeranus carunculatus*, 595-601, 603-613; 616-618; *Buteo buteo*, 539-546; *Calidris alba*, 326-330; *C. canutus*, 326-330; *C. himantopus*, 326-330; *Calonectris diomedea*, 411-416, 417; *Cathartes melambrotus*, 539-546; *Chen caerulescens caerulescens*, 177-184; *Chordeiles minor*, 539-546; *Colinus virginianus*, 492-494; *Crypturellus undulatus*, 539-546; *Cygnus buccinator*, 539-546; *Gallus gallus*, 539-546; *Geothlypis trichas*, 599; *Gruidae*, 595-601, 603-613, 616-618; *Grus americana*, 595-601, 603-613, 616-618; *G. antigone*, 595-601, 603-613, 616-618; *G. canadensis*, 595-601, 603-613, 616-618; *G. grus*, 595-601, 603-613, 616-618; *G. japonensis*, 595-601, 603-613, 616-618; *G. leucogeranus*, 595-601, 603-613, 616-618; *G. monacha*, 595-601, 603-613, 616-618; *G. nigricollis*, 603-613, 616-618; *G. rubicunda*, 595-601, 603-613, 616-618; *G. vipio*, 595-601, 603-613, 616-618; *Larus californicus*, 326-330; *Limnodromus scolopaceus*, 326-330; *Mycteria americana*, 539-546; *Passer domesticus*, 539-546, 598; *Passerella iliaca schistacea*, 157-159, 160-163; *Phalaropus fulicaria*, 326-330; *P. lobatus*, 326-330; *P. tricolor*, 326-330; *Philomachus pugnax*, 336, 337, 338; *Phytotoma raimondii*, 423-425, 426, 429-431; *P. rara*, 423-425, 426, 429-431; *P. rutila*, 423-425, 426, 429-431; *Sturnus vulgaris*, 129, 131-133; *Tringa melanoleuca*, 326-330
- Geographic variation in molts and plumages, 251, 253, 254, 255
- Geographic variation in *Aphelocoma ultramarina*, 124-128; *Calonectris diomedea*, 411-416; *Cinclocerthia gutturalis*, 249-256; *C. ruficauda*, 249-256; *Geospiza difficilis*, 145-147; *G. fuliginosa*, 145-147; *Passerella iliaca schistacea*, 157-159, 160-163; *Ramphocinclus brachyurus*, 249-256
- Geospiza difficilis*, 144-147
- G. fuliginosa*, 144-147
- Geothlypis trichas*, 599
- Geroudet, Paul, Les oiseaux du lac Léman (rev.), 346
- Gerrard, Jon M., and Gary R. Bortolotti, The Bald Eagle: haunts and habits of a wilderness monarch (rev.), 347
- Gerwin, John A., see Dittmann, Donna L., et al.
- Giant-Petrel, Antarctic, 303-312
- Gibraltar, 233-237
- Glaucidium brasiliense*, 406, 407, 408
- Good, David A., see Warheit, Kenneth I., et al.
- Goose, Canada, 218-224, 439-445; Lesser Snow, 177-184
- Goshawk, Northern, 227-230, 452
- Gottlieb, Natasha, see Breitwisch, Randall, et al.
- Grau, C. R., see Yamamoto, J. T., et al.
- Grebe, Clark's 467-470; Western, 467-470
- Grenada, 339-340
- Grimes, L. G., The birds of Ghana (rev.), 173-174
- Grosbeak, Black-headed, 452; Evening, 451, 452; Pine, 452
- Ground-Dove, Common, 742
- Grouse, Blue, 451, 452; Ruffed, 72-77, 79
- Growth and development of *Agelaius phoeniceus*, 364; *Aphelocoma ultramarina*, 124-218; *Cephus columba*,

- 198–199; *Dendroica petechia*, 364; *Empidonax minimus*, 271, 273, 274; *Eurypyga helias*, 504–505; *Molothrus ater*, 360, 361; *Picoides pubescens*, 316–317; *Sterna paradisaea*, 242, 243, 245, 246; *Vireo solitarius*, 477–478
- Grubb, Thomas C., Jr., Ptilochronology: feather growth bars as indicators of nutritional status, 314–320
- Gruidae, 595–601, 603–613, 616–618
- Grus americana*, 595–601, 603–613, 616–618
- G. antigone*, 595–601, 603–613, 616–618
- G. canadensis*, 595–601, 603–613, 616–618
- G. grus*, 595–601, 603–613, 616–618
- G. japonensis*, 595–601, 603–613, 616–618
- G. leucogeranus*, 595–601, 603–613, 616–618
- G. monacha*, 595–601, 603–613, 616–618
- G. nigricollis*, 603–613, 616–618
- G. rubicunda*, 595–601, 603–613, 616–618
- G. vipio*, 595–601, 603–613, 616–618
- Guillemot, Pigeon, 193–201
- Gull, California, 102–105, 326–330; Ring-billed, 619–624; Western, 496–497
- Guttman, Sheldon I., see Ingold, James L., et al.
- Habitat of *Agelaius phoeniceus*, 109; *Bonasa umbellus*, 75, 76–77, 79; *Branta canadensis moffitti*, 218, 222–224; *Cephus columba*, 193–201; *Cinclocerthia ruficauda*, 254; *C. gutturalis*, 254; *Haematopus palliatus*, 185–191; *Hylocichla mustelina*, 402, 405–406, 408–409; *Larus californicus*, 102, 103–104; *Parus atricapillus*, 260, 264–266, 268; *P. gambeli*, 260, 264–266, 268; *Ramphocinclus brachyurus*, 254, 255
- Haematopus palliatus*, 185–195
- Hagan, John M., and J. Michael Reed, Commentary: Response to Hill and Carr, 518–520
- Hailman, Jack, P., Common Ground-Dove's injury feigning distracts Florida Scrub Jay, 742; review by, 751–752
- Hainsworth, F. Reed, Evaluating models of crop emptying in hummingbirds, 724–726
- Hannon, Susan J., see Dunn, Peter O., and —
- Hardy, J. William, see Clench, Mary H., and —
- Harrier, Hen, 225, 227–230; Northern, 353–356
- Hatching, success of (see Breeding success)
- Hatching, timing of, 295, 378, 379, 380, 436, 437, 504, 619, 621, 622, 623, 732
- Hawk, Cooper's, 451, 452; Harris', 480–482; Red-tailed, 451, 452; Sharp-shinned, 451, 452
- Heron, Little Blue, 625–632; Tricolored, 94–100, 625–632
- Hickman, Andrew R., see Cheng, Kimberly M., et al.
- Hilgerloh, Gudrun, Autumn migration of trans-Saharan migrating passerines in the Straits of Gibraltar, 233–239; Orientation of trans-Saharan passerine migrants in southwestern Spain, 501–502
- Hill, Brad G., and M. Ross Lein, Territory overlap and habitat use of sympatric chickadees, 259–268; and M. Ross Lein, Natural and simulated encounters between sympatric Black-capped Chickadees and Mountain Chickadees, 645–652
- Hill, David, and Peter Robertson, The pheasant: ecology, management and conservation (rev.), 757–758
- Hill, Geoffrey E., and David E. Carr, Commentary: Statistical inference from color-banding data, 517–518
- Hill, Martha, Bruno Liljefors: the peerless eye (rev.), 167–168
- Himantopus mexicanus*, 94–100
- Hoglund, Jacob, and Arne Lundberg, Plumage color correlates with body size in the Ruff (*Philomachus pugnax*), 336–338
- Honeycutt, Rodney L., see Mindell, David P., and —
- Hoopoe, 726–727
- Horvath, Eric G., see Brown, Jerram L., and —
- Hosking, David, see Hosking, Eric, and —
- Hosking, Eric, and David Hosking with Jim Flegg, Eric Hosking's birds of prey of the world (rev.), 349–350
- Houston, C. Stuart, In memoriam: Betty Carnes, 140; review by, 754–756
- Hubbard, John P., review by, 760–761
- Hudak, Joann, see Breitwisch, Randall, and —
- Hudon, Jocelyn, A. P. Capparella, and Alan H. Brush, Plumage pigment differences in manakins of the *Pipra erythrocephala* superspecies, 34–41
- Hudson, Peter J., and Michael R. W. Rand (Eds.), Ecology and management of gamebirds (rev.), 756–757
- Hummingbird, Calliope, 452; Rufous, 488–489, 724–725
- Hunter, Laurie A., review by, 346–347
- Hunter, Malcolm L., Jr., Himalayan birds face uphill while singing, 728–729
- Hunting, Charles E., see Pierson, Elizabeth Cary, et al.
- Hybrids and hybridization of *Aphelocoma ultramarina* × *A. coerulescens*, 128
- Hylocichla mustelina*, 402–409
- H. ustulata*, 451, 452
- Hypsipetes madagascariensis*, 728–729
- Ibis, White, 141–143, 625–632
- Icterus galbula*, 735–737
- Idaho, 732–733
- Illinois, 492–494
- Ilyichev, V. D., V. T. Butyev, and V. M. Konstantinov, Ptitsy Moskvy i podmoskovya [Birds of Moscow and vicinity] (rev.), 174
- Incubation and incubation period of *Branta canadensis moffitti*, 439–445; *Burhinus oedicnemus*, 726, 727; *Empidonax minimus*, 273, 274–275; *Eurypyga helias*, 503–504; *Pelecanus erythrorhynchos*, 26–32; *Synthliboramphus antiquus*, 433–437; *Upupa epops*, 726–727; *Uria lomvia*, 433–437; *Vireo solitarius*, 477
- India, 728–729

- Ingold, Danny J., Nesting phenology and competition for nest sites among Red-headed and Red-bellied woodpeckers and European Starlings, 209-217
- Ingold, James L., Jack C. Vaughn, Sheldon I. Guttman, and Linda R. Maxson, Phylogeny of the cranes (Aves: Gruidae) as deduced from DNA-DNA hybridization and albumin micro-complement fixation analyses, 595-602
- Islands, colonization, 103-104, 255, 332-335
- Israel, 418-420, 726-727
- Italy, 411-416
- Ito, Yosiaki, Jerram L. Brown, and Jiro Kikkawa (Eds.), Animal societies: theories and facts (rev.), 165-166
- Jacana jacana*, 1-6
- Jacana, Wattled, 1-6
- James, Douglas A., and Joseph C. Neal, Arkansas birds, their distribution and abundance (rev.), 529-530
- James, Frances C., In memoriam: Robert James Newman, 464-465
- James, Ross D., see Peck, George K., and ———
- Japan, 292-300, 498-500
- Jay, Florida Scrub, 742; Steller's, 451-452
- Jehl, Joseph R., Jr., Biology of the Eared Grebe and Wilson's Phalarope in the nonbreeding season: a study of adaptations to saline lakes (rev.), 762-763; Nest-site tenacity and patterns of adult mortality in nesting California Gulls (*Larus californicus*), 102-106; and Bertram G. Murray Jr., Commentary: Response: Evolution of sexual size dimorphism, 155-157
- Johnsgard, Paul A., North American owls (rev.), 350
- Johnson, Ned K., review by, 347-349
- Jones, B., see Duke, G. E., et al.
- Jones, Elinor G., see Calder, William A., III, and ———
- Jones, Ian L., see Gaston, Anthony J., and ———
- Junco, Dark-eyed, 324-325, 451, 452
- Junco hyemalis*, 324-325, 451, 452
- Kamada, Tsutomu, see Taoka, Miki, et al.
- Kansas, 324-325
- Karasov, W. H., see Levey, Douglas J., and ———; see Martínez del Rio, Carlos, et al.
- Keith, Stuart, see Fry, C. Hilary, et al.
- Kemp, Kenneth E., see Shuman, Theresa W., et al.
- Keppleman, Tony, Hummingbirds (rev.), 530
- Kestrel, American, 451, 452; Eurasian, 539-546, 549-557
- Kikkawa, Jiro, and Derek J. Anderson (Eds), Community ecology: pattern and process (rev.), 168-170; see Ito, Yosiaki, et al.
- King, Ben, review by, 527
- King, James R., and L. Richard Mewaldt, In memoriam: Donald S. Farner, 710-713
- King, Warren B., review by, 526-527
- Kingbird, Western, 735, 736
- Kinglet, Golden-crowned, 451-452
- Kinkel, Linda K., Lasting effects of wing tags on Ring-billed Gulls, 619-624
- Klaassen, Marcel, Claus Bech, Dirkjan Masman, and Guri Slagsvold, Growth and energetics of Arctic Tern chicks (*Sterna paradisaea*), 240-248
- Klimstra, W. D., see Ellsworth, Darrell L., et al.
- Knot, Red, 94-100, 326-330
- Konstantinov, V. M., see Ilyichev, V. D., et al.
- Korea, 471-473
- Krajewski, Carey, Phylogenetic relationships among cranes (Gruiformes: Gruidae) based on DNA hybridization, 603-618
- Krebs, J. R., see Lessells, C. M., and ———
- Kress, Stephen W., see Podolsky, Richard H., and ———
- Kroodsma, Donald E., review by, 529
- Lanius excubitor*, 418-420
- Lanyon, Scott M., and Wesley E. Lanyon, The systematic position of the plantcutters, *Phytotoma*, 422-432
- Lanyon, Wesley E., see Lanyon, Scott M., and ———
- Larson, Terry A., see Verner, Jared, and ———
- Larus californicus*, 102-105
- L. delawarensis*, 619-624
- L. occidentalis*, 496-497
- Lauro, Brook, and Joanna Burger, Nest-site selection of American Oystercatchers (*Haematopus palliatus*) in salt marshes, 185-192
- Leduc, Alain, see Robert, Michel, et al.
- Lein, M. Ross, editor of book reviews, 165-174, 345-350, 525-531, 751-764; review by, 530; see Hill, Brad G., and ———
- Lemel, Jonas, Body-mass dependent fledgling order in the Great Tit, 49-492
- Leptotila rufaxilla*, 339-340
- L. verreauxi*, 339-340
- L. wellsi*, 339-340
- Lessells, C. M., and J. R. Krebs, Age and breeding performance of European Bee-eaters, 375-382
- Lever, Christopher, Naturalized birds of the world (rev.), 763-764
- Levey, D. J., and William H. Karasov, Digestive responses of temperate birds switched to fruit or insect diets, 675-686; see Martínez del Rio, Carlos, et al.
- Lewis, David B., see Briggs, Kenneth T., et al.
- Ligon, J. David, Cynthia Carey, Sandra H. Ligon, and Greg H. Farley, Commentary: Response to Ward et al., 343-344; and Peter B. Stacey, On the significance of helping behavior in birds, 700-705
- Ligon, Sandra H., see Ligon, J. David, et al.
- Limnodromus griseus*, 94-100
- L. scolopaceus*, 326-330
- Limpkin, 603-609, 612, 616-618
- Linder, Yosef, see Ward, David, et al.

- Lindstrom, Åke, Finch flock size and risk of hawk predation at a migratory stopover site, 225–232
- Lipaugus subularis*, 426, 428
- Livezey, Bradley C., review by, 763
- Lohrer, Fred E., John L. Trapp, David A. Wiedenfeld, Hugh A. Ford, and Innes C. Cuthill, editors of *Recent Ornithological Literature*, 1A–35A, 1B–69B, 1C–30C, 1D–44D
- Longevity (see Mortality and survival)
- Longspur, Lapland, 738–741
- Louisiana, 326–330
- Loxia leucoptera*, 483–485
- Lundberg, Arne, see Hoglund, Jacob, and —
- Luscinia megarhynchos*, 501–502
- Lyon, Bruce E., and Michael P. L. Fogden, Breeding biology of the Sunbittern (*Eurypyga helias*) in Costa Rica, 503–507
- Macronectes giganteus*, 303–312
- Magpie, Black-billed, 635–643
- Maine, 80–84, 332–335, 483–485, 687–697
- Manakin, Golden-headed, 34–39; Red-headed, 34–39; Round-tailed, 34–39
- Manitoba, 26–32, 42–47, 177–184, 269–276, 358–365, 483–485, 577–582
- Mannan, R. William, see Dawson, James W., and —
- Marks, Jeffrey S., John H. Doremus, and Richard J. Cannings, Polygyny in the Northern Saw-whet Owl, 732–734
- Martínez del Rio, Carlos, W. H. Karasov, and D. J. Levey, Physiological basis and ecological consequences of sugar preferences in Cedar Waxwings, 64–71
- Marvil, Rebecca E., and Alexander Cruz, Impact of Brown-headed Cowbird parasitism on the reproductive success of the Solitary Vireo, 476–480
- Masman, Dirkjan, see Klaassen, Marcel, et al.; see Meijer, Theo, et al.
- Mass and measurements of *Aegolius acadicus*, 732, 733; *Aegelaius phoeniceus*, 364; *Arenaria interpres*, 475; *Branta canadensis moffitti*, 442; *Calidris alba*, 475; *C. pusilla*, 475; *Catoptrophorus semipalmatus*, 475; *Charadrius*, 154; *Cinclocerthia gutturalis*, 250, 252; *C. ruficauda*, 250, 252; *Dendroica petechia*, 364; *D. striata*, 11–12, 13; *Dolichonyx oryzivorus*, 588; *Empidonax minimus*, 271, 273; *Eudocimus albus*, 141, 142; *Falco tinnunculus*, 553, 554, 555; *Geospiza difficilis*, 145–146; *G. fuliginosa*, 145–146; *Larus californicus*, 103; *Macronectes giganteus*, 304–305, 309; *Molothrus ater*, 360, 361, 478; *Numenius phaeopus*, 475; *Oceanodroma leucorhoa*, 689, 690, 693; *Oenanthe oenanthe*, 20, 21; *Parus major*, 491; *Passerculus sandwichensis*, 666–672; *Phiomachus pugnax*, 337; *Pica pica*, 639; *Pluvialis squatarola*, 475; *Pygoscelis papua*, 304–305, 309; *Ramphocinclus brachyurus*, 250, 252; *Scolopaci*, 154; *Sitta europaea*, 563–564; *Sterna paradisea*, 242, 243, 245, 246; *Sturnus vulgaris*, 676, 679; *Synthliboramphus antiquus*, 655, 656; *Turdus migratorius*, 676, 679; *Vireo solitarius*, 478
- Massa, Bruno, see Randi, Ettore, et al.
- Mating systems of *Aegolius acadicus*, 732–733; *Aegelaius phoeniceus*, 107–115; *Chen caerulescens caerulescens*, 177–184; *Coturnix japonica*, 283; *Egretta garzetta*, 292, 294–295, 297–300; *Jacana jacana*, 1–6; *Lanius excubitor*, 418–420; *Merops apiaster*, 375–381; *Mimus polyglottos*, 664; *Phiomachus pugnax*, 336–338; *Pica pica*, 635–643
- Matthysen, Erik, Territorial and nonterritorial settling in juvenile Eurasian Nuthatches (*Sitta europaea* L.) in summer, 560–567
- Maxson, Linda R., see Ingold, James L., et al.
- Mayfield, Harold R., In memoriam: Frank W. Preston, 714–717
- Mayr, Ernst, Commentary: A new classification of the living birds of the world, 508–512; Toward a new philosophy of biology: observations of an evolutionist (rev.), 751–752
- McGuire, Terry T., see Romagnano, Linda, et al.
- McIntyre, Roy F., see Cheng, Kimberly M., et al.
- McLaughlin, Robert L., and Robert D. Montgomerie, Early nest departure does not improve the survival of Lapland Longspur chicks, 738–741
- McNeil, Raymond, see Robert, Michel, et al.
- Means, Barbara, and Richard Means, Biographies for birdwatchers (rev.), 754–756
- Means, Richard, see Means, Barbara, and —
- Measurements (see Mass and —)
- Meijer, Theo, Dirkjan Masman, and Serge Daan, Energy-ergetics of reproduction in female kestrels, 549–559
- Melanerpes carolinus*, 209–216
- M. erythrocephalus*, 209–216
- Melospiza melodia*, 485–487
- Memorials of Oliver L. Austin Jr., 706–709; James Bond, 718–720; Donald J. Borror, 321–323; Betty Carnes, 140; Donald S. Farner, 710–713; Klaus Immelmann, 723; Robert James Newman, 464–465; Frank W. Preston, 714–717; Ralph W. Schreiber, 137–140; Frank B. Smith, 466; Nikolaas Tinbergen, 722; Yoshimaro Yamashina, 721
- Meng, Heinz, review by, 349–350
- Mengel, Robert M., review by, 167–168
- Merlin, 225, 227–230
- Merops apiaster*, 375–381
- Metabolism of *Oenanthe oenanthe*, 18–24; *Phoeniculus purpureus*, 342, 343
- Mewaldt, L. Richard, see King, James R., and —
- Mexico, 124–128, 402–409
- Michigan, 619–624, 730–731
- Micrastur ruficollis*, 407, 408
- Migration and orientation, 8–15, 143, 228, 231, 233–237, 488–489, 501–502, 623
- Migration and orientation of *Acrocephalus scirpaceus*, 501–502; *Dendroica striata*, 8–15; *Ficedula hypoleuca*, 501–502; *Fringilla coelebs*, 228–231; *F. montifringilla*,

- 228, 231; *Larus delawarensis*, 623; *Luscinia megarhynchos*, 501–502; *Muscicapa striata*, 501–502; *Phylloscopus trochilus*, 501–502; *Selasphorus rufus*, 488–489; *Sylvia borin*, 501–502; *S. communis*, 501–502; trans-Saharan passerines, 233–237
- Millam, J. R., see Yamamoto, J. T., et al.
- Mimus polyglottos*, 659–664
- Mindell, David P., and Rodney L. Honeycutt, Variability in transcribed regions of rDNA and early divergences in birds, 539–548
- Mississippi, 209–216
- Mockingbird, Northern, 659–664
- Molothrus ater*, 358–365, 452, 476–479, 577–582, 734–737
- Molts and plumages of *Picoides pubescens*, 314–319; *Pipra chloromerus*, 34–39; *P. erythrocephala*, 34–39; *P. rubrocincta*, 34–39
- Monasterio, Maximina, see Vuilleumier, François, and —
- Monroe, Burt L., Jr., Commentary: Response to E. Mayr, 515–516; review by, 529–530; Anne L. Stamm, and Brainard L. Palmer-Bell Jr., Annotated checklist of the birds of Kentucky (rev.), 530
- Montana, 488–489
- Montevecchi, William A., and Leslie M. Tuck, Newfoundland birds: exploitation, study, conservation (rev.), 166–167
- Montgeronie, Robert D., see McLaughlin, Robert L., and —
- Monticola cinclorhynchus*, 728–729
- Moreno, Juan, Variation in daily energy expenditure in nesting Northern Wheatears (*Oenanthe oenanthe*), 18–25
- Mortality and survival of *Calcarius lapponicus*, 738–741; *Dolichonyx oryzivorus*, 591–592; *Empidonax minimus*, 272–273, 274, 276; *Egretta garzetta*, 292, 296–297; *Fringilla coelebs*, 225, 228–229, 231; *F. montifringilla*, 225, 228–229, 231; *Hylocichla mustelina*, 402, 405, 408, 409; *Jacana jacana*, 1–6; *Larus californicus*, 102–105; *L. delawarensis*, 623; *Molothrus ater*, 360, 361–362; *Picoides borealis*, 517; *Ptychoramphus aleuticus*, 496–497; *Sitta europaea*, 565–566; *Sturnus dougalii*, 367–372; *Sturnus vulgaris*, 131; *Vireo solitarius*, 477, 478
- Morus bassanus*, 383–388
- M. capensis*, 383–388
- M. serrator*, 383–388
- Mueller, Helmut C., Commentary: Aerial agility and the evolution of reversed sexual dimorphism (RSD) in shorebirds, 154–155
- Murphy, Andrew J., and David A. Boag, Body reserve and food use by incubating Canada Geese, 439–446
- Murray, Bertram G., Jr., A critical review of the transoceanic migration of the Blackpoll Warbler, 8–17; see Jehl, Joseph R., Jr., and —
- Murre, Thick-billed, 433–437
- Murrelet, Ancient, 433–437, 653–658
- Muscicapa striata*, 501–502
- Myadestes townsendi*, 452
- Mycteria americana*, 539–546
- Myers, J. P., see Castro, Gonzalo, and —
- Neal, Joseph C., see James, Douglas A., and —
- Nebraska, 118–123
- Nelson, Douglas A., Gull predation on Cassin's Auklet varies with the lunar cycle, 495–497
- Nest and nest site of *Bucephala albeola*, 569–571, 573–574; *Cephus columba*, 193–201; *Empidonax minimus*, 270, 276; *Euryptila helias*, 503; *Haematopus palliatus*, 185–191; *Larus californicus*, 102, 103, 104; *Melanerpes carolinus*, 209–216; *M. erythrocephalus*, 209–216; *Oceanodroma leucorhoa*, 332–335; *Phalacrocorax penicillatus*, 391–393; *Sturnus vulgaris*, 209–216; *Upupa epops*, 726–727; *Vireo solitarius*, 477
- Nest parasitism, 128–134, 358–365, 476–479, 734–737
- Nestling of *Euryptila helias*, 504–505; *Oenanthe oenanthe*, 22; *Parus major*, 490–491; *Sterna paradisaea*, 240–247; *Sturnus vulgaris*, 133; *Vireo solitarius*, 477–478
- Netherlands, 549–557
- Nettleship, David N., review by, 525–526
- New Brunswick, 148–150, 483–485
- New Jersey, 185–191
- New York, 185–191, 584–592
- Nichols, Cathleen, R., see Cheng, Kimberly M., et al.
- Nichols, James D., see Spendelow, Jeffrey A., and —
- Nighthawk, Common, 451, 452, 539–546
- Night-Heron, Black-crowned, 94–100, 629; Yellow-crowned, 94–100
- Nightingale, European, 501–502
- Nomenclature, 38, 341, 613, 743–745, 750
- North Carolina, 185–191, 326–330, 747–749
- Northwest Territories, 433–437, 738–741
- Norway, 240–247
- Nuechterlein, Gary L., and Deborah P. Buitron, Diving differences between Western and Clark's grebes, 467–470
- Numerius phaeopus*, 94–100, 474–476
- Nuthatch, Eurasian, 560–566; Red-breasted, 452
- Nycticorax nycticorax*, 94–100, 629
- N. violaceus*, 94–100
- Nymphicus hollandicus*, 86–91
- Oceanodroma leucorhoa*, 80–84, 148–150, 332–335, 498–500, 687–697
- O. monorhis*, 471–473
- O'Connor, Raymond J., review by, 758–760
- Oenanthe oenanthe*, 18–24
- Ohio, 314–319
- Okumura, Hiroshi, see Taoka, Miki, et al.
- Ontario, 483–485
- Opornis tolmieii*, 451, 452, 456
- Oregon, 467–470
- Oreortyx pictus*, 452
- Orians, Gordon H., see Beletsky, Les D., et al.
- Oriole, Northern, 735–737

- Osteology of *Ampelion rubrocristata*, 426–431; *A. rufaxilla*, 426–431; *A. sclateri*, 426–431; *A. strelzmanni*, 426–431; Pelecaniformes, 383–388; *Phytotoma raimondii*, 426–431; *P. rara*, 426–431; *P. rutila*, 426–431; *Pipreola intermedia*, 426–429
- Ouellet, Henri, review by, 166–167
- Owen, Myrfyn, G. L. Atkinson-Willes, and D. G. Salmon, Wildfowl in Great Britain (rev.), 758–760
- Owl, Northern Saw-whet, 732–733; Spotted, 451, 452
- Oxyura jamaicensis*, 42–47
- Oystercatcher, American, 185–191
- Palmer-Bell, Brainard L., Jr., see Monroe, Burt L., Jr., et al.
- Panama, 1–6
- Papasula abbotti*, 383–388
- Parabuteo unicinctus*, 480–482
- Parasites (see Diseases and ——)
- Parasitism (see Nest parasitism)
- Parentage studies, 128–134, 177–184
- Parkes, Kenneth C., In memoriam: James Bond, 718–720
- Parus atricapillus*, 259–266, 268, 645–650, 652
- P. dichrous*, 728–729
- P. gambeli*, 259–266, 268, 451, 452, 459, 645–650, 652
- P. major*, 490–491
- P. melanolophus*, 728–729
- P. monticolus*, 728–729
- P. rubidiventris*, 728–729
- Passer domesticus*, 150–153, 539–546, 730–731
- Passerculus sandwichensis*, 666–672
- Passerella iliaca*, 451, 452, 456
- P. i. schistacea*, 157–159, 160–163
- Passerina cyanea*, 747–749
- Peck, George K., and Ross D. James, Breeding birds of Ontario: nidiology and distribution. Vol. 2, Passerines (rev.), 166
- Pelecanus erythrorhynchos*, 26–32, 383–388
- P. occidentalis*, 383–388
- P. onocrotalus*, 383–388
- P. philippensis*, 383–388
- P. rufescens*, 383–388
- Pelican, White, 26–32
- Pelecaniformes, 383–388
- Penguin, Gentoo, 303–312
- Pennsylvania, 150–153
- Pennycuick, C. J., and Toni de Santo, Flight speeds and energy requirements for White Ibises on foraging flights, 141–144
- Peru, 34–39
- Phaethon aethereus*, 383–388
- P. lepturus*, 383–388
- P. rubricauda*, 383–388
- Phalacrocorax africanus*, 383–388
- P. atriceps*, 383–388
- P. auritus*, 383–388
- P. bougainvillii*, 383–388
- P. capensis*, 383–388
- P. carbo*, 383–388
- P. coronatus*, 383–388
- P. gaimardi*, 383–388
- P. harrisi*, 383–388
- P. magellanicus*, 383–388
- P. melanoleucus*, 383–388
- P. neglectus*, 383–388
- P. olivaceus*, 383–388
- P. pelagicus*, 383–388
- P. penicillatus*, 383–388, 389–400
- P. pygmeus*, 383–388
- P. sulcirostris*, 383–388
- P. urile*, 383–388
- P. varius*, 383–388
- Phalarope, Red, 326–330; Red-necked, 326–330; Wilson's 326–330
- Phalaropus fulicaris*, 326–330
- P. lobatus*, 326–330
- P. tricolor*, 326–330
- Pheucticus melanocephalus*, 452
- Philomachus pugnax*, 336–338
- Phoenicircus nigricollis*, 426, 428
- Phoenicopterus ruber*, 94–100
- Phoeniculus purpureus*, 342, 343–344
- Phoenicurus frontalis*, 728–729
- Phylloscopus trochilus*, 501–502
- Phylogeny (see Taxonomy)
- Physiology, digestion (see Physiology, metabolism)
- Physiology, egg laying, 549, 552–557, 653–658, 730–731; energy cost of activities, 18–24, 141–143, 243–246, 439–445, 474–476, 549, 552–553, 555–557; flight, 141–143, 154, 474–476; growth, 22, 242, 243, 245–246, 314–319; hormones, 107–115, 485–487; incubation, 439–445, 653–658; metabolism, 18–24, 64–70, 80–84, 203–207, 243–244, 245, 303–312, 324–325, 342, 343, 353–356, 549, 552, 555–557, 675–684, 687–697; migration, 143; molt, 314–319; nutrient reserves, 314–319, 439–445, 549, 554–556, 557, 657; Nutritional status (see Physiology, nutrient reserves); reproduction, 279–383, 286–289; temperature regulation, 242–243, 244–246, 342, 343–344
- Physiology of *Agelaius phoeniceus*, 107–115; *Amazilia saucerottei*, 203–207; *Arenaria interpres*, 474–476; *Bombycilla cedrorum*, 64–70; *Branta canadensis moffitti*, 439–445; *Calidris alba*, 474–476; *C. pusilla*, 474–476; *Cardinalis cardinalis*, 324–325; *Catoptrophorus semipalmatus*, 474–476; *Chlorostilbon canivetii*, 203–207; *Circus cyaneus*, 353–356; *Coturnix japonica*, 279–283, 286–289; *Eudocimus albus*, 141–143; *Falco tinunculus*, 549, 552–557; *Junco hyemalis*, 324–325; *Macronectes giganteus*, 303–312; *Melospiza melodia*, 485–487; *Numenius phaeopus*, 474–476; *Oceanodroma leucorhoa*, 80–84, 687–697; *Oenanthe oenanthe*, 18–24; *Passer domesticus*, 730–731; *Phoeniculus purpureus*, 342, 343–344; *Picoides pubescens*, 314–319; *Pluvialis squatarola*, 474–476; *Pygoscelis papua*, 303–312; *Selasphorus rufus*, 724–725; *Spizella arborea*, 324–325; *Sterna paradisaea*, 240–247; *Sturnus vulgaris*, 675–684; *Synthliboramphus antiquus*, 653–658; Tur-

- dus migratorius*, 675–684; *Zonotrichia querula*, 324–325
- Phytotoma raimondii*, 422–431
- P. rara*, 422–431
- P. rutila*, 422–431
- Pica pica*, 635–643
- Picman, Jaroslav, Mechanism of increased puncture resistance of eggs of Brown-headed Cowbirds, 577–583
- Picoides albolarvatus*, 451, 452, 459
- P. borealis*, 517–518, 518–519
- P. pubescens*, 314–319
- P. villosus*, 451, 452, 459
- Pierson, Elizabeth Cary, Charles E. Huntington, and Nathaniel Wheelwright, Homing experiment with Leach's Storm-Petrels, 148–150
- Pigeon, Band-tailed, 452
- Pigmentation of *Aphelocoma ultramarina*, 124–128; *Cinclorhynchus gutturalis*, 251, 253, 255; *C. ruficauda*, 251, 254, 255; *Egretta garzetta*, 292, 293, 294, 298; *Lepotila rufaxilla*, 340; *L. verreauxi*, 340; *L. wellsi*, 340; *Philomachus pugnax*, 336–338; *Pipra chloromeros*, 34–39; *P. erythrocephala*, 34–39
- Pinicola enucleator*, 452
- Pinshow, Berry, see Ward, David, et al.; see Yosef, Reuven, and ———
- Pipilo chlorurus*, 451, 452, 456
- P. erythrophthalmus*, 451, 452, 456
- Pipra chloromeros*, 34–39
- P. erythrocephala*, 34–39
- P. rubrocapilla*, 34–39
- Pipreola intermedia*, 426, 429
- Piranga ludoviciana*, 451–452
- Place, Allen R., Nina C. Stoyan, Robert E. Ricklefs, and Ronald G. Butler, Physiological basis of stomach oil formation in Leach's Storm-Petrel (*Oceanodroma leucorhoa*), 687–699; see Duke, G. E., et al.; see Roby, Daniel D., et al.
- Plover, Black-bellied, 94–100; Semipalmated, 94–100; Wilson's, 94–100
- Plumages (see Molts and ———)
- Pluvialis squatarola*, 94–100, 474–476
- Podolsky, Richard H., and Stephen W. Kress, Factors affecting colony formation in Leach's Storm-Petrel, 332–336
- Poephila guttata*, 750
- Poland, 730–731
- Polyandry (see Mating systems)
- Polygyny (see Mating systems)
- Pomatorhinus erythrogenys*, 728–729
- Pomatostomidae, 341
- Pomatostominae, 341
- Portugal, 411–416
- Potter, Eloise F., Commentary: Response to P. R. Ehrlich, D. S. Dobkin, and D. Wheye, 163–164
- Powell, David W., see Gaston, Anthony J., and ———
- Power, Harry W., see Romagnano, Linda, et al.
- Predation by *Accipiter gentilis*, 225–231; *A. nisus*, 225–231; *Buteo buteo*, 225–231; *Circus cyaneus*, 225–231; *Corvus caurinus*, 197–198; *Falco columbarius*, 225–231; *Glaucidium brasilianum*, 406, 407, 408; *Larus occidentalis*, 496–497; *Micrastur ruficollis*, 407, 408
- Predation on *Aphelocoma ultramarina*, 126–127; *Calciarius lapponicus*, 739; *Casmerodius albus*, 628; *Cephus columba*, 193, 196–198, 200–201; *Egretta caerulea*, 628; *Empidonax minimus*, 269, 270, 275; *Eudocimus albus*, 628; *Fringilla coelebs*, 225–231; *F. montifringilla*, 225–231; *Haematopus palliatus*, 188–189; *Hylocichla mustelina*, 406, 407, 408, 409; *Larus californicus*, 102, 103, 104; *Molothrus ater*, 361–362; *Parus major*, 490, 491; *Pica pica*, 639; *Ptychoramphus aleuticus*, 495–497
- Ptiliogonyx cinereus*, 743–745
- Ptychoramphus aleuticus*, 495–497
- Pulich, Warren M., The birds of north central Texas (rev.), 530–531
- Pygmy-Owl, Ferruginous, 406, 407, 408
- Pygoscelis papua*, 303–312
- Pyrrhula erythrocephala*, 728–729
- Quail, Japanese, 279–283, 286–289; Mountain, 452
- Quebec, 483–485
- Quinn, Thomas W., J. Chris Davies, Fred Cooke, and Bradley N. White, Genetic analysis of offspring of a female-female pair in the Lesser Snow Goose (*Chen c. caerulescens*), 177–184
- Ramos, Mario A., see Rappole, John H., et al.
- Ramphocinclus brachyurus*, 249–256, 258
- Rand, Michael R. W., see Hudson, Peter J., and ———
- Randi, Ettore, Fernando Spina, and Bruno Massa, Genetic variability in Cory's Shearwater (*Calonectris diomedea*), 411–417
- Rappole, John H., Mario A. Ramos, and Kevin Winkler, Wintering Wood Thrush movements and mortality in southern Veracruz, 402–410
- Rassi, P., and R. Vaisanen (Eds.), Threatened animals and plants in Finland (rev.), 173
- Raven, Common, 452
- Reed, J. Michael, see Hagen, John M., and ———
- Reed-Warbler, European, 501–502
- Regulus satrapa*, 451–452
- Remsen, J. V., review by, 527–529
- Reynard, George B., and Orlando H. Garrido, Bird songs in Cuba/Cantos de Aves en Cuba (rev.), 529
- Rickard, William H., see Eberhardt, Lester E., et al.
- Ricklefs, Robert E., see Place, Allen R., et al.
- Ripley, S. Dillon, In memoriam: Yoshimoro Yamashina, 721
- Rising, James D., and Keith M. Somers, The measurement of overall body size in birds, 666–675
- Rising, R. M., Commentary: Response to R. M. Zink, 160–163
- Robel, Robert J., see Shuman, Theresa W., et al.
- Robert, Michel, Raymond McNeil, and Alain Leduc,

- Conditions and significance of night feeding in shorebirds and other water birds in a tropical lagoon, 94-101
- Robertson, Peter, see Hill, David, and —
- Robin, American, 452, 675-684, 735, 736
- Roby, Daniel D., Karen L. Brink, and Allen R. Place, Relative passage rates of lipid and aqueous digesta in the formation of stomach oils, 303-313
- Rohwer, Sievert, Carol D. Spaw, and Eivin Røsktaf, Costs to Northern Orioles of puncture-ejecting parasitic cowbird eggs from their nests, 734-738
- Romagnano, Linda, Terry R. McGuire, and Harry W. Power, Pitfalls and improved techniques in avian parentage studies, 129-136
- Roseberry, John L., see Ellsworth, Darrell L., et al.
- Røsktaf, Eivin, see Rohwer, Sievert, et al.
- Rotenberry, John T., review by, 168-170
- Roudybush, T. E., see Yamamoto, J. T., et al.
- Round, Philip D., Resident forest birds in Thailand: their status and conservation (rev.), 761
- Ruff, 336-338
- Rusch, Donald H., see Small, Robert J., and —
- Salmon, D. G., see Owen, Myrfyn, et al.
- Sanderling, 326-330
- Sandpiper, Least, 94-100; Semipalmated, 94-100; Stilt, 94-100, 326-330; Western, 94-100
- Sapsucker, Red-breasted, 452; Williamson's, 452
- Sato, Tetsu, see Taoka, Miki, et al.
- Saxicola ferrea*, 728-729
- Scolopaci, 154-155, 155-156
- Sealy, Spencer G., see Briskie, James V., and —
- Seicercus xanthoschistos*, 728-729
- Selasphorus rufus*, 488-489, 724-725
- Sellers, Timothy J. (Ed.), Bird respiration (rev.), 345
- Sex, determination of, 250, 498, 522, 523-524, 654
- Sex ratio of *Aquila chrysaetos*, 520-521, 521-522, 523-524; *Molothrus ater*, 360, 365; *Passer domesticus*, 151-152; *Picoides borealis*, 517
- Sexual size dimorphism, 154-155, 155-156, 249, 250-255, 360, 365
- Shearwater, Cory's, 411-416, 417
- Sherrod, Steve K., review by, 347
- Shields, K. M., see Yamamoto, Y. T., et al.
- Shrike, Northern, 418-420
- Shuman, Theresa W., Robert J. Robel, John L. Zimmerman, and Kenneth E. Kemp, Variance in digestive efficiencies of four sympatric avian granivores, 324-326
- Short, L. L., review by, 173-174
- Sialia currucoidea*, 451, 452, 459
- S. mexicana*, 452
- Sibley, Charles G., Commentary: Response to E. Mayr, 512-515
- Siegel-Causey, Douglas, review by, 174
- Sitta canadensis*, 452
- S. europaea*, 560-566
- S. leucopsis*, 728-729
- Slagsvold, Guri, see Klaassen, Marcel, et al.
- Small, Robert J., and Donald H. Rusch, The natal dispersal of Ruffed Grouse, 72-79
- Solitaire, Townsend's, 452
- Somers, Keith M., see Rising, James D., and —
- Song (see Vocalizations)
- Sonograms of *Dendroica graciae*, 50, 51, 54; *Leptotila rufaxilla*, 339; *L. verreauxi*, 339; *L. wellsi*, 339; *Oceanodroma leucorhoa*, 499; *O. monorhis*, 472
- South Carolina, 141-143
- Spain, 501-502
- Spaw, Carol D., see Rohwer, Sievert, et al.
- Sparrow, American Tree, 118-123, 324-325; Chipping, 452; Fox, 157-159, 160-163, 451, 452, 456; Harris', 324-325; House, 150-153, 539-546, 598, 730-731; Savannah, 666-672; Song, 485-487
- Sparrowhawk, European, 225-231
- Spendlow, Jeffrey A., and James D. Nichols, Annual survival rates of breeding adult Roseate Terns, 367-374
- Sphyrapicus ruber*, 452
- S. thyroideus*, 452
- Spina, Fernando, see Randi, Ettore, et al.
- Spizella arborea*, 118-123, 324-325
- S. passerina*, 452
- Stacey, Peter B., see Ligon, J. David, and —
- Staicer, Cynthia A., Characteristics, use, and significance of two singing behaviors in Grace's Warbler (*Dendroica graciae*), 49-63
- Stamm, Anne L., see Monroe, Burt L., Jr., et al.
- Starling, European, 129-134, 209-216, 675-684
- Statistical and analytical methods, 666-672
- Stellula calliope*, 452
- Sterna dougallii*, 367-372
- S. paradisaea*, 240-247
- Stilt, Black-necked, 94-100
- Stokes, Allen W., review by, 757-758
- Stokes, Donald W., and Lillian Q. Stokes, A guide to bird behavior, vol. 3 (rev.), 761-762
- Stokes, Lillian Q., see Stokes, Donald W., and —
- Storer, Robert W., Geographic variation and sexual dimorphism in the tremblers (*Cinclocerthia*) and White-breasted Thrasher (*Ramphocinclus*), 249-258
- Stork, Wood, 539-546
- Storm-Petrel, Leach's, 80-84, 148-150, 332-335, 498-500, 687-697; Swinhoe's, 471-473
- Stoyan, Nina C., see Place, Allen R., et al.
- Strix occidentalis*, 451, 452
- Sturnus vulgaris*, 129-134, 209-216, 675-684
- Sula dactylatra*, 383-388
- S. leucogaster*, 383-388
- S. nebulosus*, 383-388
- S. sula*, 383-388
- S. variegata*, 383-388
- Sunbittern, 503-507
- Swan, Trumpeter, 539-546
- Sweden, 18-24, 336-338, 490-491
- Sylvia borin*, 501-502
- S. communis*, 501-502

- Synthliboramphus antiquus*, 433–437, 653–658
 Systematics (see Taxonomy)
- Taeniopygia guttata*, 750
 Taigen, Theodore L., review by, 345
 Tanager, Western, 451–452
 Taoka, Miki, Tetsu Sato, Tsutomu Kamada, and Hiroshi Okumura, Sexual dimorphism of chatter-calls and vocal sex recognition in Leach's Storm-Petrels (*Oceanodroma leucorhoa*), 498–501; Pyong-Oh Won, and Hiroshi Okumura, Vocal behavior of Swinhoe's Storm-Petrel (*Oceanodroma monorhis*), 471–474
 Taxonomy, 508–512, 512–515, 515–516, 539–546, 595–601
 Taxonomy of *Ampelion rubrocristata*, 422, 426, 429–431; *A. rufaxilla*, 422, 426, 429–431; *A. sclateri*, 422, 426, 429–431; *A. stresemanni*, 422, 426, 429–431; *Bubo bubo*, 539–546; *Buteo buteo*, 539–546; *Calidris alba*, 330; *C. canutus*, 330; *C. himantopus*, 330; *Cathartes melambrotus*, 539–546; *Chordeiles minor*, 539–546; *Cinclocerthia gutturalis*, 249, 255–256; *C. ruficauda*, 249, 255–256; *Crypturellus undulatus*, 539–546; *Cygnus buccinator*, 539–546; *Falco tinnunculus*, 539–546; *Gallus gallus*, 539–546; *Geospiza difficilis*, 144–147; Gruidae, 595–601, 603–613; *Leptotila rufaxilla*, 339–340; *L. verreauxi*, 339–340; *L. wellsi*, 339–340; *Limnodromus scolopaceus*, 330; *Mycteria americana*, 539–546; *Passer domesticus*, 539–549; Pelecaniformes, 384–387; *Phalaropus fulicaris*, 326, 329–330; *P. lobatus*, 326, 329–330; *P. tricolor*, 326, 329–330; *Phytotoma raimondii*, 422–426, 429–431; *P. rara*, 422, 426, 429–431; *P. rutila*, 422, 426, 429–431; *Pipra erythrocephala*, 39; *Ramphocinclus brachyurus*, 249, 256; *Tringa melanoleuca*, 330
- Temeles, Ethan J., Effect of prey consumption on foraging activity of Northern Harriers, 353–357
 Tern, Arctic, 240–247; Roseate, 367–372
 Territory of *Jacana jacana*, 3; *Parus atricapillus*, 259–266; *P. gambeli*, 259–266
 Thrasher, White-breasted, 249–256, 258
 Thrush, Hermit, 452; Swainson's, 451, 452; Wood, 402–409
 Tiebout, Harry M. III, Tests of a model of food passage rates in hummingbirds, 203–208
 Tinamou, Undulated, 539–546
 Tit, Great, 490–491
 Tome, Michael W., Search-path characteristics of foraging Ruddy Ducks, 42–48
 Towhee, Green-tailed, 451, 452, 456; Rufous-sided, 451, 452, 456
 Trapp, John L., see Lohrer, Fred E., et al.
 Trembler, Brown, 249–256, 258; Gray, 249–256, 258
Tringa flavipes, 94–100
T. melanoleuca, 94–100, 326–330
Troglodytes troglodytes, 451, 452
 Tuck, Leslie M., see Monteverechi, William A., and —
Turdus boulboul, 728–729
- T. migratorius*, 452, 675–684, 735, 736
T. viscivorus, 728–729
 Tyler, W. Breck, see Briggs, Kenneth T., et al.
Tyrannus verticalis, 735, 736
- Udvardy, M. D. F., and M. E. Udvardy, review by, 173
 Udvardy, M. E., see Udvardy, M. D. F., and —
Upupa epops, 726–727
 Urban, Emil K., see Fry, C. Hilary, et al.
Uria lomvia, 433–437
- Vagvolgyi, Joseph, and Maria W. Vagvolgyi, The taxonomic status of the Small Ground-Finch, *Geospiza* (Aves: Emberizidae) of Genovesa Island, Galápagos, and its relevance to interspecific competition, 144–148
 Vagvolgyi, Maria W., see Vagvolgyi, Joseph, and —
 van Marle, J. G., and Karel H. Voous, The birds of Sumatra: annotated check-list (rev.), 527
 Vaughn, Jack C., see Ingold, James L., et al.
 Venezuela, 94–100
 Verbeek, N. A. M., see Emms, S. K., and —
Vermivora ruficapilla, 452
 Vermont, 483–485
 Verner, Jared, and Terry A. Larson, Richness of breeding bird species in mixed-conifer forests of the Sierra Nevada, California, 447–463
Vireo gilvus, 451–452
V. solitarius, 452, 476–479
 Vireo, Solitary, 452, 476–479; Warbling, 451–452
 Vocalizations, 49–62, 124–128, 255–256, 332–335, 339–340, 471–473, 498–500, 659, 662, 663–664; geographic variations in, 124–128; variety of, 49–62, 471–473, 498–500
 Vocalizations of *Aphelocoma ultramarina*, 124–128; *Cinclocerthia gutturalis*, 255–256; *C. ruficauda*, 255–256; *Dendroica graciae*, 49–62; *Fringilla coelebs*, 227; *F. montifringilla*, 227; *Leptotila rufaxilla*, 339–340; *L. verreauxi*, 339–340; *L. wellsi*, 339–340; *Mimus polyglottos*, 659, 662, 663–664; *Oceanodroma leucorhoa*, 332–335, 498–500; *O. monorhis*, 471–473
 von Michaëlis, H., Birds of prey: a kinship (rev.), 174
 Voous, Karel H., see van Marle, J. G., and —
 Vuilleumier, François, and Maximina Monasterio (Eds.), High altitude tropical biogeography (rev.), 527–529; review by, 346
 Vulture, Greater Yellow-headed, 539–546
- Warbler, Blackpoll, 8–15, 17; European Willow, 501–502; Garden, 501–502; Grace's, 42–62; Hermit, 451–452; MacGillivray's, 451, 452, 456; Nashville, 452; Wilson's, 452; Yellow, 358–365, 451, 452, 456; Yellow-rumped, 451–452
 Ward, David, Berry Pinshow, Daniel Afik, Yosef Linder, and Nurit Winkler, Commentary: Cavity roosting, philopatry, and cooperative breeding in the Green Woodhoopoe may not reflect a physiological trait, 342; see Afik, Daniel, and —

- Warheit, Kenneth I., David A. Good, and Kevin de Queiroz, Variation in numbers of scleral ossicles and their phylogenetic transformations within the Pelecaniformes, 383–388
- Washington, 107–115, 218–224, 734–737
- Waxwing, Cedar, 64–70
- Weather, influence on (see Environment, influence on)
- Weatherhead, Patrick J., Sex ratios, host-specific reproductive success, and impact of Brown-headed Cowbirds, 358–366
- Welsh, Daniel A., review by, 166
- West Indies, 249–256
- Westneat, David F., Intensity of nest defense in Indigo Buntings increases with stage and not number of visits, 747–749
- Wheatear, Northern, 18–24
- Wheelwright, Nathaniel T., see Pierson, Elizabeth Cary, et al.
- Whimbrel, 94–100
- White, Bradley N., see Quinn, Thomas W., et al.
- Whitethroat, Greater, 501–502
- Wiedenfeld, David A., see Lohrer, Fred E., et al.
- Willet, 94–100
- Wilsonia pusilla*, 452
- Wingfield, John C., see Beletsky, Les D., et al.
- Winker, Kevin, see Rappole, John H., et al.
- Winkler, Nurit, see Ward, David, et al.
- Wisconsin, 64–70, 72–77, 79, 675–684
- Won, Pyong-Oh, see Taoka, Miki, et al.
- Woodhoopoe, Green, 342, 343–344
- Woodpecker, Downy, 314–319; Hairy, 451, 452, 459; Pileated, 452; Red-bellied, 209–216; Red-cockaded, 517–518, 518–519; Red-headed, 209–216; White-headed, 451, 452, 459
- Woolfenden, Glen E., In memoriam: Ralph W. Schreiber, 137–139
- Wren, Winter, 451, 452
- Xanthocephalus xanthocephalus*, 577–582
- Xiphoeua punicea*, 426, 428
- Yamamoto, J. T., K. M. Shields, J. R. Millam, T. E. Roudybush, and C. R. Grau, Reproductive activity of force-paired Cockatiels (*Nymphicus hollandicus*), 86–93
- Yellowlegs, Greater, 94–100, 326–330; Lesser, 94, 100
- Yellowthroat, Common, 599
- Yosef, Reuven, and Berry Pinshow, Cache size in shrikes influences female mate choice and reproductive success, 418–421
- Zaias, Julia, see Breitwisch, Randall, et al.
- Zimmerman, Dale A., review by, 752–754
- Zimmerman, John L., see Shuman, Theresa W., et al.
- Zink, Robert M., Commentary: The study of geographic variation, 157–160; see Dittmann, Donna L., et al.
- Zonotrichia querula*, 324–325