

An Unusual Nest Addition.—A nest of Black-chinned Hummingbird (*Archilochus alexandri*) containing two eggs, apparently far advanced in incubation, was found on May 31, 1915, near Pasadena, and at that time there was nothing to distinguish it from any typical alexandri nest. It was built upon a slender branch of a sycamore and was more or less sheltered by an overhanging leaf. On June 11 the nest was again observed, in the meantime young birds having appeared, and it was noted that nest material had been added to the upper and outward part of the walls, extending upward about a half-inch. I had occasion to again visit the nest on June 18, and the tip of the overhanging leaf had been securely caught by spider webbing to the extended wall, thus forming a somewhat stationary canopy. As a protection from the sun the nest was but little better sheltered than before. Possibly the leaf drooped down to interfere with the feed-




Fig. 1. NEST OF BLACK-CHINNED HUMMINGBIRD, SHOWING ATTACHMENT OF A LEAF TO THE RIM OF THE NEST

ing of the young or to the displeasure of the parent birds for some other reason, and this means was sought for removing the trouble.—HAROLD M. HOLLAND, *Los Angeles, California.*

A Record of the Golden Plover in the State of Washington.—This interesting species (*Charadrius dominicus dominicus*) has never before, to my knowledge, been recorded as occurring in the state of Washington. Therefore it is with great pleasure that I report the capture of an adult female at Dungeness, Clallam County, Washington. This specimen was collected by Mr. F. P. McIntyre, of Tacoma, Wash., on November 14, 1915. He very kindly presented it to me, and the skin is now in my collection. Mr. McIntyre tells me that he saw about a dozen other plover that resembled this one, but that he shot no

more. It is possible that the other birds seen might also have been *C. dominicus*; but it must be stated that the Black-bellied Plover is a common migrant in Washington, and it seems equally possible that the others might have been *Squatarola squatarola*.

Since the above-mentioned specimen was obtained Mr. D. E. Brown, of Seattle, Washington, tells me that a Golden Plover was taken some time ago in the vicinity of that city, but I believe that it has never been put on record. Mr. Brown also tells me that a year or two ago he saw on the Tacoma Flats what he feels certain was a bird of this species, owing to the large amount of yellow on the upper parts. This was in the late spring and the bird appeared to be in full breeding plumage. It seems possible, therefore, that this plover is of more frequent occurrence in the state than has been supposed.—J. H. BOWLES, *Tacoma, Washington*.

A Note on the Food of the Northern Pileated Woodpecker.—The stomach contents of a specimen of the Northern Pileated Woodpecker (*Phloeotomus pileatus abieticola*) taken at Lakeport, Lake County, California, November 5, 1915, and sent to the University of California Museum of Vertebrate Zoology, was found to be of peculiar interest. The stomach contained more than fifty carpenter ants (*Camponotus herculeaneus* subsp.) and 131 seeds of poison oak (*Rhus diversiloba*). As the seeds of poison oak are hard and without a noticeable covering of softer material it is difficult to understand what there is about them that is attractive to birds. Certain it is that the seeds are incapable of complete digestion by woodpeckers. In California poison oak is a favorite food of the Red-shafted Flicker also. Ants are commonly taken by the Flicker, California Woodpecker and by Sapsuckers.

The stomachs of two Pileated Woodpeckers taken in or near Yosemite National Park (orig. nos. 1545 and 7814, in Mus. Vert. Zool.) were filled with carpenter ants (*Camponotus herculeaneus modoc* Wheeler), many of them winged. Each stomach contained more than a hundred of these ants. In addition one stomach contained a whole fruit of manzanita (*Arctostaphylos nevadensis* Gray) and the other, four large beetle larvae (*Cerambycidae*), unidentifiable as to genus or species, which had evidently been dug out of some dead tree, as the stomach contained slivers of dead wood.

The above evidence shows that the animal food of the Pileated Woodpecker in California is largely made up of carpenter ants (*Camponotus* sp.) and to a lesser extent of wood-boring larvae. Vegetable food in the shape of poison oak seeds and the fruit of manzanita is occasionally taken.—H. C. BRYANT, *Berkeley, California*.

Occurrence of Emperor Goose in Northern California.—On November 1, 1915, Mr. S. M. Gridley, of Gridley, California, brought me an Emperor Goose for mounting. The bird was in full adult plumage, and in good physical condition. Sex identification was impossible on account of the damaged state of the internal organs. I at once appreciated the rarity of the goose and endeavored to obtain possession of the specimen, but the owner declined to part with it.

The gentleman reported that for a week prior to being killed, this goose had frequented, in the company of a flock of Cackling Geese, the territory adjacent to the tules west of the town. The bird was so tame as to lead to the supposition that it was a cripple, and only upon being closely approached did it fly.

I was further told that a similar goose had been shot last year in the same locality.—CARL S. MULLER, *Marysville, California*.

Townsend Solitaire in the San Jacinto Mountains.—In the field work conducted in the San Jacinto Mountains in 1908 by the Museum of Vertebrate Zoology, and subsequently reported upon by Grinnell and Swarth (Univ. Calif. Publ. Zool., vol. 10, 1913, pp. 197-406), the Townsend Solitaire (*Myadestes townsendi*) was not met with at any point. From the number of collectors in our party, enabling us to cover a wide extent of ground, and from the length of time we spent in the mountains, we felt justified in the conclusion that our failure to encounter this species meant that it did not breed in this southern range. It was, therefore, with the greatest interest that I recently learned from Mr. O. W. Howard that this was a mistaken idea.

Messrs. O. W. Howard and H. J. Leland spent a few days of this year (1915) in Strawberry Valley and vicinity, where they discovered two nests of the Solitaire. The first was far up a narrow canyon heading on Tahquitz Peak and opening out in the