

the latter author says, page 38, "The male does not assist in incubation, but supplies its mate with food while so engaged, and she rarely leaves the nest after the first egg has been laid; at any rate I have invariably found the bird at home if there were any eggs in the nest". Major Bendire is referred to in the sentence just previous to the one first quoted above.

For more light on this subject I have recently looked up far too many authorities to mention in this brief article, but the only other reference to the matter of the incubating habits of this species that I have so far found is in Nuttall's "Manual of the Ornithology of the United States and Canada", 2nd edition, page 720, where it says "incubation, in which both parents engage, continues for 16 days". So here are two well known ornithologists responsible for absolutely contradictory statements! The matter is "side-stepped" by every other authority that I have consulted.

Now it happened on June 24, 1921, that Dr. G. Dallas Hanna, of the California Academy of Sciences, my brother, John W. Mailliard, and I were lunching at noon time in the bed of Nicasio Creek, Marin County, California, at the base of Black Mountain, and while so occupied noticed a Belted Kingfisher fly into a hole in the opposite bank. Soon after finishing lunch we proceeded to investigate this matter and discovered a nest containing five eggs, about one-third incubated, with the male bird in the tunnel and apparently on the nest. This tunnel was about ten feet long and only twelve or fourteen inches below the surface of the ground, which was rather sandy and friable, and the cavity was easily pried open by means of an old fence picket.

The bird did not attempt to leave until the nest was almost reached in the upheaving process. As it flew off it was secured for evidence and proved to be the male, with the abdominal region partially bare, as if from sitting on the eggs.

Earlier in the day two kingfishers had been observed flying up and down the creek, and shortly before lunch a female had been taken as it perched for a moment on a snag about seventy-five yards below the nest, which we had not at that time discovered. This female was apparently the other owner, yet showed practically no sign of having been incubating, as the plumage upon the abdomen was in a good state of preservation.

While this matter is not one of great importance it is one of some interest and, as part of the life history of a well known bird, might as well be cleared up if possible, hence this short paper is written in the hope that some other observer, who may have had better opportunities to study the question, may come forward with sufficient evidence to prove the point one way or the other.—JOSEPH MAILLIARD, *San Francisco, California, August 10, 1921.*

Eastern Kingbird at Mono Lake.—On July 19, 1921, I saw an Eastern Kingbird (*Tyrannus tyrannus*) near Mono Lake, Mono County, California. In Grinnell's Distributional List (1915), there are only two records for the bird from California.—RALPH HOFFMANN, *Carpenteria, California, September 25, 1921.*

Chronicle of Additions and Eliminations Pertaining to the California State List of Birds.—The present note carries the chronicle of the birds of California forward from Pacific Coast Avifauna no. 11 (1915) and from my supplementary note in THE CONDOR of January, 1919 (vol. XXI, pp. 41-42) to October 15, 1921. I have followed the rule of letting all definite proposals "ride", as if the findings set forth were final in every respect, unless and until someone has brought forward good reasons for doubting the conclusions involved. No attention is here paid to mere changes in names; only the addition or subtraction of "concepts" of species or subspecies is considered.

ADDITIONS

1. *Larus occidentalis livens* Dwight. Dark-mantled Western Gull. (See Dwight, Proc. Biol. Soc. Wash., vol. 32, February 14, 1919, p. 11.)
2. *Phaethon aethereus* Linnaeus. Red-billed Tropic Bird. (See Law, Condor, XXI, March, 1919, p. 88.)
3. *Chen caerulescens* (Linnaeus). Blue Goose. (See Grinnell, Condor, XXII, March, 1920, p. 76.)
4. *Polyborus cheriway* (Jacquin). Audubon Caracara. (See Heath, Condor, XXI, March, 1919, p. 125.)
5. *Otus asio macfarlanei* (Brewster). MacFarlane Screech Owl. (See Grinnell, Condor, XXI, July, 1919, p. 173.)