

## SKYLIGHT POLARIZATION DOES NOT AFFECT THE MIGRATORY ORIENTATION OF EUROPEAN ROBINS<sup>1</sup>

ROLAND SANDBERG

Department of Ecology, Animal Ecology, University of Lund, Ecology Building,  
S-223 62 Lund, Sweden

*Key words:* Skylight polarization; migratory orientation; robin.

Among birds, only the Rock Dove (*Columba livia*) has been shown so far to be able to perceive the E-vector direction of linearly polarized light (Kreithen and Keeton 1974, Delius et al. 1976). The adaptive significance of this ability remains largely unknown, but may be related to the birds' capacity for orientation (Brines 1980, Brines and Gould 1982). Recently, Able (1982) reported that the orientation of nocturnally migrating White-throated Sparrows (*Zonotrichia albicollis*), is affected by manipulations of skylight polarization patterns at dusk, and Moore (1986) demonstrated that polarized light influences the early morning (dawn) orientation of night migrants tested during spring on the southwest coast of Louisiana in a similar way. In addition, recent data on the relationship between sunset orientation and polarized light, includes the Yellow-rumped Warbler (*Dendroica coronata*) as a species which responds to polarized light (F. Moore, pers. comm.). Orientation experiments have indicated that there may be significant differences between bird species in their use of sun-related cues, the stars, and the geomagnetic field for the establishment and maintenance of nocturnal orientation (Wiltshcko and Wiltshcko 1978, Able and Cherry 1985, Moore 1987). In contrast to North American Savannah Sparrows (*Passerculus sandwichensis*) and White-throated Sparrows, the European Robin (*Erithacus rubecula*) seems to rely more extensively on the geomagnetic field for their orientation during the night (Wiltshcko et al. 1971, Wiltshcko and Wiltshcko 1975, but cf. Katz and Michelson 1978 and Wallraff and Gelderloos 1978 for some conflicting evidence), as well as during the twilight period after sunset (Sandberg et al., in press a). Katz (1985), however, found that robins displayed uniform orientation during twilight tests under total overcast as well as after sunset under starry skies whereas in twilight tests under clear skies, the robins were able to select an appropriate fall migratory direction. I have undertaken experiments with migrating robins captured at the Falsterbo Bird Observatory (FBO, 55°23'N, 12°50'E) in south Sweden in order to investigate the possible influence of skylight polarization patterns on the dusk orientation of this species.

### METHODS

The tests were performed during the spring (April and May) and autumn (September and October) migration periods in 1986. Experimental birds captured at FBO were transported by car approximately 55 km to the northeast, to a test site at the ecological field station Stensoffa (55°42'N, 13°25'E), close to Lund. The robins were housed indoors in individual cages under the natural photoperiod and the local geomagnetic field. During the tests, the birds were placed in modified "Emlen-funnels" with automatic counter registration of the activity in eight sectors (see Sandberg et al., in press b for details of experimental setup and equipment) and tested for 1 hr beginning 10 min after local sunset. The birds were never allowed to see the disc of the setting sun during tests, and the sunset sky below approximately 45° elevation was screened off from view. Orientation cages were covered by a linear dichroic polarizer, having a total light transmittance of 38% (Plexiglas laminated HN-38; Polaroid). Only one test per individual and condition is included in the analysis. The four different experimental conditions were:

(1) Clear skies (<5% cloud cover); (a) polarizer E-vector oriented in the seasonally normal direction at the associated test night ("E-vector = C") and (b) polarizer E-vector oriented 90° in relation to the seasonally normal direction i.e., parallel to the sunset azimuth ("E-vector = C + 90°").

(2) Simulated total overcast. A panel of 3-mm opaque diffusing Plexiglas was placed on top of the screening shield surrounding the orientation cage i.e., above the polarizer, which thereby introduced an E-vector axis in an otherwise visually cueless situation; (a) E-vector = C and (b) E-vector = C + 90°.

Vector calculation based on counter registrations yielded a mean heading for each individual and test night. Bird-hours with less than 40 registrations were excluded from analysis, as were tests which showed a highly scattered and unreliable orientation with  $P > 0.05$  according to the Rayleigh test (Batschelet 1981). This selection led to acceptance of 83% of the total number of control tests (without polarizer, cf. below) that were conducted, and of 81% of the total number of experimental tests. There was no significant difference in the proportion, of accepted tests (1), of tests excluded because of inactive birds (2), or excluded because of scattered and unreliable orientation (3), between control and experimental categories ( $\chi^2 = 0.3$ ,  $df = 2$ ,  $P > 0.5$ ). Second order mean vectors and axes of orientation were calculated on the basis of individual mean headings for each condition. A mean axis of

<sup>1</sup> Received 20 May 1987. Final acceptance 8 August 1987.




FIGURE 1. The orientation of European Robins tested under four different experimental conditions during spring and autumn migration, respectively. Polarizer E-vector directions were either in the seasonally normal direction (E-vector = C), i.e., situated 90° from the sunset azimuth, or shifted 90° from normal (E-vector = C + 90°) resulting in a polarization axis directed towards the setting sun. Experiments with the two E-vector conditions were performed under clear skies as well as under simulated total overcast. The mean direction towards the setting sun 40 min after sunset (i.e., in the middle of the test hour) is indicated for the experiments under clear skies. Open symbols denote significant headings of birds in their first year of life, and filled symbols the corresponding headings of migratory-experienced robins. Second order mean vectors (or axes) with direction (α) and length (r) are shown in the circular diagrams, and vector lengths are drawn relative to the radius of the circle = 1. Arrows outside the circles indicate the alignment of E-vectors. The preferred mean orientation of controls under natural skies is represented by broken (significant directionality according to the Rayleigh test), or dotted (mean orientation not significant) lines.

orientation is considered when it provides a better fit to the data than the unimodal mean vector (see Batschelet 1981).

If robins are able to perceive polarized light and use it for their orientation, they may be expected either to orient directly in relation to the E-vector axis, or to use the polarization patterns indirectly to localize and orient according to the position of the setting sun. This could be of particular relevance when the sun is hidden behind clouds, or situated below the horizon as is the case during twilight. In either case, the expected directions of the robins will not differ from controls under E-vector = C, but will deviate by approximately ±90° from the control direction under E-vector = C + 90°.

Robins tested at Stensoffa during the spring and autumn seasons in 1983 to 1986 under clear and simulated overcast skies, without polarizer manipulations, were used as controls. There were no statistical differences between the total sample of controls tested at this site and the subsample of controls tested during 1986 (simultaneously with polarization tests) in either

directions or scatter (according to Mardia's [1972] "one-way classification test" and "the test for the homogeneity of the concentration parameters," respectively).

### RESULTS AND DISCUSSION

The results are shown in Figure 1. From them, the following conclusions can be drawn.

Under clear skies, the robins showed almost exactly the same mean orientation whether the E-vector was in the control (C) or shifted (C + 90°) direction. The mean angles of orientation during spring and autumn are not significantly different from the corresponding directional choices of unmanipulated controls at this site (cf. Sandberg et al., in press b). These results provide clear indications that the robins do not use E-vector directions as orientational cues. This is in sharp contrast to the results presented by Able (1982) and Moore (1986), who found profound effects of imposed patterns of skylight polarization on the orientation of their test birds. White-throated Sparrows (Able 1982) as well as Northern Waterthrushes, *Seiurus nov-*

*eboracensis*, and Kentucky Warblers, *Oporornis formosus* (Moore 1986), all responded by orienting their activity parallel to the introduced E-vectors. It should be noted that the experimental design differed between my study and those of Able (1982) and Moore (1986) in that the vector alignment (C and C + 90°) in the present case did not exclude light transmittance and color variations between experimental categories (cf. also below). This difference is partly reduced by the fact that a greater proportion of the sky was screened off from the birds' view in my study.

During autumn tests under clear skies, the robins are not oriented in a seasonally appropriate migratory direction (expected direction = SW/SSW, based on ringing recoveries). A possible reason for the lack of response to E-vector manipulations may be that the birds show a nonmigratory escape orientation directed at the position of the setting sun or at the brighter part of the sky (phototactic response). This is supported by the fact that robins caught as passage migrants during autumn at FBO have low fat reserves and often engage in reoriented movements (cf. Sandberg et al., in press b). The experiments under polarizers resulted in a markedly higher degree of scatter in mean headings as compared to controls under natural twilight skies ( $r = 0.79$ ), but it was statistically significant only between controls and the E-vector = C category ( $P < 0.05$ , test for the homogeneity of concentration parameters, Mardia 1972:165-167). This may be an effect of the limited light transmission through the polarizers which in effect creates less variation in skylight intensity under E-vector = C thereby making it more difficult for the birds to detect the brighter parts of the sky. Below E-vector = C + 90°, there will be a dark band overhead, but it is still possible to recognize where the sun or the brighter part of the sky is situated (for photographs illustrating this effect, see Wehner 1976).

On the other hand, during spring tests with a clear sky overhead, the birds oriented in a seasonally appropriate direction, significantly to the right of the sunset azimuth (95% Confidence Intervals =  $\pm 24^\circ$  and  $\pm 27^\circ$ , respectively), under both polarizer conditions. Earlier experiments with artificially shifted magnetic fields have shown that the orientation of Falsterbo robins during spring is significantly influenced by manipulations of geomagnetic cues (Sandberg et al., in press a). This suggests that the spring orientation of robins at this site is related to migratory behavior. Consequently, the fact that robins under the two experimental polarizer conditions (E-vector = C and E-vector = C + 90°) select almost exactly the same directions clearly indicates that they do not use skylight polarization patterns in their migratory orientation.

Under simulated overcast, the introduction of an E-vector axis in an otherwise visually cueless situation was expected to produce clear differences in directional choice between experimental categories, if the robins use polarization patterns for their orientation. As can be seen in Figure 1, the experimental outcome is not consistent with this expectation. There are no statistical differences between experimental treatments or between birds tested under polarizers and controls. Obviously, the polarization patterns have not facilitated or changed the orientation of the robins in any decisive

way. The orientation results under simulated overcast are too scattered to allow a detailed evaluation whether there are any less pronounced effects of the imposed E-vectors.

In comparison with the clear demonstrations by Able (1982) and Moore (1986) concerning the influence of skylight polarization on the orientation of White-throated Sparrows, Kentucky Warblers, and Northern Waterthrushes, my negative results suggest that there are important differences between these species and the robin. Further support for fundamental differences between species in the mechanisms of migratory orientation is indicated by the demonstrated sensitivity of robins to the geomagnetic field (cf. Wiltschko and Wiltschko 1975; Sandberg et al., in press a), in contrast to the frequent failures to detect any involvement of magnetic cues in the orientation of North American emberizines (Moore 1985; Able and Cherry 1986; but see Bingman 1981, 1983).

I wish to thank my supervisor Thomas Alerstam for continuous encouragement, stimulating discussions, and for critically reading this manuscript. I am also very grateful to Karin Persson, Lennart Karlsson, and Göran Walinder for supplying me with robins captured at the Falsterbo Bird Observatory and to Steffi Douwes for drawing the figure. K. P. Able and F. R. Moore made useful comments on the manuscript. This study was supported by grants from the Swedish Natural Science Research Council (to T. Alerstam), and from the Swedish Ornithological Society (G. Danielsson Foundation). It is report no. 121 from the Falsterbo Bird Observatory.

#### LITERATURE CITED

- ABLE, K. P. 1982. Skylight polarization patterns at dusk influence migratory orientation in birds. *Nature* 299:550-551.
- ABLE, K. P., AND J. D. CHERRY. 1985. Laboratory and field studies of avian migratory orientation, p. 516-525. *In* M. A. Rankin [ed.], *Migration: mechanisms and adaptive significance. Contributions in marine science. Vol. 27 (Supplement)*. Port Aransas, Texas.
- ABLE, K. P., AND J. D. CHERRY. 1986. Mechanisms of dusk orientation in White-throated Sparrows (*Zonotrichia albicollis*): clock-shift experiments. *J. Comp. Physiol.* 159:107-113.
- BATSCHLET, E. 1981. *Circular statistics in biology*. Academic Press, New York.
- BINGMAN, V. P. 1981. Savannah Sparrows have a magnetic compass. *Anim. Behav.* 29:962-963.
- BINGMAN, V. P. 1983. Importance of earth's magnetism for the sunset orientation of migratory native Savannah Sparrows. *Monit. Zool. Ital.* 17:395-400.
- BRINES, M. L. 1980. Dynamic patterns of skylight polarization as clock and compass. *J. Theor. Biol.* 86:507-512.
- BRINES, M. L., AND J. L. GOULD. 1982. Skylight polarization patterns and animal orientation. *J. Exp. Biol.* 96:69-91.
- DELIUS, J., R. PERCHARD, AND J. EMMERTON. 1976. Polarized light discrimination by pigeons and an

- electroretinographic correlate. *J. Comp. Physiol. Psychol.* 90:560-571.
- KATZ, Y. B. 1985. Sunset and the orientation of European Robins (*Erithacus rubecula*). *Anim. Behav.* 33:825-828.
- KATZ, Y., AND H. MICHELSONS. 1978. (Influence of direction altering of stellar or magnetic meridians on orientation of European Robins in circular cages at planetarium), p. 182-193. In H. Michelsons, [ed.], *Orientation of birds*. Riga, Zinatne. (In Russian with English summary.)
- KREITHEN, M. L., AND W. T. KEETON. 1974. Detection of polarized light by the homing pigeon, *Columba livia*. *J. Comp. Physiol.* 89:83-92.
- MARDIA, K. V. 1972. *Statistics of directional data*. Academic Press, New York.
- MOORE, F. R. 1985. Integration of environmental stimuli in the migratory orientation of the Savannah Sparrow (*Passerculus sandwichensis*). *Anim. Behav.* 33:657-663.
- MOORE, F. R. 1986. Sunrise, skylight polarization, and the early morning orientation of night-migrating warblers. *Condor* 88:493-498.
- MOORE, F. R. 1987. Sunset and the orientation behaviour of migrating birds. *Biol. Rev. Camb. Philos. Soc.* 62:65-86.
- SANDBERG, R., J. PETTERSSON, AND T. ALERSTAM. In press a. Shifted magnetic fields lead to deflections and bimodality in the migratory orientation of Robins *Erithacus rubecula* at sunset. *Anim. Behav.*
- SANDBERG, R., J. PETTERSSON, AND T. ALERSTAM. In press b. Why do Robins *Erithacus rubecula* captured at two nearby stop-over sites orient differently? *Anim. Behav.*
- WALLRAFF, H. G., AND O. G. GELDERLOOS. 1978. Experiments on migratory orientation of birds with simulated stellar sky and geomagnetic field: method and preliminary results. *Oikos* 30:207-215.
- WEHNER, R. 1976. Polarized-light navigation by insects. *Sci. Am.* 235:106-115.
- WILTSCHKO, R., AND W. WILTSCHKO. 1978. Relative importance of stars and the magnetic field for the accuracy of orientation in night-migrating birds. *Oikos* 30:195-206.
- WILTSCHKO, W., H. HÖCK, AND F. W. MERKEL. 1971. Outdoor experiments with migrating Robins in artificial magnetic fields. *Z. Tierpsychol.* 29:409-415.
- WILTSCHKO, W., AND R. WILTSCHKO. 1975. The interaction of stars and magnetic field in the orientation system of night migrating birds. II. Spring experiments with European Robins (*Erithacus rubecula*). *Z. Tierpsychol.* 39:265-282.

*The Condor* 90:270-271  
© The Cooper Ornithological Society 1988

## RAVEN TOOL USE?

BERND HEINRICH

*Department of Zoology, University of Vermont, Burlington, VT 05405*

*Key words:* Tool use; ravens; *Corvus corax*; starvation; nutcrackers; *Nucifraga columbiana*.

Tool use by birds is well known (E. O. Wilson, 1975 *Sociobiology*, p. 172, Harvard Univ. Press, Cambridge). Recently, D. Griffin (1984 *Animal Thinking*, p. 120-121, Harvard Univ. Press, Cambridge) cites an additional "enterprising use of rocks" by Northern Ravens, *Corvus corax*. The presumptive tool use by ravens refers to observations reported by Steward W. Janes (1976 *Condor* 78:409).

Janes and a companion were climbing up a crevasse to a raven cliff nest containing six feathered young. Both adults flew at the intruders repeatedly, calling loudly, landing on the cliff above, and loosening rocks (and later also sand) and dropping them down onto the

intruders. I here report other examples of possibly similar behavior of agitated Northern Ravens dislodging substrate.

On 9 May 1987, a companion (Dick Smyth) and I were in a grove of tall (ca. 30-m) white pines near Weld, Maine, to climb to a nest of ravens containing one feathered young. On this and three subsequent occasions one of the adults of the pair immediately left the area, silently. The second became very agitated and flew within 5 m of the climber when he was near the nest. It called loudly and landed on the tree above the climber as well as on neighboring trees. The bird pecked at and snipped off about six to eight tips (2 to 3 cm) of pine twigs. These twig tips with the attached needles drifted to the ground. The bird made no apparent attempt to position itself above us while pecking at or tearing the substrate at its feet. However, the trees next to the nest afforded numerous perches.

During the previous winter (early March) I had encountered a raven in the same area at a cleanly-picked deer carcass in the snow. (The carcass had been scented

<sup>1</sup> Received 6 July 1987. Final acceptance 10 August 1987.