

# RECENT PUBLICATIONS ON WADERS 57

compiled by Henk Koffijberg

In trying to make these lists as complete as possible, and the next one even more complete and up to date, your help is most welcome. Please send new titles, abstracts, reprints, copies, reports, and (important tool) omissions or corrections to: **Henk Koffijberg, Glenn Millerweg 102, 1311 RW Almere, The Netherlands.**

Remember to include, if possible, the address of the first author, and translations of titles if they are in other languages than English. For help with this issue I am very grateful to

V.V. Morozov, T. Piersma, L.G. Underhill, B. Trolliet, A. Barbosa, G.C. Cadee, R. Deman, M. Boschert, R. Valle, D.J. Steventon, K.-M. Exo, A.-M. Blomert, M.C. Grant, J. Van Impe, and H. Hötker.

Translated titles from languages other than English are bracketed; with round brackets if the translation is by the author(s), square brackets if not.

**N.B. a copy of your paper or abstract would be very helpful to enable me to add keywords or give added annotation to the titles.**

## BREEDING

Ardamatskaya, T.S. & Kabakov, A.N. 1991 [Peculiarities of wader breeding in 1986 on the islets of Chernomorski Nature Reserve, Southern Ukraine]. *Ornitologiya* 25: 145-146. In Russian.

Boschert, M. 1990 Brutbiologie und Nahrungsökologie des Grossen Brachvogels (*Numenius arquata*) in einem Brutgebiet am südlichen Oberrhein. (Breeding biology and

feeding ecology of the Curlew (*Numenius arquata*) in a breeding area in the southern Rhine valley (Southern Germany). Thesis, University of Tübingen. In German. (Wiedigstr. 18, D-7580 Bühl, Germany). Study of breeding cycle, clutch size, breeding success, nest site selection, disturbance, territory size, feeding ecology, time-activity budget.

Colwell, M.A. 1992 Wilson's Phalarope nest success is not influenced by vegetation concealment. *Condor* 94(3): 767-772. (Biol. Dept., Univ. North Dakota, Grand Forks, ND 58201, USA). *Phalaropus tricolor*.

Crick, H.O.P. 1992 Trends in the breeding performance of Golden Plover in Britain. BTO Research Report 76. *Pluvialis apricaria*.

Estafiev, A.A. 1991 [Eurasian Oyster-catchers in European north-east of the USSR]. In: *Abstracts of the 10th All-Union Ornithological Conference, part 2, book 1*. Minsk. p. 206-207. In Russian. *Haematopus ostralegus*.

Gavrilov, V.V. 1991 (Social organization and duration difference phases of waders breeding period in tundra of lower Kolyma River). In: *Abstracts of the 10th All-Union Ornithological Conference, part 2, book 1*. Minsk. p. 130-131. In Russian.

Grant, M.C., Chambers, R.E. & Evans, P.R. 1992 The effects of re-seeding heathland on breeding whimbrel *Numenius phaeopus* in Shetland. II. Habitat use by adults during pre-laying period. *J. Appl. Ecol.* 29(2): 509-515. (Dept. Biol. Sci., Durham Univ., South Rd., Durham DH1 3LE, UK)

Grant, M.C. 1992 The effects of re-seeding heathland on breeding whimbrel *Numenius phaeopus* in Shetland. I. Nest distributions. *J. Appl. Ecol.* 29(2): 501-508.

Grant, M.C., Chambers, R.E. & Evans, P.R. 1992 The effects of re-seeding heathland on breeding whimbrel *Numenius phaeopus* in Shetland. III. Habitat-use by broods. *J. Appl. Ecol.* 29(2): 516-523.

Groen, N.M. 1992 Weidevogels in de Schaalsmeerpolder, aantalontwikkeling en broedsucces (1982-1988). [Meadowbirds in the Schaalsmeerpolder, population developments and breeding success (1982-1988)]. *Graspieper* 12(3): 94-104. In Dutch. (Paardenkamp 49, 1541 XN, Zaandam, Netherlands). Breeding waders in the province of North-Holland, Netherlands. *Haematopus ostralegus*, *Recurvirostra avosetta*, *Vanellus vanellus*, *Philomachus pugnax*, *Gallinago gallinago*, *Limosa limosa*, and *Tringa totanus*.

Ibanez, F. & Trolliet, B. 1992 Un cas de fidélité au site de nid chez la Barge "queue noire". [A case of breeding site fidelity in

Black-tailed Godwit]. *Alauda* 60(2): 111. In French. (O.N.C. Chanteloup, F-85340, L'Île D'Olonne, France). *Limosa limosa*.

Koopman, K. 1992 Biometrie, gewichtsverloop en handpenruïe van een binnenlandse populatie Scholeksters *Haematopus ostralegus*. (Biometrics, weight change and primary moult of an inland population of the Oystercatcher *Haematopus ostralegus*). *Limosa* 65(3): 103-108. In Dutch with English summary. (Diligencelaan 11, 9351 PR, Leek, Netherlands)

Kovschar, A.F. & Berezovikov, N.N. 1992 (Daylight-time life of *Burhinus oedicnemus* during reproductive period). *Bull. Moscow Soc. Naturalists (Biol. Ser.)* 97(3): 13-?. In Russian with English summary. Breeding biology of *Burhinus oedicnemus* in Kazakhstan, Alma Ata region.

Kozlov, V.P. 1991 [Little Ringed Plover in Belorussia lakes-country]. In: *Abstracts of the 10th All-Union Ornithological Conference, part 2, book 1*. Minsk. p. 289-290. In Russian.

Kretchmar, A.V., Andreev, A.V. & Kondratyev, A.Ya. 1991 (*Birds of northern plains*). Leningrad : "Nauka" Publishing House, Leningrad Branche, 288 p. In Russian.

Loegering, J.P. 1992 *Piping Plover breeding biology, foraging ecology and behavior on Assateague Island National Seashore, Maryland*. M.S. thesis, Va. Polytech. Inst. & State Univ., 248 p. *Charadrius melanotos*.

Luge, J. 1992 Zum Bestand des Grossen Brachvogels und seiner Schutzsituation im Kreis Köthen. [Breeding numbers of *Numenius arquata* and its protection in the district of Köthen]. *Beitr. Vogelkd.* 38(2): 113-117. In German. (Otto-Nuschke-Strasse 2, D-4370 Köthen, Germany). Breeding numbers of *Numenius arquata* 1964-1990.

Meininger, P.L. 1992 Kustbroedvogels en de Oosterscheldewerken. (Coastal breeding birds and the Oosterschelde engineering works). *Levende Natuur* 93(5): 152-158. In Dutch with English summary. (Rijkswaterstaat, Dienst Getijdewateren, P.O.Box 8039, 4330 EA Middelburg, Netherlands). Summary of the development in breeding populations of characteristic coastal breeding birds as *Charadrius hiaticula*, *C. alexandrinus*, *C. dubius*, and *Recurvirostra avosetta* in the Delta area of SW-Netherlands 1979-1990 after damming and other coastal engineering works.

Mnatsekanov, R.A., Emtil, M.Kh., Tilba, P.A. & Ivanenko, A.M. 1991 [Common Pratincole in the eastern coastal areas of the Azov Sea]. *Caucaskyi ornithologicheskyi vestnik* 3:131-134. In Russian. *Glareola pratincola*.

Mulder, J.L. & Swaan, A.H. 1992 Body-



- weight changes of egg-laying Curlews *Numenius arquata*, as monitored by an automatic weighing system. *Ardea* 80(2): 273-279. (IBN-DLO, P.O. Box 9201, 6800 HB Arnhem, Netherlands)
- OAG Münster 1992 Zur Situation des Kiebitzes *Vanellus vanellus* im nördlichen Münsterland : Teilaспект einer Rasterkartierung. (Situation of the Lapwing *Vanellus vanellus* in Northern Münsterland : partial aspect of a grid survey). *Vogelwelt* 113(3): 113-121. In German with English summary. (OAG Münster, Coermühle 181, 4400 Münster, Germany). Breeding distribution of *Vanellus vanellus* in large areas of Westphalia, Germany.
- O'Brien, M. & Smith, K.W. 1992 Changes in the status of waders breeding on wet lowland grasslands in England and Wales between 1982 and 1989. *Bird Study* 39(3): 165-176. (BTO, The Nunnery, Nunnery Place, Thetford, Norfolk IP24 2PU, UK). Breeding of *Vanellus vanellus*, *Gallinago gallinago*, *Tringa totanus*, *Numenius arquata*.
- Pennington, M.G. 1992 Ringed Plovers nesting in covered sites. *Brit. Birds* 85(9): 498-499. (9 Daisy Park, Baltasound, Unst, Shetland ZE2 9EA, UK). *Charadrius hiaticula*.
- Pissarev, S., Timoshenko, A. & Shapovalov, I. 1991 [Breeding of Black-winged Stilt on man-made islets]. In: *Abstracts of the 10th All-Union Ornithological Conference, part 2, book 2*. Minsk. p. 150-151. In Russian. *Himantopus himantopus*.
- Prokofyev, S.M. 1991 [Peculiarities of the Pied Avocet biology in the Minusinsk Depression]. In: *Abstracts of the 10th All-Union Ornithological Conference, part 2, book 2*. Minsk. p. 171-172. In Russian. *Recurvirostra avosetta*.
- Pulliainen, E. & Saari, L. 1992 Breeding biology of the Dotterel *Charadrius morinellus* in eastern Finnish Lapland. *Ornis Fenn.* 69(2):101-107. (Dept. Zool., Univ. Oulu, Linnanmaa, SF-90570 Oulu, Finland)
- Ranftl, H. & Dornberger, W. 1990 Brutbestand und Bruterfolg von Grossem Brachvogel, Rotschenkel und Uferschnepfe in Nordbayern 1987-1989. [Breeding populations and breeding success of Curlew, Redshank and Black-tailed Godwit in Northern Bavaria, 1987-1989]. *Anz. Ornithol. Ges. Bayern* 29: 55-61. In German. (Inst. Vogelkunde, Sandrinaweg 1, D-8825 Triesdorf-Weidenbach, Germany). *Numenius arquata*, *Tringa totanus* and *Limosa limosa* in Germany.
- Scharringa, C.J.G. 1992 Meetnet weidevogels Noord-Holland in 1992. [Monitoring meadowbirds in the province of North-Holland in 1992]. *Graspieper* 12(4): 134-140. In Dutch. (Provincie Noord-Holland, Dienst Ruimte en Groen, Afd. Onderzoek en Informatie, P.O. Box 6090, 2001 HB Haarlem, Netherlands). Breeding numbers of *Haematopus ostralegus*, *Vanellus vanellus*, *Limosa limosa*, *Tringa totanus*, *Philomachus pugnax*, and *Gallinago gallinago*.
- Valle, R. & D'Este, A. 1992 (Three years of ornithological investigations in the area of Lido Harbour (Venice) with notes on the breeding biology of Kentish Plover (*Charadrius alexandrinus*) and White Wagtail (*Motacilla alba*)). *Lavori* 17: 121-129. In Italian with English summary. (Castello 618/E, 30122 Venezia, Italy). Breeding of *Charadrius alexandrinus*, and wintering of *C. hiaticula* and *Calidris alpina* in the Venice Lagoon, Italy.
- Witt, K. 1992 Bestandsentwicklungen einiger ausgewählter gefährdeter Vogelarten im westlichen Deutschland 1977-1989. (Population changes of some selected endangered breeding bird species of western Germany 1977 to 1989). *Vogelwelt* 113(6): 289-310. In German with English summary. (Hortensienstr. 25, D-1000 Berlin 45, Germany). Breeding waders: *Limosa limosa*, *Numenius arquata*, *Tringa totanus*, *Tringa ochropus*.
- BEHAVIOUR, ECOLOGY & EVOLUTION**
- Alberico, J.A.R., Reed, J.M. & Oring, L.W. 1992 Non-random philopatry of sibling Spotted Sandpipers *Actitis macularia*. *Ornis Scand.* 23(4): 504-508. (Ecology, Evolution and Conservation Biology Program, RWF 1000 Valley, Rd., Univ. Nevada, Reno, NV 89512, USA). Philopatry data from a long-term study of Spotted Sandpipers were used to test the hypothesis that yearling returns were independent, and to evaluate explanations for non-random return.
- Alerstam, T., Gudmundsson, G.A. & Johannesson, K. 1992 Resources for long distance migration : intertidal exploitation of *Littorina* and *Mytilus* by knots *Calidris canutus* in Iceland. *Oikos* 65(2): 179-189. (Dept. Ecol., Ecology Building, Univ. Lund, S-223 62 Lund, Sweden)
- Andreeva, T.R. 1991 (Trophic relations of the Terek Sandpiper (*Xenus cinereus*) in the Stchuch River basin, Yamal Peninsula). *Ornitologiya* 25: 35-39. In Russian with English summary.
- Andreeva, T.R. & Tomkovich, P.S. 1992 (On the role of chick food items in populations structure formation of tundra waders). In: *Cenotic interactions in tundra ecosystems: Collection of papers*. Moscow : Nauka, p. 70-78. In Russian with English summary.
- Bacetti, N., De Faveri, A. & Serra, L. 1992 Spring migration and body condition of Common Sandpipers *Actitis hypoleucus* on a small Mediterranean island. *Ring. & Migr.* 13(2): 90-94. (Istituto Nazionale di Biologia della Selvaggina "A. Ghigi", via Ca Forncasetta 9, 40064 Ozzano Emilia BO, Italy). Information on body mass and timing of spring migration of *Actitis hypoleucus*.
- Baines, D., Grant, M., Jackson, D.B. & Evans, P.R. 1991 Effects of habitat loss through changes in agricultural practice on waders breeding in western Europe. *Acta XX Congress. Intern. Ornithol. Vol. IV*: 2236. (Dept. Biol. Sci., Univ. Durham, South Rd., Durham DH1 3LE, UK)
- Blitzblau, S. 1990 [Migration of waders along the Carmel Coast, spring 1990]. *Torgos* 8(2): 59-68, 73. In Hebrew with English summary. (SPNI Carmel Coast Field Study Ctr., SPNI Ma'agan-Mikhael, D.N. Monashe 37805, Israel). *Calidris minuta* and *Tringa glareola* in Israel.
- Cadée, G.C. 1989 Size-selective transport of shells by birds and its palaeoecological implications. *Palaeontology* 32(2): 429-437. (NIOZ, P.O. Box 59, 1790 AB Den Burg, Texel, Netherlands). Size-selective transport of shells is demonstrated in the Dutch Wadden Sea for *Haematopus ostralegus*: only larger shells were transported.
- Calinin, S.S. 1991 [Impact of abiotic factors on egg-laying and morphologic characteristic wader eggs in areas eastern to the Ural Mountains and Kama River basin near Ural Mountains]. In: *Ornithological problems of Siberia. Abstracts to the Conference*. Barnaul. p. 15-17. In Russian.
- Ctyroky, P. 1992 Der Weissenschwanzsteppenkiebitz (*Chettusia leucura* [Lichtenstein]) - ein endemischer Kiebitz Südwestasiens. [White-tailed Plover (*Chettusia leucurus*) - an endemic plover of southwest Asia]. *Beitr. Vogelkd.* 38(2): 92-98. In German. (Na Julisce 13, CS-16000 Praha 6-Dejvice, Czechoslovakia). Observations on *Vanellus leucurus* in Iraq.
- Dubois, P.J. 1992 Migration et hivernage de l'Echasse blanche (*Himantopus himantopus*) dans l'Ouest du Paléarctique et de l'Afrique. (Migration and wintering of the Black-winged Stilt in the west of the Western Palearctic). *Nos Oiseaux* 41(6): 347-366. In French with German and English summaries (29 Rue Saint Didier, Epiais Rhus, 95810 Frizyles, Plalettes, France). Breeding Stilt population of Italy, France, Spain, Portugal and the Magreb is estimated at about 12000 pairs, the total population being between 36000 and 44000 individuals in early autumn and reducing to about 30000 birds during winter.
- Evans, P.R., Davidson, N.C., Uttley, J.D. & Evans, R.D. 1992 Premigratory hypertrophy of flight muscles : an ultrastructural study. *Ornis Scand.* 23(3): 238-243. (Dept. Biol. Sci., Univ. Durham, Sci. Lab., South Rd., Durham, England DH1 3LE). Results are



described of studies on the fine structure and composition of pectoral muscles of *Calidris alpina*, *C. alba*, and *C. canutus*. The studies were undertaken to identify the functions of flight muscle hypertrophy.

Evans, P.R., Davidson, N.C., Piersma, T. & Pienkowski, M.W. 1991 Implications of habitat loss at migration staging posts for shorebird populations. *Acta XX Congress. Intern. Ornithol.* Vol. IV: 2228-2235.

Gavrilov, E.I. 1991 (Energetical maintenance in the Little Stint (*Calidris minuta* Leisler) during autumn migration in Kazakhstan). *Ornitologiya* 25: 15-25. In Russian with English summary.

Gerasimov, Yu.N. 1991 [Spring wader migration in Western Kamchatka]. In: *Abstracts of the 10th All-Union Ornithological Conference, part 2, book 1*. Minsk. p. 142. In Russian.

Gorban, I.M., Bokotei, A.A. & Pogranichnyi, V.A. 1991 (Autumn migration of waders in Lvov and Volyn Regions, western Ukraine). *Ornitologiya* 25: 190-191. In ussian.

Gordienko, N.S. 1991 (Biology and number of the Sociable Lapwing (*Chettusia gregaria*) in Kustanai Steppe, Northern Kazakhstan). *Ornitologiya* 25: 54-61. In Russian with English summary.

Goss-Custard, J.D. & Cayford, J.T. 1990 How the foraging behaviour of Oystercatchers, *Haematopus ostralegus*, changes with age. In: *Proceedings of the International Centennial Meeting of the Deutsche Ornithologen Gesellschaft. Current topics in Avian Biology*. p. 335-340. (Furzebrook Res. Stn., Inst. Terrestrial Ecol., Wareham, Dorset BH20 5AS, UK)

Granval, P. 1992 [Worm predation by birds : an example, the Woodcock *Scolopax rusticola*]. *Alauda* 59(1): 40. In French. (O.N.C. Avifaune Migatrice, Lab. Zoecol. sol (INRA), Cent. Ecol Fonctionnelle & Evol.(C.N.R.S.), B.P. 5051, 34033 Montpellier Cedex, France)

Hoglund, J., KŒlEs, J.A. & Fiske, P. 1992 The costs of secondary sexual characteristics in the lekking Great Snipe (*Gallinago media*). *Behav. Ecol. Sociobiol.* 30(5): 309-315. (Dept. Zool., Uppsala Univ., Box 561, S-751 22 Uppsala, Sweden)

Van Impe, J. 1992 Waarnemingen bij een gemeenschappelijke slaapplaats van Regenwulpalen *Numenius phaeopus* en Wulpalen *N. arquata*. (Observations at a collective roost in Whimbrel *Numenius phaeopus* and Curlew *N. arquata*). *Oriolus* 58(1): 1-5. In Dutch with English summary. (K.B.I.N., Vautierstraat 29, B-1040 Brussel, Belgium). During the spring of four consecutive years 1988-1991, time of arrival and activity among

Whimbrel and Curlew were quantitatively and comparatively investigated at a collective roost situated on a man-made deposit on the left bank of the river Scheldt near Antwerp, Belgium.

Jukema, J. & Piersma, T. 1992 Mannelijke en vrouwelijke Goudplevieren *Pluvialis apricaria* zijn wat betreft hun uitwendige en inwendige afmetingen volkomen identiek. (Golden Plovers *Pluvialis apricaria* do not show sexual dimorphism in external and internal linear dimensions). *Limosa* 65(4): 147-154. In Dutch with English summary. (Haerdawei 44, 8854 AC Oosterbierum, Netherlands)

Kaletja, B. 1992 *Aspects of the ecology of migrant shorebirds (Aves: Charadrii) at the Berg River estuary, South Africa*. Ph.D thesis, University of Cape Town

Kasperek, M. 1992 Zum Zug des Steinwülers, *Arenaria interpres*, in der Türkei. [Migration of Turnstone, *Arenaria interpres*, in Turkey]. *Beitr. Vogelkd.* 38(3): 200-208. In German with English summary. (Bleichstr. 1, W-6900 Heidelberg, Germany). *Arenaria interpres* is a regular but uncommon passage migrant through Turkey. Migration periods extend from late March to late May and from mid-July to late October.

Korzukov, A.I. & Petrovskaya, L.P. 1991 (Distribution and biology of *Himantopus himantopus* in the North-Western Black Sea coastal areas). In: *Rare birds of the Black Sea Coastal Area*, Kiev-Odessa. p. 165-181. In Russian with English summary.

Lank, D.B. & Smith, C.M. 1992 Females prefer larger leks : field experiments with Ruffs (*Philomachus pugnax*). *Behav. Ecol. Sociobiol.* 30(5): 323-329. (Dept. Biol., Queen's Univ., Kingston, ON, Canada K7L 3N6)

Marks, J.S. 1992 Longevity record for the Bristle-thighed Curlew : an extension. *J. Field Ornithol.* 63(3): 309-310. With Spanish summary. (Div. Biol. Sci., Univ. Mont., Missoula 59812, USA). *Numenius tahitiensis* in Hawaii.

Van der Merwe, D., McLachlan, A. & De Ruyck, A.M.C. 1992 Partitioning of feeding habitats by Whitefronted Plovers *Charadrius marginatus* at a dune/beach interface. *Ostrich* 63: 49-57. (Coastal Res. Inst. & Dept. Zool., Univ. Port Elizabeth, P.O. Box 1600, Port Elizabeth 6000, South Africa)

Morris, P.I. 1992 Escape reaction of Knot from Peregrine. *Brit. Birds* 85(8): 465. (Caughall Farmhouse Cottage, Caughall Rd., Upton-by-Chester, Cheshire CH2 4BW, UK). *Calidris canutus*.

Ostatshenko, A.N. 1991 [The artificial irrigation influence on shorebirds in Chu River

valley of Kirgizia]. In: *Abstracts of the 10th All-Union Ornithological Conference, part 2, book 2*. Minsk. p. 130. In Russian.

Pannach, G. 1992 Phänologische Untersuchungen des Limikolenzuges 1979 bis 1991 in den Braunschweiger Rieselfeldern. (Phenological investigations of the wader migrations in the brunswick sewage farm (Lower Saxony) from 1979 to 1991). *Braunschweig. naturkd. Schr.* 4(1): 27-57. In German with English summary. (Oppelnstr. 17, D-3300 Braunschweig, Germany). Characteristic, distribution, winter ground, home direction, off migration and a comparative avifaunistic are described. (Paper contains 147 lit. refs.)

Sallaberry, M.A. 1991 *Genetic and morphological differentiation of Sanderling () populations on the wintering grounds*. Ph.D. dissertation, Univ. Pa., 83 p. From Diss Abstr. Int. B Sci. Eng. 51(12):5762, Order no. DA9112616.

Shubin, A.O. 1991 [Visual wader migration on the western coast of the Caspian Sea]. In: *Abstracts of the 10th All-Union Ornithological Conference, part 2, book 2*. Minsk. p. 300-301. In Russian.

Smith, K.W., Reed, J.M. & Trevis, B.E. 1992 Habitat use and site fidelity of Green Sandpipers *Tringa ochropus* wintering in Southern England. *Bird Study* 39(3): 155-164. (RSPB, The Lodge, Sandy, Bedfordshire SG19 2DL, UK). Study of seasonal patterns of site usage, site fidelity and return rates of a colour marked population of *Tringa ochropus* wintering in an extensive area.

Tomkovich, P.S. 1991 (Relation between breeding dates and parental care in some monogamous waders). In: *Abstracts of the 10th All-Union Ornithological Conference, part 2, book 2*. Minsk. p. 251-252. In Russian.

Tomkovich, P.S. 1991 (External morphology of the Spoon-billed Sandpiper (*Eurynorhynchus pygmeus*) at Chukotski Peninsula). *Ornitologiya* 25: 135-144. In Russian with English summary.

Velasquez, C.R. & Hockey, P.A.R. 1992 The importance of supratidal foraging habitats for waders at a south temperate estuary. *Ardea* 80(2): 243-253. (Percy FitzPatrick Inst. African Ornithol., Univ. Cape Town, Rondebosch 7700, South Africa). This study quantifies the extent to which waterbirds, mainly waders, utilize foraging areas other than intertidal mudflats during high and low tides at the Berg River estuary, South Africa.

Wenink, P.W., Smit, C.J., Tilanus, M.G.J., Muiswinkel, W.B. van & Baker, A.J. 1992 DNA-analyse : achter de grenzen van de biometry. (DNA analysis : beyond the limits of morphometrics). *Limosa* 65(3): 109-115.



- In Dutch with English summary. (Johan de Wittlaan 99, 6828 XE, Arnhem, Netherlands). A method of DNA analysis has been developed in order to identify subspecies and populations of wader species. This is exemplified by breeding *Calidris alpina* from Iceland, Norway, and Siberia.
- Yalden, D.W. & Holland, P.K. 1992 Relative contributions of Common Sandpiper *Actitis hypoleucus* parents to guarding their chicks. *Ring. & Migr.* 13(2): 95-97. (Dept. Environ. Biol., Univ. Manchester, M13 9PL, UK)
- Zwarts, L. & Blomert, A.-M. 1992 Why Knot *Calidris canutus* take medium-sized *Macoma balthica* when six prey species are available. *Mar. Ecol. Prog. Ser.* 83: 113-128. (Rijks-waterstaat Flevoland, P.O.Box 600, 8200 AP Lelystad, Netherlands). Prey selection and intake rate of *Calidris canutus* has been quantified when 6 different, intertidal prey species were abundant.
- Zwarts, L., Blomert, A.-M. & Wanink, J.H. 1992 Annual and seasonal variation in the food supply harvestable by Knot *Calidris canutus* staging in the Wadden Sea in late summer. *Mar. Ecol. Prog. Ser.* 83: 129-139.
- DISTRIBUTION & FAUNISTICS**
- Ardamatskaya, T.B. 1991 (Rare and disappearing birds of the Ukrainian Black Sea State Biosphera Reservation and its neighbouring territories). In: *Rare birds of the Black Sea Coastal Area*, Kiev-Odessa. p. 54-68. In Russian with English summary.
- Beckon, W.N. 1990 First record of the Pectoral Sandpiper in Fiji. *Notornis* 37(2): 106. (Dept. Zool., Univ. California, Davis 95616, USA). *Calidris melanotos*.
- Blokhin, Yu.Yu. 1991 (On the northern limits of breeding of some birds on the northern Lena River). *Ornitologiya* 25: 145-150. In Russian.
- By, R.A. de, Berg, A.B. van den & CDNA 1992 Zeldzame en schaarse vogels in Nederland in 1990. (Rare and scarce birds in The Netherlands in 1990). *Limosa* 65(4): 137-146. In Dutch with English summary. Waders: *Pluvialis fulva*, *Chettusia gregaria*, *Calidris melanotos*, *Limicola falcinellus*, *Tringitea subruficollis*, *Limnodromus scolopaceus*, *Tringa stagnatilis*.
- Deman, R. 1992 De overwintering van de Bonte Strandloper in het natuurreervaat "Groot Buitenschoor" te Zandvliet. [Wintering of *Calidris alpina* in the nature reserve "Groot Buitenschoor" near Zandvliet]. *Wielewaal* 58: 164-165. In Dutch. (Kwade Veldenplein 8 bus 3, B-2170, Antwerpen-Merksem, Belgium). Numbers of wintering *Calidris alpina*, 1981-1992, in a nature reserve near Antwerp, Belgium.
- Dorogoi, I.V. 1991 (To the fauna and distribution of birds at North-eastern Chukotski Peninsula). *Ornitologiya* 25: 102-109. In Russian.
- Dunnett, J.B. 1992 Long-toed Stint : new to Britain and Ireland. *Brit. Birds* 85(8): 431-436. (43 Hemlington Rd., Stainton, Middlesbrough, Cleveland TS8 9AG, UK). *Calidris subminuta*. With colour plates of juvenile.
- Eigenhuis, K.J. 1992 Grote Kanoet in Oostvaardersplassen en bij Camperduin in september-oktober 1991. (Great Knot in Oostvaardersplassen and near Camperduin in September-October 1991). *Dutch Birding* 14(4): 126-131. In Dutch with English summary. (Seringstr. 6, 1431 BJ Aalsmeer, Netherlands). *Calidris tenuirostris* in The Netherlands.
- Ellis, P.M. 1992 Great Knot : new to Britain and Ireland. *Brit. Birds* 85(8): 426-430. (Seaview, Sandwich, Shetland ZE2 9HP, UK). *Calidris tenuirostris*.
- Emtil, M.Kh. ...[et al.] 1991 [Recent status of Black-winged Stilt breeding population in Krasnodar Region]. *Caucaskyi ornithologicheskyi vestnik* 3: 77-79. In Russian. *Himantopus himantopus*.
- Faanes, C.A. 1990 A recent record of Eskimo Curlew in Nebraska. *Prairie Nat.* 22(2): 137-138. (U.S. Fish & Wildl. Serv., 203 W. Second St., Grand Island, NE 68801, USA). *Numenius borealis*.
- Fraser, M., McMahon, L., Harris, M. & Wanless, S. 1992 American Golden Plover at the Cape of Good Hope nature reserve. *Promerops* No. 203, p. 11. *Pluvialis dominica* in South Africa.
- Fraser, P.A. & Ryan, J.F. 1992 Scarce migrants in Britain and Ireland. Part 1. Numbers during 1986-90 : seabirds to waders. *Brit. Birds* 85(12): 631-635. (12 Redland Court Rd., Bristol BS6 7EQ, UK). Waders: *Burhinus oedicnemus*, *Charadrius alexandrinus*, *C. morinellus*, *Calidris temminckii*, *C. melanotos*, *Tringites subruficollis*, *Phalaropus lobatus*, *P. fulicarius*.
- Gill, R.E. Jr. & Redmond, R.L. 1992 Distribution, numbers, and habitat of Bristle-thighed Curlews (*Numenius tahitiensis*) on Rangiroa Atoll. *Notornis* 39(1): 17-26. (U.S. Fish & Wildl. Serv., Alaska Fish & Wildl. Res. Ctr., 1011 E. Tudor Rd., Anchorage, Alaska 99503, USA)
- Gorban, I.M. 1991 [The status of Eurasian Curlew in western Ukraine]. In: *Abstracts of the 10th All-Union Ornithological Conference, part 2, book 1*. Minsk. p. 161-162. In Russian.
- Grishanov, G.V. 1991 [Waders in the man-made landscapes destruction in the Kalinograd Region]. In: *Abstracts of the 10th All-Union Ornithological Conference, part 2, book 1*. Minsk. p. 168-169. In Russian.
- Gunderson, M., Gunderson, E., Lease, E. & Moriarty, P. 1992 Nesting American Avocet in Big Stone County. *Loon* 64(2): 125-126. (463 Marshall Ave., St. Paul, MN 55102, USA). *Recurvirostra americana* in Minnesota.
- Helbig, A.J., Barth, R. & Bauer, H.-G. 1991 [First record of Spotted Sandpiper *Actitis macularia* for Austria]. *Limicola* 5(6): 299-302. In German with English summary. (Inst. Pharm. Biol., Univ. Heidelberg, Im Neuenheimer Feld 364, D-6900 Heidelberg, Germany)
- Histsov, A.P. & Ivasenko, A.N. 1991 [Shorebird numbers on the north-eastern coast of the Caspian Sea]. In: *Ornithological problems of Siberia. Abstracts to the Conference*. Barnaul. p. 139-140. In Russian.
- Hockey, P.A.R. & SAOS Rarities Committee 1992 Rare birds in South Africa 1989-1990. *Birding in South Africa* 44(2): 38-44. (Percy Fitzpatrick Inst. African Ornithol., Univ. Cape Town, Rondebosch 7700, South Africa).
- Waders: *Haematopus ostralegus*, *Charadrius mongolus*, *C. asiaticus*, *Pluvialis dominica*, *P. dominica/fulva*, *Tringa ochropus*, *T. totanus*, *Calidris melanotos*, *Limosa falcinellus*, *L. limosa*, *L. haemastica*, *Phalaropus fulicarius*.
- Hötker, H. & Dietrich, S. 1991 *Rapport dé'tude sur l'é'cologie des Avocettes et d'autre oiseaux d'eau dans le delta du Sénégal, en 1988. Considérations particulières pour la Réserve Spéciale de Faune de Guembeul*. (Report on ecological studies on Avocets and other wetland birds in the delta of River Senegal in 1988, with special reference to the 'Réserve Spéciale de Faune de Guembeul'). Husum : report, 76p. In French with English summary. (c/o WWF-Wattenmeerstell, Norderstr. 3, 2250 Husum, Germany). Total wintering population of *Recurvirostra avosetta* was estimated to be 8000 birds. Most of them wintered in the lagoon system of the lower part of the delta of the River Senegal. Other ornithological observations are given as well.
- Hut, R.M.G. van der 1992 Aantalsontwikkeling van weidevogels in de winter : Wulp, Goudplevier en Kievit in de Zaanstreek, 1976-1991. [Numbers of wintering meadowbirds : Curlew, Golden Plover and Lapwing in the Zaan area, 1976-1991]. *Graspieper* 12(4): 146-154. In Dutch. (Geersbroekstr. 13, 1507 KR, Zaandam). *Numenius arquata*, *Vanellus vanellus* and *Pluvialis apricaria* in the province of North-Holland, Netherlands.
- Ivey, G. & Baars, C. 1990 A second Semi-palmated Plover nest in Oregon. *Oreg. Birds* 16(3): 207-208. (Malheur Natl. Wildl.



- Refuge, HC-72, Box 245, Princeton, OR 97721, USA). *Charadrius semipalmatus*.
- Jönsson, P.E. 1992 (The Kentish Plover (*Charadrius alexandrinus*) in Scania, South Sweden, 1991 - a report from a conservation project). *Anser* 31(1): 1-10. In Swedish with English summary. (Dept. Ecol., Ecology Building, S-223 62 Lund, Sweden)
- Khokhlov, A.N. & Kulikov, V.G. 1991 [Summer avifauna of the northern Stavropol Region]. In: *Fauna, number populations and ecology of birds of Northern Caucasus (Data of Scientific-practical conference 23-27 April 1991)*. Stavropol. p. 30-35. In Russian.
- Khokhlov, A.N., Telpov, V.A. & Bicherev, V.N. 1991 [Winter avifauna of Kislovodsk city and its vicinity (Stavropol Region)]. In: *Fauna, number populations and ecology of birds of Northern Caucasus (Data of Scientific-practical conference 23-27 April 1991)*. Stavropol. p. 123-135. In Russian.
- Koshelev, A.I., Korzukov, A.I. & Lobkov, V.A. 1991 (The analysis of the amount of rare species of birds in the district of Odessa). In: *Rare birds of the Black Sea Coastal Area, Kiev-Odessa*. p. 9-36. In Russian.
- Kovshar, A.F. 1991 [Ibisbill in Kazakhstan]. In: *Abstracts of the 10th All-Union Ornithological Conference, part 2, book 1*. Minsk. p. 285-286. In Russian. *Ibidorhyncha struthersii*.
- Kydyraliev, A. & Kydyralieva, R.A. 1991 [Waders in Kyrgyzstan (species composition and habitat distribution)]. In: *Abstracts of the 10th All-Union Ornithological Conference, part 2, book 2*. Minsk. p. 14-15. In Russian.
- Lebed, E.A., Knysh, N.P. & Khomenko, S.V. 1991 [About some rare waders of Sumy Region (Ukraine)]. In: *Abstracts of the 10th All-Union Ornithological Conference, part 2, book 2*. Minsk. p. 28-29. In Russian.
- Luckner, G.D. 1992 Marbled Godwit. *Spoonbill* 41(8): 7. *Limosa fedoa* in Texas.
- Lysenko, V.I. & Siokhin, V.D. 1991 (The present state of the amount and distribution of rare birds of the Northern Sea of Azov coastal areas). In: *Rare birds of the Black Sea Coastal Area, Kiev-Odessa*. p. 69-78. In Russian.
- Marrack, P.A. 1992 Nearctic vagrant waders in the Cap Lopez area, Gabon. *Malimbus* 14: 7-10. With English and French summaries. (Van Alkemadeaan 350/401, 2597 AS Den Haag, Netherlands). *Bartramia longicauda*, *Calidris melanotos*, *Tryngites subruficollis*, and a possible *Pluvialis dominica*.
- Martin, P. & Martin, J. 1992 Too late the Phalarope. *Bee-eater* 43(1): 6-7. *Phalaropus fulicarius* in South Africa.
- Molodan, G.N., Bukreev, S.A., Dyakov, V.A., Zalevsky, V.D., Kabakov, A.N., Pozhidaeva, S.I. & Chernyshov, A.I. 1991 (New data about distribution amount and biology of near-water nonsparrowlike birds of the northern coastal areas of Azov Sea). In: *Rare birds of the Black Sea Coastal Area, Kiev-Odessa*. p. 193-211. In Russian.
- OAG Münster 1992 Wegzugbestände des Kampflüfers *Philomachus pugnax* 1990 in Deutschland. (Numbers of Ruffs during autumn migration in Germany 1990). *Vogelwelt* 113(3): 102-113. In German with English summary. (OAG Münster, Coermühle 181, 4400 Münster, Germany).
- Nakhodkin, N.A. & Issaev, A.P. 1991 [Ornithological study of Alyssardaakh Lakes (Yakutia)]. In: *Ornithological problems of Siberia. Abstracts to the Conference*. Barnaul. p. 150-151. In Russian. Records of *Numenius madagascariensis*, *N. phaeopus*, *N. minutus*.
- Namassivayan, L. & Venugopalan, R. 1989 Avocet *Recurvirostra avosetta* in Kerala. *J. Bombay Nat. Hist. Soc.* 86(3): 447.
- Namassivayan, L., Uthaman, P.K. & Venugopalan, R. 1989 Four additions to the birds of Kerala. *J. Bombay Nat. Hist. Soc.* 86(3): 458-459. *Calidris temminckii*, *C. alpina*, *Limosa lapponica*, *Haematopus ostralegus*
- Nechaev, V.A. 1991 (Records of new and rare birds in Sakhalin Island). *Ornitologiya* 25: 167-168. In Russian.
- Nikiforov, M.E., Kozulin, A.V., Yaminski, B.V. & Zuynok, S.V. 1991 (New data on breeding of the Ringed Plover (*Charadrius hiaticula*), the Oystercatcher (*Haematopus ostralegus*) and the Terek Sandpiper (*Xenus cinereus*) in Belorussia). *Ornitologiya* 25: 168-169. In Russian.
- Norman, D. & Coffey, P. 1992 The importance of the Mersey estuary for waders in the cold weather of February 1991. In: Barber, S. & D.J. Steventon (eds.). *Cheshire and Wirral Bird Report 1991*, p. 85-88. (Rowswood Cottage, Riddings Lane, Sutton Weaver, Runcorn, Cheshire WA7 6PF, UK). Weight changes of *Tring totanus*, *Arenaria interpres* and *Calidris alpina* were examined during a cold spell.
- Osipova, M.A. & Golovushkin, M.I. 1991 [New data on Oriental Pratincole in southern Transbaikalia]. In: *Abstracts of the 10th All-Union Ornithological Conference, part 2, book 2*. Minsk. p. 127-128. In Russian.
- Panchenko, V.A. & Balatskyi, K.L. 1991 (Rare and disappearing birds of the delta of the Danube, the Dniester and the neighbouring territories). In: *Rare birds of the Black Sea Coastal Area, Kiev-Odessa*. p. 37-53. In Russian.
- In Russian.
- Patrikeev, M.V. 1991 [On spring-summer avifauna of south-eastern Shirvan and adjacent territories (Azerbaijan)]. In: *Fauna, number populations and ecology of birds of Northern Caucasus (Data of Scientific-practical conference 23-27 April 1991)*. Stavropol. p. 30-35. In Russian.
- Patrikeev, M.V. 1991 [About birds of southern Makhmud-Chala Lake and its vicinity (southern Azerbaijan)]. In: *Fauna, number populations and ecology of birds of Northern Caucasus (Data of Scientific-practical conference 23-27 April 1991)*. Stavropol. p. 60-61. In Russian.
- Payevski, V.A. & Karelina, L.D. 1991 (Ornithological observations on the south-eastern coast of Onega Gulf, White Sea). *Ornitologiya* 25: 169-170. In Russian.
- Prys-Jones, R.P., Corse, C.J. & Summers, R.W. 1992 The role of the Orkney Islands as a spring staging post for Turnstones *Arenaria interpres*. *Ring. & Migr.* 13(2): 83-89. (Estuaries Unit, BTO, The Nunnery, Nunnery Place, Thetford, Norfolk IP24 2PU, UK). Patterns of change in numbers and mass of Turnstones in two parts of Orkney during winter and spring were compared.
- Rahmani, A.R. & Manakadan, R. 1989 Breeding records of Cream-coloured Courser *Cursorius cursor* (Latham) from India. *J. Bombay Nat. Hist. Soc.* 86(3): 447-448. *Calidris temminckii*, *C. alpina*, *Limosa lapponica*, *Haematopus ostralegus*
- Reay, P. 1991 Wintering Avocets in Devon and Cornwall. *Devon Birds* 44(1): 3-11. *Recurvirostra avosetta*.
- Ruitenbeek, W. 1992 Waarnemingen van de Morinelpievier in Noord-Holland van augustus 1988 tot en met juli 1990. [Records of Dotterel in the province of North-Holland from August 1988 - July 1990]. *Graspieper* 12(3): 86-88. In Dutch. (Nassauplein 17, 1815 GN Alkmaar, Netherlands). *Eudromias morinellus* in North-Holland, Netherlands.
- Schekkerman, H., Meininger, P. & Meire, P. 1992 Watervogels en de Oosterschelde-werken. (Waterbirds and coastal engineering works in the Oosterschelde). *Levende Natuur* 93(5): 147-152. In Dutch with English summary. (DLO-IBN, P.O.Box 9201, 6800 HB Arnhem, Netherlands). Effects of a storm-surge barrier and two compartment dams in the Oosterschelde/Krammer-olkerak estuary (SW Netherlands) on birds have been studied.
- Shubin, A.O. 1991 [Autumn stay of migrating waders in Kyzyl-Agach Nature State Reserve]. In: *Abstracts of the 10th All-Union Ornithological Conference, part 2, book 2*. Minsk. p. 301-302. In Russian.



- Smirenskyi, S.M., Sumyaa, D. & Boldbaator, C. 1991 (Ornithological investigations in Dornod Aimak, Mongolian People's Republic). *Ornitologiya* 25: 116-126. In Russian with English summary.
- Smit, C.J. 1989 Perspectives in using shorebird counts for assessing long-term changes in wader numbers in the Wadden Sea. *Helgoländer Meeresuntersuchungen* 43: 367-383. (IBN-DLO, P.O. Box 167, 1790 AB Den Burg, Texel, Netherlands)
- Smith, T.C. Jr. 1990 A Ruff sight record for South Carolina. *Chat* 54(2): 64. (313 Abercorn St., Savannah, GA 31401, USA). *Philomachus pugnax*.
- Smith, F.T.H. 1992 An aberrant Red-capped Plover *Charadrius ruficollis* at Lake Connewarre, Victoria. *Aust Bird Watcher* 14(7): 260-261. (71 Cobden St., Kew, Vic. 3101, Australia)
- Spek, A. van der & Spek, V. van der 1992 Kleine Geelpootruiter bij Flaauwersinlaag in oktober 1991. (Lesser Yellowlegs near Flaauwersinlaag in October 1991). *Dutch Birding* 14(2): 50-52. In Dutch with English summary. (Dasseburcht 39, 3892 WC Zeeuwse, Netherlands). *Tringa flavipes* in SW Netherlands.
- Stashaitis, Yu.N. 1991 [Golden Plover in Lithuania]. In: *Abstracts of the 10th All-Union Ornithological Conference, part 2, book 2*. Minsk. p. 229. In Russian. *Pluvialis apricaria*.
- Stishov, M.S. 1991 [New data on Buff-breasted Sandpiper distribution in the USSR]. In: *Abstracts of the 10th All-Union Ornithological Conference, part 2, book 2*. Minsk. p. 230-231. In Russian. *Tryngites subruficollis*
- Stishov, M.S. 1991 [New data on some bird species distribution at the arctic coast of Chukotski Peninsula]. In: *Ornithological problems of Siberia. Abstracts to the Conference*. Barnaul. p. 132-133. In Russian.
- Stishov, M.S., Pridatko, V.I. & Baranyuk, V.V. 1992 [*The birds of Wrangel Island*]. Novosibirsk : Nauka, 252p. In Russian.
- Stotz, D.F., Bierregaard, R.O., Cohn-Haft, M., Petermann, P., Smith, J., Whittaker, A. & Wilson, S.V. 1992 The status of North American migrants in central Amazonian Brazil. *Condor* 94(3): 608-621. (Museu de Zoologia, Universidade de São Paulo, SP, CEP 01064, Brasil). Important migration route for: *Pluvialis dominica*, *Calidris fuscicollis*, *Calidris melanotos*.
- Sumyaa, D. & Skryabin, N.G. 1989 [*The birds of Khubsugul Lake and adjacent territories, the Mongolian People's Republic*]. Irkutsk. 198p. In Russian.
- Tomkovich, P.S. 1992 Spoon-billed Sandpiper in north-eastern Siberia. *Dutch Birding* 14(2): 37-41. (Zool. Mus., Moscow State Univ., Herzen St. 6, 103009 Moscow K-9, Russia). *Eurynorhynchus pygmaeus*.
- Tomkovich, P.S., Masterov, V.B. & Soloviev, M.Yu. 1991 (To avifauna of Shmidt Cape, Chukchi Sea). *Ornitologiya* 25: 175-176. In Russian.
- Tomkovich, P.S., Syroechkovskyi, E.E., Soloviev, M.Yu., Esu, P. & Lappo, E.G. 1991 [New data on avifauna of northern Taimir]. In: *Ornithological problems of Siberia. Abstracts to the Conference*. Barnaul. p. 134-135. In Russian.
- Trolliet, B., Girard, O., Fouquet, M., Ibanez, F., Triplet, P. & Léger, F. 1992 L'Effectif de Combattants *Philomachus pugnax* hivernant dans le delta du Sénégal. [Census of Ruff wintering in the Senegal delta]. *Alauda* 60(3): 159-163. In French with English summary. (O.N.C. Chanteloup, F-85340 Ile d'Olonne, France). Wintering numbers in January of 1991 and 1992, 180 000 and 200 000 respectively.
- Underhill, L.G. 1992 Waders (*Charadrii*) and other shorebirds at Dassen Island, and the Algoa Bay islands, South Africa. *Ostrich* 63: 128-129. (Avian Demography Unit, Dept. Stat. Sci., Univ. Cape Town, Rondebosch 7700, South Africa)
- Warbrick, S., Clark, N.A. & Donald, P.F. 1991 *Distribution studies of Waders and Shelduck on the Severn Estuary with reference to sediment mobility*. BTO Research Report 78.
- Waters, R. 1992 Wintering waders in the United Kingdom 1991-1992. *BTO News* 183: 10-11. Summary of results of the Birds of Estuaries Enquiry.
- Zhuravlev, V.E., Podkopyrov, V.A., Skryabin, N.G., Tupitsyn, I.I. & Shinkarenko, A.V. 1991 [Sketch of wader's fauna of the Selenga River delta]. In: *Ecology and fauna of birds of Eastern Siberia*. Ulan-Ude. p. 93-100. In Russian.
- Zielinski, M. 1992 Egyptian Plover in Poland in October-November 1991. *Dutch Birding* 14(3): 99. (Ornithol. Station, Nadwislanska 108, 80-680 Gdańsk 40, Poland). *Pluvianus aegyptius*.
- MISCELLANEOUS
- Barbosa, A. 1990 Identification key of Iberian waders (*Charadriiformes*) based on the os quadratum. *Misc. Zool.* 14: 181-185. (UEI Ecología Evolutiva, Mus. Nacional de Ciencias Naturales CSIC, C/ José Gutiérrez Abascal 2, 28006 Madrid, Spain). Skulls of 28 wader species were analysed to elaborate an identification key based on the os quadratum. The key identifies 75% of the species concerned at specific level.
- Barbosa, A. 1991 European waders identification key on the bases of the cranial morphology. *Ardeola* 38(2): 249-263. With English and Spanish summaries. Skulls of 32 wader species were analysed to elaborate an identification key based on the cranial and postcranial morphometry.
- Beintema, A. 1992 Mayfield moet : oefeningen in het berekenen van uitkomstsucces. (Mayfield, a must : exercises in calculation of nesting success). *Limosa* 65(4): 155-162. In Dutch with English summary. (DLO-Inst. Bos-en Natuuronderzoek, P.O.Box 9201, 6800 HB Arnhem, Netherlands).
- Boschert, M. 1992 Rückstünde chlororganischer Verbindungen in Gelegen des Grossen Brachvogels (*Numenius arquata*) aus einem Brutgebiet am südlichen Oberrhein. (Organochloro-residues in clutches of the Curlew (*Numenius arquata*) from a breeding area in the southern Rhine Valley). *Ecol. Birds* 14: 101-111. In German with English summary. (Wiedigstr. 18, D-7580 Bühl, Germany). "DDT, PCB and HCB were detected in all samples, sometimes in relatively high concentrations. Aldrin, dieldrin and heptachlor were also found but less frequently.
- Ching, H.L. 1990 Some helminth parasites of Dunlin (*Calidris alpina*) and Western Willet (*Catoptrophorus semipalmatus inornatus*) from California. *J. Helminthol. Soc. Wash.* 57(1): 44-50. (Hydra Enterprises Ltd., P.O. Box 2184, Vancouver, BC, Canada V6B 3V7)
- Christian, P.D., Christidis, L. & Schodde, R. 1992 Biochemical systematics of the *Charadriiformes* (shorebirds) : relationships between the *Charadrii*, *Scolopaci* and *Lari*. *Aust. J. Zool.* 40(3): 291-302. (CSIRO, Div. Entomol., P.O. Box 1700, Lyneham, ACT 2601, Australia)
- Christian, P.D., Christidis, L. & Schodde, R. 1992 Biochemical systematics of the Australian Dotterels and Plovers (*Charadriiformes: Charadriidae*). *Aust. J. Zool.* 40(2): 225-233.
- Deblinger, R.D., Vaske, J.J. & Rimmer, D.W. 1992 An evaluation of different predator exclosures used to protect Atlantic coast Piping Plover nests. *Wildl. Soc. Bull.* 20(3): 274-279. (The Trustees of Reservations, 572 Essex St., Beverly, MA 01915, USA). *Charadrius melanotos*.
- Denny, M. 1992 Black Turnstone seen eating millet. *Oreg. Birds* 18(3): 74. (323 Scenic View Dr., College Place, WA 99324, USA). *Arenaria melanocephala*.
- Dickin, G. 1992 A highveld episode. *Birding*



in South Africa 44(2): 34-35. (Box 826, Delmas 2210, South Africa). Among others, *Glaureola nordmanni* as crop follower.

Exo, K.-M. & Scheiffarth, G. 1990 Are motion-sensitive transmitters suitable to record activity patterns and time-budgets of waders?. In: *Proceedings of the European Telemetry Conference Garmisch-Partenkirchen, Germany, May 15 to 17, 1990*. p. 401-403. With German summary. (Inst. Vogelforschung, An der Vogelwarte 21, D-2940 Wilhelmshaven). *Haematopus ostralegus*.

Exo, K.-M., Eggers, U., Laschinski-Sievers, R. & Scheiffarth, G. 1992 Monitoring activity patterns using a microcomputer-controlled radiotelemetry system tested for waders (*Charadrii*) as an example. In: Priede, I.G. & S.M. Swift (eds.). *Wildlife telemetry, remote monitoring and tracking of animals*. Chichester : Ellis Horwood, p. 79-87. Data obtained from *Haematopus ostralegus* showed that it is possible to record the overall locomotor activity as well as foraging behaviour by the position of the mercury switch of the system, as long as the transmitters are calibrated by visual observations. After a transmitter had been calibrated the system proved highly reliable for continuous long-term monitoring.

Gretton, A. 1991 *Conservation of the Slender-billed Curlew*. Cambridge : ICBP, 159p. ICBP Monograph No. 6. Monograph on *Numenius tenuirostris*, all records of the Western Palearctic are summarized and important resting areas are described.

Hill, D. & Player, A. 1992 Behavioural responses of Black-headed Gulls and Avocets to two methods of control of gull productivity. *Bird Study* 39(1): 34-42. (British Trust for Ornithol., The Nunnery, Nunnery Place, Thetford, Norfolk IP24 2PU, UK). Behaviour of *Larus ridibundus* and *Recurvirostra avosetta* was analysed in order to compare the effects of two methods of Gull productivity control, 1. formalin injection of eggs and 2. nest raking.

Johnson, A. & Pineau, O. 1992 Longévité record pour un Gravelot ° collier interrompu *Charadrius alexandrinus*. (Longevity record for the Kentish Plover *Charadrius alexandrinus*). *Alauda* 60(2): 111-112. In French. (Str. Biol. de la Tour du Valat, Le Sambuc, 13200 Arles, France)

Klomp, N.I. & Furness, R.W. 1992 A technique which may allow accurate determination of the age of adult birds. *Ibis* 134(3): 245-249. (School. Appl. Sci., Charles Stuart Univ.-Murray, P.O. Box 789, Albury, NSW 2640, Australia). A.o. *Tringa totanus*.

Schneider, D.C. 1992 Thinning and clearing of prey by predators. *Am. Nat.* 139(1): 148-160. (Ocean Sci. Ctr., Memorial Univ. New Foundland, St. John's A1B 3X7, Canada).

The purpose of the article is to present a general model for calculating the direction and magnitude of change in patchiness of immobile prey after a period of predation by *Limnodromus griseus*.

ABU Soest 1992 Bedeutung von Feuchtwiesen als Rastgebiet für Watvögel. (Significance of wet meadows as stop-over areas for migrating waders). *Vogelwelt* 113(3): 122-133. In German with English summary. (Arbeitsgemeinschaft Biol. Umweltschutz im Kreis Soest, Teichstr. 19, 4772 Bad Sassendorf-Lohne, Germany). *Vanellus vanellus*, *Pluvialis apricaria*, *Gallinago gallinago*, *Philomachus pugnax*, *Tringa glareola*.

Székely, T. & Bamberger, Z. 1992 Predation of waders (*Charadrii*) on prey population : an enclosure experiment. *J. Anim. Ecol.* 61(2): 447-456. (Dept. Zool., Kossuth Univ., Debrecen, H-4010, Hungary)

Yésou, P. 1992 Importance de la Baie de l'Aiguillon et de la Pointe d'Arcay (Vendée, France) pour les limicoles. (The importance of Baie de l'Aiguillon and Pointe d'Arcay (Vendée, France) for the waders). *Oiseau Rev. Française Ornithol.* 62(3): 213-233. In French with English summary. (O.N.C. Chanteloup, F-85340 L'Île d'Olonne, France)

Ziewitz, J.W., Sidle, J.G. & Dinan, J.J. 1992 Habitat conservation for nesting Least Terns and Piping Plovers on the Platte River, Nebraska. *Prairie Nat.* 24(1): 1-20. (J.G. Sidle: Platte River Whooping Crane Habitat Maintenance Trust, 2550 N. Diers Ave., Suite H, Grand Island, NE 68803, USA). *Charadrius melanotos*.

