

Recent Publications on Waders 58

COMPILED BY HENK KOFFIJBERG

In trying to make these lists as complete as possible, and the next one even more complete and up to date, your help is most welcome. Please send new titles, abstracts, reprints, copies, reports, and (important too!) omissions or corrections to: **Henk Koffijberg, Glenn Millerweg 102, 1311 RW Almere, The Netherlands.**

Remember to include, if possible, the address of the first author, and translations of titles if they are in other languages than English.

For help with this issue I am very grateful to D. Blanco, T. Piersma, K. van Scharenburg, I. Bainbridge, M. Stock, B. Trollet, P. Yesou and M.B. Pedersen

Translated titles from languages other than English are bracketed; with round brackets if the translation is by the author(s), square brackets if not.

N.B. a copy of your paper or abstract would be very helpful to enable me to add keywords or give added annotation to the titles.

BREEDING

Berg, Å., Lindberg, T. & Källebrink, K.G. 1992. Hatching success of Lapwings on farmland: differences between habitats and colonies of different sizes. *J. Anim. Ecol.* 61(2): 469-476. (Dept. Wildl. Ecol., Swedish Univ. Agric. Sci., Box 7002, S-750 07 Uppsala, Sweden).

Ens, B.J., Kersten, M., Breninkmeijer, A. & Hulscher, J.B. 1992. Territory quality, parental effort and reproductive success of Oystercatchers (*Haematopus ostralegus*). *J. Anim. Ecol.* 61(3): 703-715. (DLO Inst For. & Nat. Res., P.O. Box 167, 1790 AD Den Burg, Texel, Netherlands).

Flemming, S.P., Chiasson, R.D. & Austin-Smith, P.J. 1992. Piping Plover nest selection in New Brunswick and Nova Scotia. *J. Wildl. Manage.* 56(3): 578-583. *Charadrius melodus*.

Groen, N.M. 1993. De broedbiologie van de Grutto in een Noordhollands weidevogelreservaat. [Breeding biology of the Black-tailed Godwit in a meadowbird reserve in the province of North Holland]. *Graspieper* 13(1): 13-19. In Dutch. (Paardenkamp 49, 1541 XN Koog aan de Zaan, Netherlands). *Limosa limosa* in the province of North Holland, Netherlands.

Hut, R.M.G. van der 1993. Grutto's in de Zaanstreek

: ontwikkeling van de broedvogelstand in de jaren zeventig en tachtig. [Black-tailed Godwits in the Zaanstreek: breeding population in the seventies and eighties]. *Graspieper* 13(1): 20-28. In Dutch. (Geersbroekstr. 13, 1507 KR Zaandam, Netherlands). *Limosa limosa* in the province of North Holland, Netherlands.

Koiker, G. 1993. Phänologie und Brutbiologie des Kiebitzes (*Vanellus vanellus*): 17 jährige Beobachtungen in Nordwestdeutschland. (Phenology and breeding biology of Lapwing (*Vanellus vanellus*): results of a 17 years' study in NW-Germany). *J. Orn.* 134(1): 43-57. In German with English summary. (Univ. Osnabrück, FB Biologie/Chemie, Barbarastr. 11, 4500 Osnabrück, Germany).

L'Hyver, M.-A. & Miller, E.H. 1991. Geographic and local variation in nesting phenology and clutch size of the Black Oystercatcher. *Condor* 93(4): 892-903. (Biol. Dept., Univ. Victoria, P.O. Box 1700, Victoria, BC, Canada V8W 2Y2). *Haematopus bachmani*.

Newman, M. 1992. Pied Oystercatcher breeding at Mortimer Bay, Tasmania. *Emu* 92(2): 87-92. (Pascimco Metals-EZ, GPO Box 377D, Hobart, Tas., 7001, Australia). *Haematopus longirostris*.

Newman, O.M.G. & Park, P. 1992. Pied Oystercatcher breeding success in the 1991/92 summer. *Tasm. Bird Rep.* No. 21, pp. 13-22. (98 Nowra Rd., Lauderdale, Tasm. 7021, Australia). *Haematopus longirostris* in Tasmania.

Scharringa, C.J.G. 1993. De Grutto in Noord-Holland in 1979-1992. [Black-tailed Godwits in the province of North Holland]. *Graspieper* 13(1): 29-36. In Dutch. (Provincie Noord-Holland, Dienst Ruimte en Groen, Afd. Onderzoek en Informatie, P.O. Box 6090, 2001 HB Haarlem). Breeding population of *Limosa limosa* in the province of North Holland, Netherlands.

Schotsman, N. 1992. De Grutto in Friesland, 1970-1990. [The Black-tailed Godwit in the province of Friesland, 1970-1990]. *Vanellus* 45: 2-12. In Dutch. Breeding population of *Limosa limosa* in the province of Friesland, Netherlands.

Trollet, B., Levesque, A. & Ibanez, F. 1992. Repartition, chronologie et devenir des pontes chez le vanneau: exemple du marais breton en 1992. [Distribution, chronology and fate of the clutches of Lapwing: an example of the Marais breton]. B.M. O.N.C. No. 173: 2-7. In French. (O.N.C., Chanteloup, 85340 L'Île D'Olonne, France). *Vanellus vanellus*, Vendee, France.

Vander Haegen, W.M. 1992. *Bioenergetics of American Woodcock during the breeding season on Moosehorn National Wildlife Refuge, Maine*. Ph.D. dissertation, Univ. Maine (Orono), 116p. From Diss. Abstr. Int. B Sci. Eng. 53(5):2120-2121. *Scolopax minor*.

Vermeer, K., Morgan, K.H. & Smith, G.E.J. 1992. Black Oystercatcher habitat selection, reproductive success, and their relationship with Glaucous-winged Gulls. *Colon. Waterbirds* 15(1): 14-23. *Haematopus bachmani*.

BEHAVIOUR, ECOLOGY & EVOLUTION

Baccetti, N., Magnani, A. & Serra, L. 1991. Migration pattern and fattening of Curlew Sandpiper at an Italian stopover site. *Ist. Nazl. Biol. Selvaggina* No. 19: 3p. In English with Italian summary. (Ist. Nazl. Biol. della Selvaggina "A Ghigi", via Ca Fornacetti 9, 40064 Ozzano Emilia, Bologna, Italy). *Calidris ferruginea*.

Barter, M. 1992. Morphometrics and moult of Little Curlew *Numenius minutus* in north western Australia. *Stilt* 21: 20-21. (21 Chivalry Ave., Glen Waverley, VIC 3150, Australia).

Barter, M. 1992. Morphometrics and moult of Victorian Masked Lapwings *Vanellus miles novaehollandiae*. *Stilt* 21: 24-25.

Beintema, A.J. 1992. [On population dynamics of upland sandpipers: a model of probable survival]. *LOLF-Mitt.* 17(3): 26-28. In German. (DLO-Inst. Bos- en Natuuronderzoek, P.O. Box 9201, 6800 HB Arnhem, Netherlands).

Berg, Å. 1992. Habitat selection by breeding Curlews *Numenius arquata* on mosaic farmland. *Ibis* 134(4): 355-360. (Dept. Wildl. Ecol., Swedish Univ. Agric. Sci., Box 7002, S-750 07 Uppsala, Sweden).

Blanco, D., Goñi, H.R. & Pugnali, G. 1992. La importancia de Punta Rasa, Buenos Aires, en la migración del Chorlo Rojizo, (*Calidris canutus*). (Importance of Punta Rasa (Buenos Aires province) for the migration of the Red Knot). *Hornero* 13(3): 203-206. In Spanish with English summary. (Grupo Argentino de Limícolas, Monroe 2124, (1428) Buenos Aires, Argentina). The south end of the Samborombon Bay is considered as an important site in the Red Knot migration. Seasonal occurrence of Knots is analysed in order to evaluate the importance of the area as a stopover site. Results show that it is an important stopover site during April but not during September-October. It seems that Knots use different migration routes in northward and southward migration.

Brinkkemper, J. 1993. Grutto's kijken. [Watching Black-tailed Godwits]. *Graspieper* 13(1): 5-12. In Dutch. (Oranjeboomstraat 27, 1551 BH Westzaan, Netherlands). *Limosa limosa* in the province of North Holland, Netherlands. Observations and pictures of territorial and breeding behaviour.

Clark, N. 1993. Wash Oystercatchers starving. *BTO News* 185: 1, 24. *Haematopus ostralegus*, unusually large numbers found dead in the Wash (UK), due to small numbers of cockles and mussels in the autumn of 1992.

Dittberner, H. & Dittberner, W. 1991. [Epimeletic behaviour in *Vanellus vanellus*]. *Beitr. Vogelkd.* 37(5/6): 346-347. In German. (Rotenseestr. 6, O-2330 Bergen, Germany).

Fabritius, H.E. 1993. Slaapplaatsen van de Grutto tijdens de voorjaarstrek in de Noordhollandse veenweiden ten noorden van het Noordzeekanaal: een nieuw verschijnsel. [Communal roosting of the Black-tailed Godwit during spring migration in the province of North Holland: a new phenomenon]. *Graspieper* 13(1): 58-63. In Dutch. (Oost Mijzen 6,



1633 WB Avenhorn, Netherlands). *Limosa limosa* in the Netherlands. Numbers and distribution of communal roosting sites in the province of North Holland during spring migration. Peak numbers increased from 10,000 in March 1990 to 45,000 in March 1992.

Gerritsen, T. 1992. Oecomorfologie : tastzin en smaakzin in strandlopers. [Ecomorphology: tactile sense and taste in sandpipers]. *Limosa* 65(3): 128-129. In Dutch. *Calidris alpina*, *C. alba*, *C. maritima*, and *C. canutus*.

Gudmundsson, G.A. & Lindström, Å. 1992. Spring migration of Sanderlings *Calidris alba* through SW Iceland: wherefrom and whereto?. *Ardea* 80(3): 315-326. (Dept. Ecol., Univ. Lund, Ecol. Building, S-223 62, Lund, Sweden). Ringing recoveries and controls, together with a large number of sightings of colour-ringed birds revealed that many Sanderlings staging in Iceland in spring spend winter on the British Isles. Departure directions suggest breeding grounds in northeast Greenland.

Hampshire, J.S. & Russell, F.J. 1993. Oystercatchers rearing Northern Lapwing chick. *Brit. Birds* 86(1): 17-19. (Rose Cottage, Beach Road, Lessingham, Norfolk NR12 0SU, UK). *Haematopus ostralegus* with *Vanellus vanellus* chick.

Holmgren, N., Ellegren, H. & Pettersson, J. 1993. The adaptation of moult pattern in migratory Dunlins *Calidris alpina*. *Ornis Scand.* 24(1): 21-27. (Dept. Ecol., Theoretical Ecol., Lund Univ., Ecol. Bldg., S-223 62 Lund, Sweden). Post-nuptial primary moult of adult Dunlins was investigated along the Baltic coast at Ottenby and Falsterbo, S. Sweden. Secondly, the influence of a decreased wing area due to moult on migration energy reserves was studied.

Hulscher, J. 1992. Is de snavel van de mannelijke Scholekster een beter instrument om mossels te openen dan die van het vrouwtje? [Is the bill of the male Oystercatcher a better instrument for opening mussels than that of the female?]. *Limosa* 65(3): 129-130. In Dutch. (Zool. Lab., Univ. Groningen, P.O. Box 14, 9750 AA Haren, Netherlands). *Haematopus ostralegus*.

Jessop, R. 1992. Biometrics of Sanderling *Calidris alba* caught in Victoria, Australia. *Stilt* 21: 22-23. (14 Myrtle Grove, Altona, VIC 3018, Australia). Bill length, wing length and weights were similar to those reported for birds breeding near Spitsbergen [and]¹ Siberia.

Kaletja, B. 1992. Time budgets and predatory impact of waders at the Berg River estuary, South Africa. *Ardea* 80(3): 327-342. (Percy FitzPatrick Inst. African Ornithol., Univ. Cape Town, Rondebosch 7700, South Africa).

Kramer, D. & Palmer, J.C. 1993. Grey Phalarope feeding in association with ducks. *Brit. Birds* 86(1): 19. (7 Little Headlands, Putnoe, Bedford MK41 8JT,

UK). *Phalaropus fulicarius*.

Larsen, T. & Moldsvor, J. 1992. Antipredator behavior and breeding associations of Bar-tailed Godwits and Whimbrels. *Auk* 109(3): 601-608. (Dept. Anim. Ecol., Zool. Mus., Univ. Bergen, N-5007 Bergen, Norway). *Limosa lapponica* and *Numenius phaeopus*.

Mouritsen, K.N. & Jensen, K.T. 1992. Choice of microhabitat in tactile foraging Dunlins *Calidris alpina*: the importance of sediment penetrability. *Mar. Ecol. Prog. Ser.* 85(1/2): 1-8. (Inst. Biol. Sci., Dept. Ecol. & Genet., Univ. Aarhus, Ny Munkegade, DK-8000 Aarhus C, Denmark).

Mundkur, T. 1991. Prey items of the Great Thick-knee *Esacus recurvirostris*. *Forktail* 6: 71-72. (Asian Wetland Bureau, Inst. Advanced Studies, Univ. Malaya, Lembah Pantai, 59100 Kuala Lumpur, Malaysia).

Newman, O.M.G. 1992. Nesting association between Hooded Plover and Pied Oystercatcher. *Stilt* 21: 26. (98 Nowra Rd., Lauderdale, Tasmania, 7021, Australia). *Haematopus longirostris* and *Charadrius rubricollis*.

Oring, L.W., Fleischer, R.C., Reed, J.M. & Marsden, K.E. 1992. Cuckoldry through stored sperm in the sequentially polyandrous Spotted Sandpiper. *Nature* (Lond.) 259(6396): 631-633. (Ecol., Evol. & Conserv. Biol. Program, Univ. Nevada, Reno 89512, USA). *Tringa macularia*. See review article this *Bulletin*.

Piersma, T. & Sant, S. van de 1992. Pattern and predictability of potential wind assistance for waders and geese migrating from West Africa and the Wadden Sea to Siberia. *Ornis Svecica* 2(2): 55-66. (NIOZ, P.O. Box 59, 1790 AB Den Burg, Texel, Netherlands).

Piersma, T. & Jukema, J. 1993. Red breasts as honest signals of migratory quality in a long-distance migrant, the Bar-tailed Godwit. *Condor* 95: 163-177. Describes in detail the pre-nuptial moult of *Limosa lapponica* and analyzes it in relation to plumage colour, nutritional status, and migration schedule. Analysis suggests links between moulting during long-distance migration and breeding performance.

Serasinghe, R. 1992. Communal roosting of the Common Sandpiper (*Actitis hypoleucos*). *Loris* 19(5): 174-175.

Sidle, J.G., Carlson, D.E., Kirsch, E.M. & Dinan, J.J. 1992. Flooding mortality and habitat renewal for Least Terns and Piping Plovers. *Colon. Waterbirds* 15(1): 132-136. *Charadrius melodus*.

Summers, R.W., Underhill, L.G., Nicoll, M., Rae, R. & Piersma, T. 1992. Seasonal, size- and age related patterns in body-mass and composition of Purple Sandpipers *Calidris maritima*. *Ibis* 134(4): 346-354. (Lismore Mill Crescent, North Kessock, Easter-Ross, Scotland).

Thompson, P.S. & W.G. Hale 1993. Adult survival and numbers in a coastal breeding population of Redshank *Tringa totanus* in northwest England. *Ibis*

135(1): 61-69. (Univ. Durham, Dept. Biol. Sci., Sci. Lab., South Rd., Durham DH1 3LE, England).

Tomkovich, P.S. 1992. Migration of the Spoon-billed Sandpiper *Eurynorhynchus pygmeus* in the Far East of the Russian Federation. *Stilt* 21: 29-33. (Zool. Mus., Moscow Univ., Herzen St. 6, Moscow 103009, Russia).

Tüllinghoff, R. & Bergmann, H.-H. 1993. Zur Habitatnutzung des Grossbrachvogels (*Numenius arquata*) im westlichen Niedersachsen: Bevorzugte und gemiedene Elemente der Kulturlandschaft. (Curlew habitats (*Numenius arquata*) in Lower Saxonia - preferred and avoided structures of agricultural landscapes). *Vogelwarte* 37(1): 1-11. In German with English summary. (Gartenstr. 4, D 4553 Voltlage, Germany).

Weggler, M. 1992. Nahrungssuchverhalten von vier Limikolenarten: Fortbewegungsmuster, Sucherfolg und Raumnutzung. (Foraging behaviour of four species of waders: patterns of locomotion, foraging success and spatial distribution). *Orn. Beob.* 89(3): 177-190. In German with English summary.

Wenink, P.W., Baker, A.J. & Tilanus, M.G.J. 1993. Hypervariable- control-region sequences reveal global population structuring in a long-distance migrant shorebird, the Dunlin (*Calidris alpina*). *Proc. Natl. Acad. Sci. USA* 90(1): 94-98. (A.J. Baker: Dept. Ornithol., Royal Ontario Mus., Toronto, ON, Canada M55 2C6).

Weston, M. & Rush, M. 1992. Probable pseudo copulation of a Red-necked Avocet. *Stilt* 21: 28. (28 Craig Rd., Donvale, Vic. 3111, Australia). *Recurvirostra novaehollandiae*.

Zomerdijk, P. & Groen, N. 1993. De Grutto buiten de broedtijd. [The Black-tailed Godwit outside the breeding season]. *Graspieper* 13(1): 46-52. In Dutch. (Kalmanstr. 21, 1817 HV Alkmaar, Netherlands). *Limosa limosa*, migration routes and strategies, and stop-over sites in Europe and Africa.

Zwarts, L. 1993. Het voedsel van de Grutto. [Food of the Black-tailed Godwit]. *Graspieper* 13(1): 53-57. In Dutch. (Rijkswaterstaat- Flevoland, P.O. Box 600, 8200 AP Lelystad, Netherlands). *Limosa limosa*, prey choice during different stages of autumn migration at several stop-over sites in Europe and in Africa.

DISTRIBUTION & FAUNISTICS

Appert, O. 1992. Der Flussuferläufer *Actitis hypoleucos* Brutvogel im Kanton Luzern. (The Common Sandpiper *Actitis hypoleucos* as a breeding bird in the canton Lucerne). *Orn. Beob.* 89(3): 196-197. In German with English summary.

Barter, M. 1992. Changing wader numbers in Swan Bay, Victoria: a cause for concern? *Stilt* 21: 8-12. (21 Chivalry Ave., Glen Waverley, VIC 3150, Australia). Decline in waders from 12,000 to 7,000 over the period 1983-1992. Fall has mainly occurred with smaller waders as *Calidris ferruginea* and



Calidris ruficollis.

Beusekom, R. van 1993. IJslandse Grutto's in Noord-Holland: herkenning en voorkomen. [Icelandic Black-tailed Godwits in the province of North Holland: recognition and occurrence]. *Graspijper* 13(1): 64-72. In Dutch. (Dienst Ruimte en Groen, P.O. Box 6090, 2001 HB Haarlem). *Limosa limosa islandica* in the Netherlands.

Blasco, A. 1992. [Probable over-wintering of Red-necked Phalarope *Phalaropus lobatus* in the Camargue (southern France) during the 1990-91 winter]. *Alauda* 60(3): 174-175. In French. (Le clovert, chemin du Chaland-Les Carabins, 13270 Fos-sur-Mer, France).

Boldt, B. 1992. Black-necked Stilt (*Himantopus mexicanus*) - 10 October 1991, Milwaukee County, at the Milwaukee coast guard impoundment. *Passenger Pigeon* 54(2): 174-175. (1832 Jeffery Ln., Waukesha, WI 53186, USA).

Burcar, K. 1992. Black-necked Stilt (*Himantopus mexicanus*) - 11 October 1991, Milwaukee County, at the Milwaukee coast guard impoundment. *Passenger Pigeon* 54(2): 175-176. (5136 Enchanted Valley Rd., Cross Plains, WI 53528, USA).

Burcar, K. 1992. Purple Sandpiper (*Calidris maritima*) - 4 November 1991, Sheboygan County, Northpoint Park in Sheboygan. *Passenger Pigeon* 54(2): 176.

Campbell, J. 1992. Anderson Inlet wader & waterbird survey report No. 1 1991/92. *Stilt* 21: 13-16. (8/5 Wattle Ave., Glen Huntly, VIC 3163, Australia).

Carlson, S. 1992. Female Ruff in Laq Qui Parle County. *Loon* 64(3): 169. (2705 Dupont Ave., S., Minneapolis, MN 55408, USA). *Philomachus pugnax* in Minnesota.

Clements, J.F. 1992. Sight record of Lesser Yellowlegs in French Polynesia. *Elepaio* 52(11): 81. (3420 Fredas Hill Rd, Vista, CA 92804, USA). *Tringa flavipes*.

Cowart, B. 1992. Red Phalarope (*Phalaropus fulicaria*) - 4 November 1991, Racine County, at Wind Point. *Passenger Pigeon* 54(2): 177. (3518 N. Murray, Milwaukee, WI 53211, USA).

Domagalski, R.C. 1992. Black-necked Stilt (*Himantopus mexicanus*) - 10 October 1991, Milwaukee County, at the Milwaukee coast guard impoundment. *Passenger Pigeon* 54(2): 175. (W140 N8508 Lilly Rd., Menomonee Falls, WI 53051, USA).

Dufourny, H. & Lebailly, J.L. 1991. Premiere observation d'un Becasseau de Bonaparte (*Calidris fuscicollis*) en Belgique. [First observation of White-rumped Sandpiper (*Calidris fuscicollis*) in Belgium]. *Aves* 28(4): 220-222. In French. (Rue du Rambaix 20, B-7387 Angre, Belgium).

Frank, J.C. 1992. Black-necked Stilt (*Himantopus mexicanus*) - 10 October 1991, Milwaukee County, at the Milwaukee coast guard impoundment.

Passenger Pigeon 54(2): 175. (4339 W. Laverna Ave., Mequon, WI 53092, USA).

Hall, G.A. 1992. Lesser Golden Plover on Allegheny Front Mountain, West Virginia. *Redstart* 59(2): 46-47. (Rt. 12, Box 89, Morgantown, WV 26505, USA). *Pluvialis dominica*.

Hirschfeld, E. 1991. First record of Long-toed Stint *Calidris subminuta* in Bahrain. *Sandgrouse* 13(2): 108-109.

Hutton, K. 1992. Long-toed Stints in New South Wales. *Aust. Birds* 26(1): 20-28. (9 Karri Rd., Leeton, NSW 2705, Australia). *Calidris subminuta*.

Kipp, M. 1992. [The status of the Curlew in Northrhine-Westphalia]. *LOLF-Mitt.* 17(3): 28-32. In German. (Kuhdamm 95, 4540 Lengerich, Germany). *Numenius arquata*.

Kirby, J.S. & Lack, P.C. 1993. Spatial dynamics of wintering Lapwings and Golden Plovers in Britain and Ireland, 1981/82 to 1983/84. *Bird Study* 40(1): 38-50. (Wildfowl & Wetlands Trust, Slimbridge, Gloucester GL2 7BT, UK). Distribution of *Vanellus vanellus* and *Pluvialis apricaria* has been analysed using data from The Atlas of Wintering Birds in Britain and Ireland, and the Birds of Estuaries Enquiry. In all three winters, Lapwings were more abundant in Britain than in Ireland, but Golden Plovers were more abundant in Ireland. The possible reasons for this are discussed.

Korducki, M. 1992. Purple Sandpiper (*Calidris maritima*) - 29 November 1991, Racine County, at Myers Park. *Passenger Pigeon* 54(2): 176-177. (4410 S. 21st. St., Milwaukee, WI 53221, USA).

Krych, S. 1992. Minnesota's third record of Black-necked Stilt. *Loon* 64(3): 166-167. (1045 Hiawatha Ave., #310, Hopkins, MN 55343, USA). *Himantopus mexicanus*.

Lehmann, W. 1992. Austernfischerbrut bei Torgau. [Oystercatcher breeding at Torgau]. *Falke* 39(9): 308-309. In German. *Haematopus ostralegus* in Germany.

Lethaby, N. & Gilligan, J. 1992. Great Knot in Oregon. *Am. Birds* 46(1): 46-47. *Calidris tenuirostris*.

McLaren, I.A. & Maybank, B. 1992. Apparent Broad-billed Sandpiper in Nova Scotia. *Am. Birds* 46(1): 48-50. *Limicola falcinellus*.

McLaughlin, M. 1992. Almost stilted over a reclusive Stilt. *Pa. Birds* 6(2): 66. *Himantopus mexicanus* in Pennsylvania.

Miller, J.C. 1988/1989. First nesting of a Black-necked Stilt in Pennsylvania. *Cassinia* No. 63: pp. 73. (1220 Prospect Ave., Prospect Park, PA 19076, USA). *Himantopus mexicanus*.

Murray, R. & McDavit, M.W. 1993. First nesting records of Piping Plover in South Carolina. *Chat* 57(1): 10-11. (4387 8th Ave., Little River, SC 29566, USA). *Charadrius melodus*.

Naismith, L. 1992. Results of the 1991. summer population monitoring count. *Stilt* 21: 6-7. (40 Wattle Valley Rd., Mitcham 3132, Australia)

Neubeck, P. 1992. Ruff in Carver County. *Loon* 64(3): 173-174. (1504 S. Tyrol Trail, Golden Valley, MN 55416, USA). *Philomachus pugnax* in Minnesota.

Nikolaus, G. 1992. Some observations on Dunlin *Calidris alpina* wintering in the Sudan. *Scopus* 15(2): 124-126.

O'Brien, M. 1993. The survey of breeding waders on Scottish agricultural land. *BTO News* 184: 6-7. (BTO, The Nunnery, Nunnery Place, Thetford, Norfolk IP24 2PU, UK). First results of the survey on non-upland areas of Scotland for *Haematopus ostralegus*, *Vanellus vanellus*, *Gallinago gallinago*, *Numenius arquata*, *Tringa totanus*.

Olioso, G. & le Comite d'Homologation National 1992. [The Marsh Sandpiper *Tringa stagnatilis* in France]. *Alauda* 60(3): 143-147. In French with English summary. (Cent. Reg. Bague Provence, Le Mail, 26320 Grignan, France).

Ouwerkerk, R. van 1992. Grey Phalarope *Phalaropus fulicarius* in northern Cameroon. *Malimbus* 14(1): 25.

Pearson, D.J., Coverdale, M.A.C. & Archer, A.L. 1992. Recent surveys of wintering waders at Lake Turkana, Kenya with details of 1989 counts on the northeast shore. *Scopus* 16(1): 29-33.

Pedersen, M.B. 1992. *Distribution, population size and ecology of the Jack Snipe in Southern Sweden*. Project report 1992. 4 pp. (Bygager 24, DK-7190 Billund, Denmark). WWF Project No. 326. *Lymnocyptes minimus*.

Pedersen, M.B. 1992. *Breeding status of the Jack Snipe in South Baltic: a preliminary study in Lithuania*. WWF Project No. 326 (Suppl. 1992.) 2 pp. *Lymnocyptes minimus* breeding data for Estonia, Latvia, Lithuania, Poland. No information from RSFSR Kaliningrad.

Pittaway, R. 1992. Short-billed Dowitcher subspecies. *Birding* 24(5): 309-311. (Box 619, Minden, ON, Canada K0M 2K0). *Limnodromus griseus* in Canada.

Roos, M.T. & Schrijvershof, P.G. 1993. Red-wattled Lapwing in Israel in winter of 1991/92. *Dutch Birding* 15(1): 13-15. In English with Dutch summary. (M.T. Roos: Zevenkampse Ring 742, 3068 HG Rotterdam, Netherlands). *Hoplopterus indicus*.

Safford, R.J. 1992. A record of Great Knot *Calidris tenuirostris* from Mauritius, Indian Ocean. *Bull. Brit. Orn. Club* 112(2): 134-135.

Sagar, P. 1992. New Zealand wader counts - 1990 & 1991. *Stilt* 21: 34-36. (38A Yardley St., Christchurch 4, New Zealand).

Scebba, S., Vitolo, A. & Moschetti, G. 1991. [First



confirmed summer nesting of *Himantopus himantopus* in Campania]. *Avocetta* 15(1-2): 67. In Italian. (Via Posillipo 276/2-800123 Napoli, Italy).

Schwoppe, W. & Schwoppe, M. 1992. [Status of breeding upland sandpipers in the Borken district]. *LOLF-Mitt.* 17(3): 32-35. In German. (Biol. Stn. Zwillbrock, Zwillbrock 10, 4426 Vreden, Germany).

Smith, F.T.H. 1992. Second Australian sighting of the Stilt Sandpiper *Micropalama himantopus*. *Aust. Bird Watcher* 14(8): 313-317. (71 Cobden St., Kew, Vic. 3101, Australia).

Spencer, R. and the Rare Breeding Birds Panel 1993. Rare breeding birds in the United Kingdom in 1990. *Brit. Birds* 86(2): 62-90. (Rare Breeding Birds Panel, c/o Fountains, Park Lane, Blunham, Bedford MK44 3NJ, UK). Breeding waders: *Recurvirostra avosetta*, *Burhinus oedicnemus*, *Charadrius morinellus*, *Calidris temminckii*, *Calidris maritima*, *Philomachus pugnax*, *Limosa limosa*, *Tringa glareola*, and *Phalaropus lobatus*.

Steva, S. 1992. Ruffs in Polk County. *Loon* 64(3): 167-169. (Rt. 2, Box 51, Thief River Falls, MN 56701, USA). *Philomachus pugnax* in Minnesota.

Tipper, R. 1993. Waders in Hong Kong. *Brit. Birds* 86(5): 231-242. (The Old Bakehouse, Hanging Birch Lane, Waldron, East Sussex TN21 0PA, UK). Waders at the Mai Po Nature Reserve. Colour photographs of *Limicola falcinellus*, *Calidris tenuirostris*, *Glareola pratincola*, *Calidris subminuta*, *Limnodromus semipalmatus*, *Tringa guttifer*.

Tubbs, C.R., Tubbs, J.M. & Kirby, J.S. 1992. Dunlin *Calidris alpina alpina* in The Solent, southern England. *Biol. Conserv.* 60(1): 15-30.

Uhlig, R. 1991. [New findings about *Arenaria interpres* in Bulgaria]. *Beitr. Vogelkd.* 37(5/6): 348. In German. (Hufelandstr. 10, -1055 Berlin, Germany).

Vieweg, A. 1992. Der Flussuferläufer, Brutvogel an der Zschopau. [Common Sandpiper, breeding bird of Zschopau]. *Falke* 39(9): 310. In German. *Actitis hypoleucos* in Germany.

Vijayagopal, K. & Chacko, S. 1991. Hitherto unrecorded nesting site of Yellow-wattled Lapwing *Vanellus malabaricus* (Boddaert). *J. Bombay Nat. Hist. Soc.* 88(3): 451.

MISCELLANEOUS

Brinkmann, R., Köhler, B., Heins, J.-U. & Rösler, S. 1991. *Menderes-Delta: Zustand und Gefährdung eines ostmediterranen Flussdeltas*. [Menderes-Delta: position and threats of an East Mediterranean river delta]. Arbeitsbericht des Fachbereichs Stadt- und Landschaftsplanung der Gesamthochschule Kassel, Heft 99, 240 pp. In German. Available from: Infosystem Planung, Gesamthochschule Kassel, Henschelstrasse 2, W-3500, Kassel, Germany. Costs: 20,- DM). With information on staging, wintering, and breeding waders in the Menderes Delta, SW Turkey.

Cadée, G.C. 1992. Raadsel L/R sortering bij door de wind getransporteerde *Mya*-schelpen opgelost. (Wind of left/right sorting in eolian transported *Mya*-shells solved). *Corresp.-blad Ned. Malac. Ver.* No. 272: 68-70. In Dutch with English summary. (Waterweg 12, 1791 LH Den Burg, Texel, Netherlands). *Arenaria interpres* causing saltational transportation of *Mya arenaria*.

Calvo, B. & Furness, R.W. 1992. A review of the use and the effects of marks and devices on birds. *Ring. & Migr.* 13(3): 129-151. (Applied Ornithology Unit, Dept. Zool., Univ. Glasgow, Glasgow G12 8QQ, UK). First, a quantitative review of marking methods and devices used in studies published in several leading bird journals is presented. Then the authors consider the extent to which the methods have been demonstrated to provide unbiased results, by reviewing research published throughout the scientific literature. Containing 374 references and appendix with references on effects of different marks and devices.

Fleet, D.M. 1993. Merlin stalking Dunlins on foot. *Brit. Birds* 86(4): 181. *Calidris alpina*.

Gewalt, W. 1992. Hohes Alter eines Schwarzen Austernfischers (*Haematopus ater*) im Zoo Duisburg. [A very old Blackish Oystercatcher in the zoo of Duisburg (Germany)]. *Beitr. Vogelkd.* 38(2): 138. In German.

Kölsch, G. 1992. Erfahrungen und Ergebnisse der Internationalen Wasservogel- und Limikolenzählungen in Deutschland. (Experiences and some results of the international waterfowl and wader census in Germany). *Vogelwelt* 113(4-5): 273-277. In German with English summary. (Biol. Stn. Münster, Coermühle 181, W-4400 Münster, Germany). Several aspects both of the International Waterfowl Census and the Inland Wader Counts are described: extent of census work, count sites, organisation and financial matters.

Lane, B.A. 1992. The impact of *Spartina* on international migratory waders. *Stilt* 21: 17-19. (Asian Wetland Bureau, Inst. Advanced Studies, Univ. Malaya, Lembah Pantai, 59100 Kuala Lumpur, Malaysia).

Mackereth, B. 1992. Avian pox in Variable Oystercatcher chick. *Notornis* 39(4): 248. (3 Harbour Lights Tce, R.D. 2, Whitanga, New Zealand). *Haematopus unicolor*, New Zealand.

Manakadan, R. 1991. Greenshank *Tringa nebularia* (Gunner) feeding on large fish. *J. Bombay Nat. Hist. Soc.* 88(3): 451-452.

Manakadan, R. 1991. A flock of one-legged Greenshanks *Tringa nebularia*. *J. Bombay Nat. Hist. Soc.* 88(3): 452.

Morton, J.M., Kirkpatrick, R.L. & Smith, E.P. 1991. Comments on estimating total body lipids from measures of lean mass. *Condor* 93(2): 463-465. (Dept. Fish. & Wildl. Sci., VPI&SU, Blacksburg, VA 24061, USA). *Scelopax minor*.

Pfister, C., Harrington, B.A. & Lavine, M. 1992. The impact of human disturbance on shorebirds at a migration staging area. *Biol. Conserv.* 60(2): 115-126.

Reed, J.M. & Oring, L.W. 1992. Reconnaissance for future breeding sites by Spotted Sandpipers. *Behav. Ecol.* 3(4): 310-317. (Program Ecol., Evol. & Conserv. Biol., RWF, 1000 Valley Rd., Univ. Nev., Reno 89512, USA). *Tringa macularia* in Minnesota, USA.

Scharenburg, K. van, Meijering, J. & Koning, R. de 1992. *Rapportage Biologisch Meetnet Provincie Groningen 1986-1990*. [Report of the Biological Monitoring Project in the province of Groningen 1986-1990]. Groningen: Provincie Groningen, Dienst RO, Afd. Groen, Bureau NLO, 73 p. In Dutch. With results of monitoring meadowbirds, 1986-1990, related to agricultural and nature management in the province of Groningen, Netherlands.

Schulz, M. 1992. Hooded Plover taken by a Peregrine Falcon. *Stilt* 21: 27. (167 South Beach Rd., Bittern, Vic. 3918, Australia). *Charadrius rubricollis*.

Stock, M. 1992. Ungestörte Natur oder Freizeitnutzung? : das Schicksal unserer Strände. [Unspoiled nature or leisure ground? : the faith of our beaches]. In: Prokosch, P. (ed.): *Ungestörte Natur - was haben wir davon?* WWF-Tagungsbericht 6, Husum, pp. 223-249. In German. (Landesamt für den Nationalpark Schleswig-Holsteinisches Wattenmeer, 2253 Tönning, Schlossgarten 1, Germany). Containing information on breeding numbers of *Charadrius alexandrinus* in the Wadden Sea area of Schleswig-Holstein, Germany, and the effects of human disturbance in breeding areas.

Tanger, D. 1993. De Grutto en het agrarisch beheer. [The Black-tailed Godwit and agricultural management]. *Graspieper* 13(1): 37-45. In Dutch. (Stichting Vrijwillig Natuur- en Landschapsbeheer Noord-Holland, P.O. Box 6090, 2001 HB Haarlem, Netherlands). The role of agriculture and meadowbird conservation is discussed for *Limosa limosa* in the province of North Holland, Netherlands.

Telford, S.R. Jr., Spalding, M.G. & Forrester, D.J. 1992. Hemoparasites of wading birds (*Ciconiiformes*) in Florida. *Can. J. Zool.* 70(7): 1397-1408. In English with French summary. (D.J. Forrester, Dept. Infectious Dis., Coll. Vet. Med., Univ. Florida, Gainesville 32610, USA).

Watson, A. 1993. Dotterels evading Peregrine Falcon by settling near human being. *Brit. Birds* 86(3): 131. (c/o Inst. Terrest. Ecol., Banchory, Kincardineshire AB31 4BY, UK). *Charadrius morinellus*.

Zingo, J.M. 1992. Common Tern adopts American Oystercatcher egg at Falkner Island. *Conn. Warbler* 12(4): 146-149. (Queens Coll. Cent., Environ. Teaching & Res., 31 Lloyd Harbor Rd., Huntington, NY 11743, USA). *Haematopus palliatus* and *Sterna hirundo*.

