

Brown Thrasher, Oven-bird, Mourning Dove, Crested Flycatcher, and Catbird Age Records.

Brown Thrasher Return-2 S (Summer) (<i>Toxostoma rufum</i>)		
Adult A322167, banded August 27, 1929.		Age
A Return-1 on May 8, 1930.		
A Return-2 on July 3, 1934.		Six years.
Over-bird Return-1 S (<i>Seiurus aurocapillus</i>)		
Adult A161565, banded on August 30, 1930.		
A Return-1 on May 20, 1934.		Five years.
Mourning Dove Return-1 S (<i>Zenaidura macroura carolinensis</i>)		
Adult A410063, banded May 8, 1930.		
A Return-1 on June 21, 1934.		Five years.
Northern Crested Flycatcher Return-2 S (<i>Myiarchus crinitus crinitus</i>)		
Adult A147214, banded June 30, 1929.		
A Return-1 on June 12, 1933, using same nest-box as in 1929.		
A Return-2 on July 8, 1934, using another nest-box.		Six years.
Catbird Return-4 S (<i>Dumetella carolinensis</i>)		
Immature A222905, banded July 5, 1929.		
Return-1 on May 18, 1930.		
Return-2 on September 29, 1931.		
Return-3 on May 12, 1932.		
Return-4 on July 3, 1933, found dead an eighth of a mile from traps.		Four years.

—RAYMOND J. MIDDLETON, Norristown, Pennsylvania.

A Coastal Group of Canada Geese.—In the course of trapping Black Ducks at the Austin Ornithological Research Station a few Canada Geese (*Branta c. candensis*) have been banded. These birds squeezed through the narrow funnel entrances of the heart-shaped duck traps. The first Goose was captured on January 20, 1932, and by April the total was 43.

Of these 43 Geese, 17 (39.5 per cent) have been reported as shot during the following periods: three in Newfoundland the spring subsequent to banding; twelve the next fall at scattered points along the coast from New Brunswick to New Jersey; one the second autumn in Florida; and the last bird in Yarmouth, Massachusetts, on November 23, 1934.

During the winter of 1932-33 only eighteen Geese were banded. Within a few days five were shot at a stand two miles to the south, one was found dead near here the following autumn, and one was shot in Old Lyme, Connecticut, in December, 1934. Of three birds banded in the spring of 1934, one was shot at Nantucket, Massachusetts, the following December and one retrapped here the following January.

The points of recovery are:

Newfoundland	465555	Terenceville.
	465583	Morrisville, Conne River.
	A717006	" " " "
Nova Scotia	465599	Long Island, Cape Sable Island.
Quebec	465600	Lobster Bay, Saguenay Bay.
New Brunswick	A717005	Inkerman.
	A717009	Lamegue.
Massachusetts	465568	Great Pond, Eastham, Cape Cod.
	465571	Chilmark, Martha's Vineyard.
	465573	Marshfield.
	465574	Mill Pond, Yarmouth, Cape Cod.
	465585	East Falmouth, Cape Cod.
	A717010	Barnstable, Cape Cod.
	A717103	Nantucket.
	A717026	Old Lime, Connecticut.
New York	465570	East Moriches, Long Island.
New Jersey	465556	Middle Island, Tuckerton Bay.
	465578	between Ocean City and Sea Isle City.
Florida	465576	St. George's Island near Apalachicola.

These recoveries, although few, are very interesting from the point of view of the source of the New England Geese. The data obtained from the Geese banded by Jack Miner at Kingsville, Ontario, show that his birds come from the whole