

## INDEX TO VOLUME 44

Compiled by BRIAN A. HARRINGTON  
AND THE  
MANOMET BIRD OBSERVATORY

This index is for new papers published in this volume of *Bird-Banding* as well as a subject and species index to papers reviewed in Volume 44 Recent Literature sections, and a title and author index to book and monograph reviews. Items indexed from Recent Literature are identified by (r), those from books and monographs by (rev.).

- Acanthis flammea*, sex determination, 13-21  
  *A. hornemannii*, sex determination, 13-21  
*Accipiter gentilis*, mercury contamination, 72(r)  
  *A. nisus*, migration and banding, 67(r)  
Activity patterns, see behavior  
*Aegolius acadicus*, sense of touch, 125  
AGE, CRITERIA  
  Grosbeak, Evening, 240(r)  
  Flycatcher, Western, 205-220  
  Jay, Beechey, 81-90  
  —, Bushy-crested, 81-90  
  —, San Blas, 81-90  
  —, Yucatan, 81-90  
  Mockingbird, 62, 124  
  Ovenbird, 239-40  
  Puffin, Common, 238(r)  
  Vireo, White-eyed, 63  
—, LIFE EXPECTANCY AND LONGEVITY  
  Birds, 66(r)  
  Bluebird, nestlings, 110-123  
  Bobolink, 57-58  
  Finch, Purple, 227  
  Hawk, Sparrow, 66(r)  
  Honeyeater, Fuscous, 127(r)  
  —, White-naped, 127(r)  
Longevity records, 319(r)  
Mortality and weather, 61-62, 67(r), 110-123  
Osprey, juveniles, 180-186  
Parulidae, 221-224  
Pelican, White, 66(r)  
Scaup, Greater, 127(r)  
Stilt, Black-winged, 67(r)  
Tern, Caspian, 127(r)  
  —, Common, 65  
Waterfowl, 68(r)
- , STUDIES OF KNOWN-AGE BIRDS  
  Bobolink, 57-58  
  Western Flycatcher, migration 205-220  
  Mockingbird, movements, 62  
*Agelaius phoeniceus*, populations and migration, 68(r)  
  —, recognition of eggs, young, and nests, 130(r)
- , changes of seasonal breeding behavior, 70(r)  
*Alca torda*, New England population changes, 267-313  
Alison, R. M. Delayed nesting in Old-squaws, 61-62  
Allen, R. T., and F. C. James (eds.). A symposium on ecosystematics (rev.), 135-136  
*Amblyornis*, distribution, taxonomy, 327(r)  
*Anas acuta*, distribution, 240(r)  
  *clypeata*, reproduction, 233(r)  
  *platyrhynchos*, gamma radiation of —, 73(r)  
  —, heat loss when oiled, 323(r)  
Anatidae, sound production, 239(r)  
  —, summer migrations, 128(r)  
  —, traps, 230(r)
- ANATOMY AND PHYSIOLOGY  
Air-sacs, *Anhinga* and *Phalacrocorax*, 236(r)  
Avian retina, ultrastructure, 238(r)  
Blood sugar, migrants and residents, 237(r)  
Extraocular muscles, 74(r)  
Eyes, UV light damage, 324-25(r)  
Growth and gamma radiation, 73(r)  
Hearing, neural centers, 74(r)  
Heart rate, 73(r)  
Heat loss, oiled ducks, 323(r)  
Migrants, body fat chemistry, 237(r)  
Mortality, gamma radiation, 73(r)  
Nest temperatures, 237(r)  
Neuro-hormonal heat regulation, 324(r)  
Pecten oculi, function, 324(r)  
Pigeon's milk, nutrition, 73(r)  
Rearing young, energy expended, 73(r)  
Syrinx, 239(r)  
Toe pads, 325(r)  
Tongue structure, 325(r)
- Anhinga*, see *Anhinga anhinga*  
*Anhinga anhinga*, air-sac system, 236(r)  
Antbird, Spotted, see *Hylophylax naevioides*  
*Apteryx australis*, chick biometrics, 325(r)  
*Apus apus*, migration and night flights, 319(r)

- A. pacificus*, food, 241(r)
  - Ardea herodias*, breeding biology, 320-21(r)
  - Athene noctua*, taxonomy, 326-27(r)
  - AVICULTURE**
    - Caique, Blue-headed, 322(r)
    - Aythya fuligula*, activity and feeding, 70(r)
    - A. marila*, heat loss when oiled, 323(r)
      - , longevity, 127(r)
  - Band loss, see banding
  - BANDING, BAND LOSS**
    - Fulmar, 127(r)
    - Gull, Red-billed, 67(r)
  - , COMPUTERIZED RECORDS, 187-195, 249-257
  - , RECOVERIES
    - Accipiter nisus*, 67(r)
    - Australian birds, 127(r)
    - Finch, Purple, 227
    - Pintail, 240(r)
    - Siskin, Pine, 227-228
    - Tern, Common, 127(r)
      - , Royal, 240-41(r)
  - , REPORTS
    - Barn Owls in Australia, 231(r)
    - Galapagos Islands, 230(r)
    - Gulls in Montana, 231(r)
    - South Africa, 230(r)
  - , RETURNS
    - Finch, Purple, 227
    - North American birds in Mexico, 228-229
    - North American Parulidae in Jamaica, 221-224
  - , STATIONS, 66(r)
  - , TECHNIQUES
    - Statistics, 66(r)
      - Tattooing nestlings, 63
    - Worksheets, 230(r)
  - , TRAPS AND NETS
    - For Anatidae
    - Cannon net, 66(r), 67(r)
    - Live trap, 66(r)
      - Nest box trap, 56-57
  - BEHAVIOR, ACTIVITY PATTERNS**
    - Duck, Tufted, 70(r)
    - Magpie, Yellow-billed, 321(r)
  - , ANTING, 234(r), 322(r)
  - , COMPARATIVE
    - Cictoniidae, 321(r)
  - , DOMINANCE AND AGGRESSION
    - Interspecific, see ecology
    - Sparrow, White-throated, 314-15
  - , FLOCKING
    - Cowbirds, 315-16
    - Grackle, Common, 315-16
    - Jay, Piñon, 69(r)
    - Starling, 315-16
  - , FORAGING, see food and feeding
  - , GENERAL
    - Antbird, Spotted, 75-76(r)
    - Hummingbirds, hermit, 322(r)
- , MAINTENANCE
  - Burrow-digging, Barn Owl, 59-60
  - Cliff-nesting behavior, Kittiwake and Herring Gull, 322(r)
  - Dustbathing, Bobwhite, 234(r)
  - Food-holding with feet, 91-99
  - Huddling, 238(r), 321(r)
  - Sleep, Burrowing Owl, 234(r)
    - , Domestic Pigeon, 234(r)
  - Thermoregulatory, 24, 59-60
- , PAIRING
  - Grackle, Boat-tailed, 68(r)
- , PLAY, 321(r)
- , POSTURES
  - Food-holding, 91-99
  - Footedness, 24, 94
  - Unipedal, 22-26
- , RECOGNITION
  - Nests, eggs and young, Red-winged Blackbird, 70(r), 130(r)
- , ROOSTING
  - Blackbirds, 315-16
- , TERRITORIAL, see territory
- , TOOL USE
  - Cormorant, Double-crested, 130(r)
  - Jay, Blue, 321(r)
  - Sittella, Orange-winged, 129(r)
- Bergstrom, Edward Alexander.
  - Tributes to—, 153-157
- Bittern, American, see *Botaurus lentiginosus*
- Blackbird, Red-winged, see *Agelaius phoeniceus*
- Blindfolds, see orientation
- Bluebird, Eastern, see *Sialia sialis*
- Bobolink, see *Dolichonyx oryzivorus*
- Bonasa umbellus*, parasites of—, 72(r)
- Botaurus lentiginosus*, caught by pelecypods, 65
- Bowerbirds, gardiner, see *Amblyornis brachyramphus marmoratum*, distribution, 326(r)
- Bray, O. E. et al. Migration and seasonal distribution of Common Grackles banded in North and South Dakota. 1-12
- Breeding seasons, see nesting and reproduction
- Breeding success, see nesting and reproduction
- Brooks, W. S. A tentative key for sex determination of Common Redpolls (*Acanthis flammea*) in the northern United States during winter. 13-21
- Browder, J. A. Long-distance movements of Cattle Egrets. 158-170
- Bubulcus ibis*, feeding rates, 158-70
  - , foraging success, 327(r)
  - , long-distance movements, 158-70
  - , starvation, 158-70
- Bucephala* sp., possible hybrids, 326(r)
- Bulbuls, see *Pycnonotidae*
- Bunting, Lazuli, see *Passerina amoena*
  - , Painted, see *Passerina ciris*

- , Reed, see *Emberiza schoeniclus*  
 Caigüe, Blue-headed, see *Pionites melanopephala*  
*Calidris alpina*, classification, 241(r)  
*Capella gallinago*, banding worksheet, 230(r)  
*Carpodacus purpureus*, longevity and distant movement, 227  
*Cassidix mexicanus*, breeding success, territorial behavior, 68(r)  
 —, food habits, 134(r)  
*Cephus grylle*, New England population changes, 267-313  
 Chaffinch, see *Fringilla coelebs*  
 Chat, Yellow-breasted, see *Icteria virens*  
*Chen hyperborea*, arctic fox relationships, 128(r)  
 Ciconiidae, comparative ethology, 321(r)  
*Clangula hyemalis*, delayed nesting, 61-62  
 Clark, G. A., Jr. Unipedal postures in birds, 22-26  
 —. Holding food with the feet in passerines, 91-99  
 Clutch size, see nesting and reproduction  
*Colaptes auratus*, nest spacing and timing, 317-18  
 Columbidae, food-holding with feet, 96  
*Colinus virginianus*, dustbathing, 234(r)  
*Columba livia*, orientation in flight, 102-109  
 —, sleep, 234(r)  
 Columbidae, food-holding with feet, 96  
**COMPETITION, FOOD**  
 Blackbirds and waterfowl, 68(r)  
 —, NEST SITES  
 Bluebird, Eastern, 110-123  
 Woodpecker, Pileated, and Screech Owl, 316  
 Computer records, see banding  
 Conner, R. N. Screech Owl displaces nesting Pileated Woodpeckers, 316  
 Cormorant, Double-crested, see *Phalacrocorax auritus*  
*Corvus coronooides*, storing food, 70(r)  
*Coturnix japonica*, effects of chemical pollutants, 237(r)  
 Cowardin, L. M., and D. A. Davenport. Computerized system for organizing and maintaining files of banding data, 187-195  
 Cowbird, see *Molothrus ater*  
 Crow, see *Corvus*  
 Cocho, New Zealand Long-tailed, see *Eudynamis taitensis*  
 Cuculidae, food-holding with feet, 96  
 Cuthbert, F. J., and W. E. Southern. Ring-billed Gull relocates nest as a result of egg displacement, 225-227  
*Cyanocitta cristata*, tool use, 321(r)
- Cyanocorax beecheii*, age and sex criteria, 81-90  
*C. melanocyanus*, age and sex criteria, 81-90  
*C. sanblasiana*, age and sex criteria, 81-90  
*C. yucatanica*, age and sex criteria, 81-90  
 Daveyport, D. A., see Cowardin, L. M., and —.  
 DeGrazio, J. W., see Bray, O. E., and —.  
*Dendragapus obscurus*, predation on—, 322(r)  
*Dendrocopos pubescens*, age-related eye color, 100-101  
*Dendrocygna eytoni*, tongue structure, 325(r)  
*Dendrocoptes caeruleocephalus*, kills in central Florida, 258-66  
 —, winter area fidelity and survival, 221-224  
*D. coronata*, winter returns, Jamaica, 221-224  
*D. discolor*, winter returns, Jamaica, 221-224  
*D. dominica*, winter returns, Jamaica, 221-224  
*D. magnolia*, winter returns, Mexico, 228-229  
*D. pensylvanica*, winter returns, Jamaica, 221-224  
*D. pharetra*, winter returns, Jamaica, 221-224  
*D. pinus*, winter returns, Jamaica, 221-224  
*D. striata*, kills in central Florida, 258-266  
*D. tigrina*, kills in central Florida, 258-266  
 Diamond, A. W., and R. W. Smith. Returns and survival of banded warblers wintering in Jamaica, 221-224.  
 Diesselhorst, G. Khumbu Himal (rev.) 242  
 Distribution, see Zoogeography and —  
*Dolichonyx oryzivorus*, longevity and breeding, 57-58  
 Drury, W. H. Population changes in New England seabirds, 267-313  
*Dryocopus pileatus*, nesting displacement, 316  
 Duck, Common Eider, see *Somateria mollissima*  
 —, goldeneye, see *Bucephala* sp.  
 —, Greater Scaup, see *Aythya marila*  
 —, Mallard, see *Anas platyrhynchos*  
 —, Pintail, see *Anas acuta*  
 —, Plumed Whistling, see *Dendrocygna eytoni*  
 —, Ruddy, see *Oxyura jamaicensis*  
 —, Shoveller, see *Anas clypeata*  
 —, Tufted, see *Aythya fuligula*  
 Ducks, see Anatidae

- Dunlin, see *Calidris alpina*  
 Eagle, White-tailed, see *Haliaeetus albicilla*  
**ECOLOGY, INTERSPECIFIC RELATIONSHIPS**  
 In St. Croix, 234(r)  
 In Puerto Rico, 234(r)  
 —, GENERAL  
 Arctic-nesting birds, 322(r)  
 Pied Flycatcher breeding, 235(r)  
 Hermit hummingbirds, 322(r)  
 Piñon Jay, 69(r)  
 Mat-grassland birds, 235(r)  
 Egret, Cattle, see *Bubulus ibis*  
 —, Snowy, see *Leucophoyx thula*  
 Ely, C. A. Returns of North American birds to their wintering grounds in southern Mexico, 228-229  
*Emberiza schoeniclus*, biogeography, 240(r)  
*Empidonax difficilis*, migration, age and sex ratios, 205-220  
 E. *flaviventris*, winter returns in Mexico, 228-229  
 E. *minimus*, winter returns in Mexico, 228-229.  
**ENVIRONMENTAL QUALITY**  
 Bird deaths by human litter, 322(r)  
 Chemical influence on egg production, 237(r)  
 Eggshell thinning, owls, 72(r), 323(r)  
 —, White-tailed Eagle, 236(r)  
 Heat loss in oiled ducks, 323(r)  
 Oiled bird care, 72(r)  
 Mercury contamination, 72(r)  
 Erskine, A. J., Buffleheads (rev.), 242-43  
*Eudynamus taitensis*, parasitic and predatory habits, 129(r)  
**EVOLUTION AND GENETICS**  
 Caryotypes of Turdidae, 132(r)  
 Evolution of the genus *Tetrao yattus*, 241(r)  
 — of predator-prey systems, 319(r)  
 — of gardiner bowerbirds, 327(r)  
*Falco sparverius*, breeding biology, 69(r)  
 —, longevity, 66(r)  
 —, predator of Bank Swallows, 327-28(r)  
**FALCONIFORMES**  
 Autumn counts in N. J., 231(r)  
 Autumn counts in Penna., 231(r)  
 Food-holding with feet, 96  
 Foraging by three species, 328-29(r)  
 Winter population trends, 233(r)  
 Farner, D. S., and J. R. King (eds.), Avian biology, vol. 2 (rev.), 329-31  
**FEATHERS**  
 Filoplumes in petrels and shearwaters, 238(r)  
 Substructure, iridescent feathers, 325(r)  
 Water-holding, Sandgrouse feathers, 325(r)  
 Finch, Purple, see *Carpodacus purpureus*  
 Finfoot, American, see *Heliois fulica*  
 Fisk, E. J. Fall movement and probable migrant returns of Mockingbirds in south Florida, 62  
 —, Further notes on the color of Mockingbird eyes, 124  
 Flicker, Yellow-shafted, see *Colaptes auratus*  
 Flight, see locomotion  
 Flocking, see behavior  
 Florida, Bluebird breeding, 110-123  
 Flycatcher, Crested, see *Myiarchus crinitus*  
 —, Least, see *Empidonax minimus*  
 —, Pied, see *Muscicapa hypoleuca*  
 —, Western, see *Empidonax difficilis*  
 —, Yellow-bellied, see *Empidonax flaviventris*  
**FOOD AND FEEDING, CACHING**  
 Crow, Australian, 70(r)  
 Nuthatch, Rock, 134(r)  
 Owl, Barn, 225  
 Woodpecker, Great Spotted, 70(r)  
 —, DIET  
 Pigeon's milk, 73(r)  
 —, FOOD HABITS  
 Duck, Tufted, 70(r)  
 Egret, Cattle, 158-170  
 —, Snowy, bill-vibrating, 328(r)  
 Falconiformes, three species, 328-29(r)  
 Grackle, Great-tailed, 134(r)  
 Jay, nestling Common, 320(r)  
 —, Piñon, 69(r)  
 Kittiwake, 68(r)  
 Kiwi, 328(r)  
 Orioles, nectar-feeding, 134(r)  
 Owl, Barn, 75(r)  
 Swift, White-rumped, 241(r)  
 Terns with multiple fish, 328(r)  
 Wood warblers, 328(r)  
 —, FORAGING EFFICIENCY  
 Egret, Cattle, 327(r)  
 Parent birds with nestlings, 73(r)  
 Footedness, see behavior  
 Franz Josef Land, ornithology of —, 133(r)  
*Fratercula arctica*, New England population changes, 267-313  
 —, sex and age determination, 238(r)  
*Fregata magnificens*, distribution, 129(r)  
 Frigatebird, *Magnificent*, see *Fregata magnificens*  
*Fringilla coelebs*, energy balance, 73(r)  
*Fulmarus glacialisoides*, recoveries, 127(r)  
*Garrulus glandarius*, breeding biology and nestling food, 320(r)  
*Geothlypis trichas*, winter area fidelity, survival, 221-224  
 Goldfinch, Lesser, see *Spinus psaltria*  
 Goose, Snow, see *Chen hyperborea*  
 Grackle, Boat-tailed, see *Cassidix mexicanus*

- , Common, see *Quiscalus quiscula*  
 Grosbeak, Evening, see *Hesperiphona vespertina*  
 —, Rose-breasted, see *Pheucticus ludovicianus*  
 Grouse, Blue, see *Dendragapus obscurus*  
 —, Ruffed, see *Bonasa umbellus*  
 Growth and development, see nesting and reproduction  
 Gruson, E. S. Words for birds (rev.), 243  
 Guarino, J. L. See Bray, O. E., et al.  
 Guillemot, Black, see *Cephus griseus*  
 Gull, Great Black-backed, see *Larus marinus*  
 —, Herring, see *Larus argentatus*  
 —, Red-billed, see *Larus novaehollandiae*  
 —, Ring-billed, see *Larus delawarensis*  
*Gymnorhinus cyanocephalus*, flocking and annual cycle, 69(r)  
*Halcyornis taliapicus*, affinities from lower Eocene, 133(r)  
*Haliaeetus albicilla*, eggshell thinning, 236(r)  
 Hardy, J. W. Age and sex differences in the black-and-blue jays of middle America, 81-90  
 Harrington, B. A. Aggression in winter resident and spring migrant White-throated Sparrows in Massachusetts, 314-15  
 Hatching success, see nesting and reproduction  
 Hawk, Sparrow, see *Falco sparverius*  
 Hawks, see Falconiformes  
 Hearing, see sensory organs  
*Heliothis fulica*, biology in Mexico, 233(r)  
*Helminthesh vermicivorus*, winter returns in Jamaica, 221-224  
 —, winter returns in Mexico— 228-29  
 Heron, Great Blue, see *Ardea herodias*  
*Hesperiphona vespertina*, age determination, 240(r)  
 Hilditch, C. D. M., et al. Autumnal bird migration over Antigua, W. I. 171-179  
 *Himantopus himantopus*, deaths by hail, 67(r)  
*Hirundo rustica*, nest materials, 319-20(r)  
 HISTOLOGY of bird leucocytes, 132(r)  
 Homing, see migration  
 Honeyeater, Fuscous, see *Meliphaga fusca*  
 —, White-naped, see *Melithreptus lunatus*  
 Hummingbirds, see Trochilidae  
 HYBRIDS AND HYBRIDIZATION  
   In the genus *Acanthis*, 13-21  
   In the genus *Bucephala*, 326(r)  
   In the genus *Lanius*, 134(r)  
   In the genus *Myzomela*, 75(r)
- Hydroprogne caspia*, breeding success, 196-204  
 —, comparison of laying date with Common Tern, 129(r)  
 —, longevity, 127(r)  
*Hylocichla mustelina*, winter returns in Mexico, 228-29  
*Hylophylax naevioides*, behavior, 75-76(r)  
*Icteria virens*, winter returns in Mexico, 228-29  
*Icterus*, nectar-feeding by—, 134(r)  
 —, winter identification in the northeast, 132(r)  
 Incubation, see nesting and reproduction  
 Individual distance, see territory  
 INJURIES AND ACCIDENTS  
   Budgerigars, 24  
   Ruddy Ducks, 130(r)  
*Iridoprocne bicolor*, banding worksheet, 230(r)  
 —, individual distance, 321(r)  
 Jamaica, North American Parulidae wintering in—, 221-224  
 James, F. C., see Allen, R. T., and—  
 Jay, Blue, see *Cyanocitta cristata*  
 —, Bushy-crested, see *Cyanocorax melanocyanea*  
 —, Common, see *Garrulus glandarius*  
 —, Piñon, see *Gymnorhinus cyanocephalus*  
 —, San Blas, see *Cyanocorax sanblasianus*  
 —, Yucatan, see *Cyanocorax yucatanicus*  
 Johnson, N. K. Spring migration of the Western Flycatcher, with notes on seasonal changes in sex and age ratios, 205-220  
*Junco hyemalis*, orientation in flight, 102-109  
 Junco, Slate-colored, see *Junco hyemalis*  
 Kaufman, D. W. Captive Barn Owls stockpile prey, 225  
 Kennedy, R. S. Notes on the migration of juvenile Ospreys from Maryland and Virginia, 180-186  
*Ketupa blakistoni*, parental care, 233(r)  
 Kilham, L. Colonial-type nesting in Yellow-shafted Flickers as related to staggering of nesting times, 317-18  
 Kittiwake, see *Rissa*  
 Kiwi, North Island Brown, see *Apteryx australis*  
 Koch, T. Anatomy of the chicken and domestic birds (rev.), 331-32  
*Lanius collurio*, hybrids with *L. phoenicuroides*, 134(r)  
   *L. phoenicuroides*, hybrids with *L. collurio*, 134(r)  
*Larus argentatus*, cliff-nesting behavior, 322(r)

- , New England population changes, 267-313
- , orientation in flight, 102-109
- , predator of swallows, 327-28(r)
- L. delawarensis*, capture method, 67(r)
- , relocates nest, 225-227
- L. marinus*, New England population changes, 267-313
- L. novachollandiae*, band loss, 67(r)
- Laskey, A. R. A longevity and round trip record of Purple Finches, 227
- Leucophoyx thula*, bill-vibrating to attract prey, 328(r)
- Limnothlypis swainsonii*, winter returns in Jamaica, 221-224
- Lockyer, N. F. et al. A device for removing birds from nest boxes, 56-57
- LOCOMOTION**
  - Flight of blindfolded birds, 102-109
  - Flight speed of autumnal migrants, 171-179
  - Flight of swifts at night and in migration, 319(r)
- Longevity, see age
- Magpie, Black-billed, see *Pica pica*
- Magpie, Yellow-billed, see *Pica nuttalli*
- MANAGEMENT AND ECONOMIC ORNITHOLOGY**
  - Aircraft birdstrikes, 235(r)
  - Blue Grouse predation, 322(r)
  - Seabirds and management, 267-313
  - Predation induced by human nest visits, 319(r)
  - Raptor and corvid deterrence, 71(r)
- Martin, D. J. Burrow digging by Barn Owls, 59-60
- Martin, S. G. Longevity surprise: the Bobolink, 57-58
- Massachusetts, tern populations, 27-55
- Measurements, see morphology
- Megapodiidae, food holding with feet, 96
- Meliphaga fusca*, longevity, 127(r)
- Melithreptus lunatus*, longevity, 127(r)
- Melospiza melodia*, orientation in flight, 102-109
- Meyerriecks, A. J. Man and birds: evolution and behavior (rev.), 243-44
- MIGRATION, MECHANICS**
  - Pigeons, 67(r)
  - , NOMADISM
    - Jay, Piñon, 69(r)
  - , PHYSIOLOGY
    - Blood sugar change, 237(r)
    - Body fat chemical change, 237(r)
  - , REGIONAL
    - Antigua, W. I. in autumn, 171-179
    - Florida in autumn, tower kills, 232(r), 258-266
    - N. J., hawks in autumn, 231(r)
    - Ontario, spring diurnal, 232(r)
    - Penna., hawks in autumn, 231(r)
- South China, 232(r)
- Sweden, nocturnal in autumn, 231(r)
- , diurnal in autumn, 231-32(r)
- SPECIES**
  - Anatidae, 128(r)
  - Bobolink, 57-58
  - Egret, Cattle, 158-70
  - Flycatcher, Western, 205-220
  - Grauke, Common, 1-12
  - Hawks, 231(r)
  - Mockingbird, 62
  - Osprey, juveniles, 180-196
  - Pelican, White, 66(r)
  - Sparrowhawk, European, 67(r)
  - Swifts, 319(r)
  - Warbler, Cape May, 258-66
    - , Black-throated Blue, 258-66
    - , Blackpoll, 258-66
  - Mimus polyglottos*, eye color, 124
  - , fall movement in Florida, 62
  - Mix, H. See Williams, T. C., and—
  - Mockingbird, see *Mimus polyglottos*
  - Mniotilla varia*, winter returns in Jamaica, 221-224
  - , winter returns in Mexico, 228-229
  - Molothrus ater*, capture method, 66(r)
  - , pre-roost gatherings, 315-16
  - , populations and migrations, 68(r)
  - Molt, see plumages and—
  - Moreau, R. E. The Palaearctic-African bird migration systems (rev.), 136-37.
  - Morphology and measurements, Redpoll, 13-21
  - Mortality, see age, life expectancy
  - Murre, Common, see *Uria aalge*
  - Murrelet, Marbled, see *Brachyramphus marmoratum*
  - Muscicapa hypoleuca*, breeding ecology, 235(r)
  - Myiarchus crinitus*, winter returns in Mexico, 228-229.
  - Myzantha*, hybridization, 75(r)
  - Myzantha melanoccephala*, nest loss from hail, 67(r)
  - Navigation, see migration; see also orientation
  - Neositta chrysoptera*, tool use, 129(r)
  - NESTING AND REPRODUCTION, BREEDING BIOLOGY**
    - Bluebird, Eastern, 110-123
    - Bulbul, 71(r)
    - Hawk, Sparrow, 69(r)
    - Heron, Great Blue, 320-21(r)
    - Jay, Common, 320(r)
    - Kittiwake, 68(r)
    - Shoveller, 233(r)
    - Swallow, Bank, 128(r)
    - Tern, Caspian, 196-204
    - Weather effects on—, 61-62, 67(r), 110-123
  - CLUTCH SIZE**
    - Bluebird, Eastern, 110-123
    - Seasonal variations, Caspian Tern, 196-204

- , GROWTH AND DEVELOPMENT
  - Effect of gamma radiation on—, 73
  - (r)
  - Kittiwake, 68(r)
  - Oldsquaws, 61-62
- , HABITS
  - Bluebird, Eastern, 110-123
  - Finfoot, American, 233(r)
  - Flycatcher, Pied, 235(r)
  - Hole-nesting birds, 235(r)
  - Oldsquaw, 61-62
  - Owl, Barn, 59-60
  - , Short-eared, 59
- , INCUBATION
  - Bluebird, Eastern, 110-123
  - Effects of weather, 61-62, 67(r), 110-123
  - Oldsquaw, 61-62
  - Tern, Caspian, 196-204
  - In the genus *Sylvia*, 320(r)
- , MULTIPLE BROODS
  - Bluebird, Eastern, 110-123
  - Grosbeak, Rose-breasted, 321(r)
- , NESTS
  - Materials, Barn Swallow, 319-320(r)
  - Nest temperature, 237(r)
  - Relocation, Ring-billed Gull, 225-27
- , PARENTAL CARE, Fishing Owl, 233(r)
- , SEASONS
  - Bluebirds in Florida, 110-123
  - Oldsquaw, 61-62
  - Tern, Arctic, 27-55
  - , Caspian, 129(r), 196-204
  - , Common, 129(r), 27-55
  - , Least, 27-55
  - , Roseate, 27-55
- , SUCCESS
  - Artificial and natural sites, 110-123
  - Bluebird, Eastern, 110-123
  - Grackle, Boat-tailed, 68(r)
  - Tern, Caspian, 196-204
- New England, seabird population changes, 267-313
- Nisbet, I. C. T. Terns in Massachusetts: present numbers and historical changes, 27-55
- , also see Hilditch, C. D. M., and—
- Nuthatch, Rock, see *Sitta tephronota*
- Oceanites oceanicus*, smell and foraging, 127(r)
- Oceanodroma leucorhoa*, New England populations, 267-313
  - , smell and foraging, 127(r)
- Okinawa, birds of—, 326(r)
- Oldsquaw, see *Clangula hyemalis*
- Oporornis agilis*, migration, 261
  - O. formosus*, winter returns in Mexico, 228-29
  - O. philadelphica*, migration, 261
  - O. tolmei*, winter returns in Mexico, 228-29
- ORIENTATION
  - Gull, Herring, 102-109
- Junco, Slate-colored, 102-109
- Pigeon, Domestic, 102-109
- Sparrow, House, 102-109
  - , White-throated, 102-109
- Orioles, see *Icterus*
- Osprey, see *Pandion haliaetus*
- Otus asio*, displacing Pileated Wood-pecker, 316
  - , thin shells of—, 72(r), 323(r)
- Ovenbird, see *Seiurus aurocapillus*
- Owl, Barn, see *Tyto alba*
  - , Blakiston's Fish, see *Ketupa blakistoni*
- , Burrowing, see *Speotyto cunicularia*
- , Little, see *Athene noctua*
- , Saw-whet, see *Aegolius acadicus*
- , Screech, see *Otus asio*
- Owls, extraocular muscles, 74(r)
- , neural hearing centers, 74(r)
- , stockpiling prey, 225
- Oxyura jamaicensis*, colliding with wires, 130(r)
- Pandion haliaetus*, migration, 180-186
- PARASITES AND DISEASE
  - Chiggers, host records, 236(r)
  - Freyanella*, feather mite genus, 323(r)
  - of Ruffed Grouse, 72(r)
  - Mites in birds' nests, 72(r)
  - Mites in House Sparrow nests, 130-31(r)
  - Nasal mites, key, 236(r)
  - Nutrition of feather mites, 236(r)
- PARASITISM BY BIRDS
  - New Zealand Cuckoo, 129(r)
  - Parrot, Thick-billed, see *Rhynchopsitta pachyrhyncha*
  - Parrots, see *Psittacidae*
  - Parulidae, wintering and survival in Jamaica, 221-24
  - Passer domesticus*, orientation, 102-109
  - , nest mites, 130-31(r)
  - Passerina amoena*, banding worksheet, 230(r)
    - P. ciris*, winter returns in Mexico, 228-29
  - Patterson, B. N. More on one-night mileage of migrants, 227-28
  - Pelecanus erythrorhynchos*, banding and recoveries, 66(r)
  - Pelican, White, see *Pelecanus erythrorhynchos*
  - Petrel, Leach's, see *Oceanodroma leucorhoa*
    - , Silver-grey, see *Fulmarus glacialis*
    - , Wilson's, see *Oceanites oceanicus*
  - Phalacrocorax auritus*, air-sacs, 236(r)
  - , New England populations, 267-313
  - , Tool use, 130(r)
  - Pheucticus ludovicianus*, brood overlap, 321(r)
  - Physiology, see Anatomy and—
  - Pica pica*, capture method, 66(r)
    - P. nuttalli*, time budgets, 321(r)
  - Picathartes*, relationships of—, 327(r)

- Pigeon, Domestic, see *Columba livia*  
*Pionites melanocephala*, plumage care, 322(r)  
*Piranga rubra*, banding worksheet, 230(r)
- PLUMAGES AND MOLT  
Bobolink, 57-58  
Flycatcher, Western, 205-220  
Jay, Beechey, 81-90  
—, Bushy-crested, 81-90  
—, San Blas, 81-90  
—, Yucatan, 81-90  
Kiwi, 325(r)  
Oldsquaw, 61-62  
Orioles, 132(r)  
Vireo, White-eyed, 63-65
- POLYMORPHISM  
Shearwater, Wedge-tailed, 128(r)  
Sparrow, White-throated, 314-15
- POPULATIONS AND SOCIAL BIOLOGY  
Blackbirds, 68(r)  
Estimating, 28, 66(r), 68(r)  
Jay, Piñon, 69(r)  
Seabirds in New England, 267-313  
Shearwaters of Shark Bay, 128(r)  
Terns in Mass., 27-55  
—, PRODUCTIVITY  
Bluebird, Eastern, 110-123  
Latitudinal variation, 71(r)  
Swallow, Rough-winged, 71(r)  
—, RECRUITMENT  
Swallow, Rough-winged, 71(r)  
Post, W. Some encounters between birds and pelecypods., 65  
Postures, see behavior
- PREDATION  
of Bank Swallows by Sparrow Hawks, 327-28(r)  
of Blue Grouse, 322(r)  
of Bluebird nests, 110-123  
Nest predation induced by humans, 319(r)  
by New Zealand Cuckoo, 129(r)  
Predator-prey evolution, 319(r)  
of Shearwaters, 128(r)  
of Snow Geese by fox, 128(r)
- Psittacidae, extinct and vanishing species, 234(r)  
—, food-holding with feet, 96
- Puffin, Common, see *Fratercula arctica*  
*Puffinus gravis*, smell and foraging, 127(r)  
*P. pacificus*, breeding populations, 128(r)
- Pyenonotidae, biology, 71(r)
- Quail, Bobwhite, see *Colinus virginianus*  
—, Japanese, see *Coturnix japonica*  
*Quiscalus mexicanus*, see *Cassidix mexicanus*  
*Q. quiscula*, migration and distribution, 1-12  
—, roosting associates and behavior, 315-16
- Radar, see migration; also see research methods  
Rallidae, food-holding with feet, 96  
Razorbill, see *Alca torda*  
Recoveries, see banding  
Redpoll, Common, see *Acanthis flammea*  
—, Hoary, see *Acanthis hornemannii*  
Redstart, American, see *Setophaga ruticilla*  
*Regulus regulus*, huddling, 238(r)
- RESEARCH METHODS  
Computation of pesticide residues, 323(r)  
Recording data from radar, 124-25  
*Rhynchositta pachyrhyncha*, sex determination, 325-26(r)
- Ricklefs, R. E. Tattooing nestlings for individual recognition, 63  
*Riparia riparia*, nesting in sawdust, 128(r)  
—, prey of Sparrow Hawk, 327-28(r)  
*Rissa tridactyla*, breeding biology in Newfoundland, 68(r)  
—, cliff-nesting behavior, 322(r)  
Roosting, see behavior  
Royall, W. C. See Bray, O. E., and—  
*Rynchops nigra*, human-caused deaths, 323(r)  
—, New England populations, 267-313  
Ryukyu Island, birds of—, 326(r)
- Salvadori, A., and K. A. Youngstrom. A system survey of a bird observatory: Part I. A recording form for banding data. 249-257
- Scaup, Greater, see *Aythya marila*  
Schaeffer, F. S. Tactile bristles of Saw-whet Owls are sensitive to touch, 125  
Seabirds, population changes in New England, 267-313  
—, zoogeography, 132(r)  
*Sciurus aurocapillus*, aging criteria, 239-40(r)  
—, winter returns in Jamaica, 221-24  
—, winter returns in Mexico, 228-29  
*S. noveboracensis*, winter returns in Jamaica, 221-24
- SENSORY ORGANS  
Hearing, neural centers, 74(r)  
Smell, in procellariids, 127(r)  
Smell and touch in Kiwis, 328(r)  
Touch, Saw-whet Owl, 125
- Setophaga ruticilla*, winter returns in Jamaica, 221-24
- SEX AND AGE RATIOS  
of Black-throated Blue and Cape May warblers, 258-66  
of Sparrow Hawks, 69(r)  
of Western Flycatchers, 205-220
- SEX CRITERIA AND DIMORPHISM  
Flycatcher, Western, 205-220  
Jay, San Blas, 83-85  
—, Yucatan, 85-87

- Parrot, Thick-billed, 325-26(r)  
 Puffin, Common, 238(r)  
 Redpoll, Common, 13-21  
 Shearwater, Greater, see *Puffinus gravis*  
 —, Wedge-tailed, see *Puffinus pacificus*  
 Shearwaters, see *Puffinus*  
 Shorebirds, autumnal migration, 171-79  
*Sialia sialis*, breeding biology, 110-123  
 Sickness, see injuries  
 Siskin, Pine, see *Spinus pinus*  
 Site tenacity, see territory  
*Sitta tephronota*, food storing, 134(r)  
 Sittella, Orange-winged, see *Neositta chrysopetra*  
 Skimmer, Black, see *Rynchops nigra*  
 Smith, R. W. See Diamond, A. W., and —  
 Snipe, Common, see *Capella gallinago*  
 Social biology, see populations and —  
 Soikkeli, M. Breeding success of the Caspian Tern in Finland, 196-204  
*Somateria mollissima*, New England population changes, 267-313  
 Song, see vocalizations  
 Southern, W. E. See Cuthbert, F. J., and —  
 Sparrow, House, see *Passer domesticus*  
 —, Song, see *Melospiza melodia*  
 —, White-crowned, see *Zonotrichia leucophrys*  
 —, White-throated, see *Zonotrichia albicollis*  
*Speotyto cunicularia*, nest locations, 59(r)  
 —, sleep in —, 234(r)  
*Spinus pinus*, movements, 227-228  
 —, *S. psaltria*, banding worksheet, 230(r)  
 Starling, see *Sturnus vulgaris*  
*Stelgidopteryx ruficollis*, breeding productivity, 71(r)  
*Sterna albifrons*, Massachusetts populations, 27-55  
 —, New England population changes, 267-313  
 —, *S. bengalensis*, display with *S. sandvicensis*, 326(r)  
 —, *S. dougallii*, New England populations, 27-55, 267-313  
 —, *S. hirundo*, clamped by clam, 65  
 —, clutch initiation, 129(r)  
 —, human-caused deaths, 323(r)  
 —, New England populations, 27-55, 267-313  
 —, recoveries of — banded in Sask., 127(r)  
 —, *S. paradisea*, New England populations, 27-55, 267-313  
 —, *S. sandvicensis*, display with *S. bengalensis*, 326(r)  
 —, sp., carrying multiple fish, 328(r)  
*Sturnus vulgaris*, populations and migration, 68(r)  
 —, pre-roost gatherings, 315-16  
 Stewart, P. A. Basis for pre-roost gatherings of Starlings and Brown-headed Cowbirds, 315-16  
 Stone, C. P. See Lockyer, N. F., and —  
 Strigiformes, food-holding with feet, 96  
 Swallow, Bank, see *Riparia riparia*  
 —, Barn, see *Hirundo rustica*  
 —, Rough-winged, see *Stelgidopteryx ruficollis*  
 —, Tree, see *Iridoprocne bicolor*  
 —, Violet-green, see *Tachycineta thalassina*  
 Swifts, see *Apus apus*  
*Sylvia* sp., incubation behavior and patches, 320(r)
- SYSTEMATICS AND TAXONOMY**
- Dunlin, 241(r)  
*Halcyornis toltecus*, 133  
 Owl, Little, 326-27(r)  
*Picathartes*, relationships, 327  
 Use of rare characters in —, 327(r)  
*Tachycineta thalassina*, Banding worksheet, 230(r)  
 Tanager, Summer, see *Piranga rubra*  
 Tatooing, see banding  
 Taylor, W. K. Black-throated Blue and Cape May warblers killed in central Florida, 258-266  
 Teal, J. M. See Williams, T. C., and —  
 Tern, Arctic, see *Sterna paradisaea*  
 —, Caspian, see *Hydroprogne caspia*  
 —, Common, see *Sterna hirundo*  
 —, Least, see *Sterna albifrons*  
 —, Roseate, see *Sterna dougallii*  
 —, Royal, see *Thalasseus maximus*
- TERRITORY AND AREA**
- FIDELITY**
- Blackbird, Red-winged, seasonal changes, 70(r)  
 Bobolink, 57  
 Grackle, Boat-tailed, 68(r)  
 Individual distance, Tree Swallow, 321(r)  
 Nest spacing, Yellow-shafted Flicker, 317-18  
 North American Parulidae wintering in Jamaica, 221-224  
 North American birds wintering in Mexico, 228-229  
 Territorial songs of birds, 134(r)  
*Tetraogallus*, evolution of the genus, 241(r)  
*Thalasseus maximus*, recoveries of — banded in Virginia, 240-241(r)  
 Thermoregulation, see behavior  
 Thompson, C. F. Postjuvenile molt in the White-eyed Vireo, 63-65  
 Thrush, Wood, see *Hylocichla mustelina*  
 Thrushes, see Turdidae  
 Time budgets, see behavior  
 Trapping, see banding techniques  
 Trochilidae, behavior and ecology, 322(r)  
 Troganidae, food-holding with feet, 96

- Tuck, L. M. The snipes: a study of the genus *Capella* (rev.), 244
- Turdidae, comparative karyotypes, 132(r)
- Tyto alba*, banding report, Australia, 231(r)
- , behavior, 59
  - , burrow-digging, 59-60
  - , prey, 75(r)
  - , stockpiling prey, 225
- Unipedal postures, see behavior
- Uria aalge*, New England population changes, 267-313
- Vartapetyan, B. B., Molecular oxygen and water in cell metabolism (rev.), 244-45
- Vermivora peregrina*, wintering in Jamaica, 221
- V. ruficapilla*, migration, 261
- Vireo griseus*, postjuvenile molt, 63-65
- V. solitarius*, winter returns in Mexico, 228-229
- Vireo, Solitary, see *Vireo solitarius*
- , White-eyed, see *Vireo griseus*
- VISION
- Alteration of avian eye growth, 131(r)
  - Avian retina, ultrastructure, 238(r)
  - Extraocular muscles of birds, 74(r)
  - Lens damage in sharks by UV light, 324-25(r)
  - Pecten oculi in birds, 132(r), 324(r)
  - Photopic spectral sensitivity, 131(r)
  - Retinal damage in pigeons, 131(r)
  - Subcortical vision center in pigeons, 131(r)
- VOCALIZATIONS AND SOUND
- Anatidae, sound production, 239(r)
  - Blackbird, Red-winged, nestlings, 70(r)
  - Musical patterns, evolution in birds, 329(r)
  - Gackle, Boat-tailed, 68(r)
  - Passerines, syrinx, 239(r)
  - Territorial and courtship songs, 134(r)
  - Sound and orientation, 102-109
- Warbler, Black-and-white, see *Mniotilla varia*
- , Black-throated Blue, see *Dendroica caeruleascens*
  - , Cape May, see *Dendroica tigrina*
  - , Connecticut, see *Oporornis agilis*
  - , Hooded, see *Wilsonia citrina*
  - , Kentucky, see *Oporornis formosus*
  - , Magnolia, see *Dendroica magnolia*
  - , MacGillivray's, see *Oporornis tollimie*
  - , Mourning, see *Oporornis philadelphica*
  - , Nashville, see *Vermivora ruficapilla*
  - , Parula, see *Parula americana*
  - , Prairie, see *Dendroica discolor*
- , Swainson's, see *Limnothlypis swainsonii*
- Warblers, see Parulidae
- Waterthrush, Northern, see *Seiurus noveboracensis*
- Weather, effects on breeding, see nesting and reproduction
- , effects on bird mortality, see age
- Weights, migrant warblers, 258-266
- West Virginia, Birds of—, 326(r)
- White, S. C., and G. E. Woolfenden. Breeding of the Eastern Bluebird in central Florida, 110-123
- Williams, T. C., and J. M. Teal. The flight of blindfolded birds, 102-109
- , and H. Mix. An inexpensive apparatus for recording data from radar Plan Position Indicator displays, 124-125
  - , see Hilditch, C. D. M., and —
- Willis, E. O. The behavior of Spotted Antbirds (rev.), 75-76
- Wilsonia citrina*, winter studies in Mexico, 228-229
- pusilla*, migration, 261
- Wood, D. L., see Wood, D. S., and —
- Wood, D. S., and D. L. Wood. Quantitative iris color change with age in Downy Woodpeckers, 100-101
- Wood, M. Birds of Pennsylvania (rev.), 245
- Wood warblers, food of—, 328(r)
- Woodpecker, Downy, see *Dendrocopos pubescens*
- , Great Spotted, see *Dendrocopos major*
  - , Pileated, see *Dryocopus pileatus*
- Woolfenden, G. E., see White, S. C., and —
- Youngstrom, K. A., see Salvadori, A., and —
- Zonotrichia albicollis*, aggression, 314-315
- , orientation in flight, 102-109
- Z. leucophrys*, in Ohio, 74(r)
- ZOOGEOGRAPHY AND DISTRIBUTION, REGIONAL
- Franz Josef Land, 133(r)
  - Okinawa and Ryukyu Islands, 326(r)
  - Southern Appalachian, 74(r)
  - West Virginia, 326(r)
- , SPECIES
- Accipiter nisus* in winter, 67(r)
  - Bowerbirds, 327(r)
  - Bunting, Reed, 240(r)
  - Duck, Pintail, 240(r)
  - Egret, Cattle, 158-170
  - Flycatcher, Western, 205-220
  - Frigatebird, Magnificent, 129(r)
  - Gackle, Common, 1-12
  - Murrelet, Marbled, 326(r)
  - Seabirds, 132(r)
  - Sparrow, White-crowned, 74(r)