

- GRIFFIN, D. R. 1943. Homing experiments with Herring Gulls and Common Terns. *Bird-Banding*, **14**: 7-33.
- HARRIS, S. W. 1952. A throw net for capturing female waterfowl on the nest. *J. Wildl. Manage.*, **16**: 515.
- HILDÉN, O., AND S. VUOLANTO. 1972. Breeding biology of the Red-necked Phalarope *Phalaropus lobatus* in Finland. *Ornis Fenn.*, **49**: 57-85.
- HOWE, M. A. 1975. Behavioral aspects of the pair bond in Wilson's Phalarope. *Wilson Bull.*, **87**: 248-270.
- JOHNS, J. E. 1963. A new method of capture utilizing the mist net. *Bird-Banding*, **34**: 209-213.
- MARTIN, S. G. 1969. A technique for capturing nesting grassland birds with mist nets. *Bird-Banding*, **40**: 233-237.
- ORING, L. W., AND M. L. KNUDSON. 1972. Monogamy and polyandry in the Spotted Sandpiper. *Living Bird*, **11**: 59-73.
- SCHAMEL, D., AND D. TRACY. 1977. Polyandry, replacement clutches and site tenacity in the Red Phalarope (*Phalaropus fulicarius*) at Barrow, Alaska. *Bird-Banding*, **48**: 314-324.
- SOWLS, L. K. 1949. A preliminary report on reneating in waterfowl. *Trans. 14th N. Am. Wildl. Conf.*, **14**: 260-275.
- WEAVER, D. K., AND J. A. KADLEC. 1970. A method for trapping breeding adult gulls. *Bird-Banding*, **41**: 28-31.
- WELLER, M. W. 1957. An automatic nest-trap for waterfowl. *J. Wildl. Manage.*, **21**: 456-458.
- CYNTHIA M. KAGARISE, *Division of Biological Sciences, University of Michigan, Ann Arbor, MI 48109*. Received 12 January 1978, accepted 18 May 1978.

**Play by Ring-billed Gulls.**—Following upon Jarvis and Southern's account (*Wilson Bull.*, **88**: 621-631, 1976) of the food habits of the Ring-billed Gull (*Larus delawarensis*), Ficken's review of avian play (*Auk*, **94**: 573-582, 1977) prompts me to report activities of this gull that appear to be play. The activities seem adaptive as practice of catching or retrieving prey.

Jarvis and Southern found that fish comprise 76% (volume) of the diet of Ring-bills at two Michigan breeding colonies. Ring-bills use a pond at Woodlawn, Baltimore County, MD, as a daytime loafing place in winter, and from 1969 to 1972, when Southern was color-tagging at his colonies, individuals from both were among the birds at Woodlawn. At this pond fish 4 to 33 cm long are commonly caught by herons in summer and by Herring Gulls (*L. argentatus*) in winter, yet in 10 years of visiting the pond I have never seen Ring-bills make a catch, although occasionally one picks briefly at a carcass abandoned by a Herring Gull. (Where they do their feeding, I do not know.)

Nevertheless the Ring-bills, both subadults and adults, do a great deal of diving. Sometimes they lunge forward from the surface, sometimes rise 1 m or more in the air to plunge vertically into the water, which is probably nowhere more than 30 cm deep. When they bring up anything, it is always trash (sodden leaves or twigs), but most often they bring up nothing at all. Several times, when their diving has been persistent at a particular spot, a Herring Gull has been attracted but after swimming about the area, has gone away, confirming that no prey was there.

It would seem, then, that this diving is a mere pastime. This must also be true of the chases that occur when a gull flies up with a piece of trash; the chases are so close and twisting that the pursuer must surely see that it is not food the other bird is carrying. And still a third activity consists of dropping things from the air and catching them before they reach the ground or water; examples from my notes are:

9 March 1970: From heights of perhaps 13 m above the water an adult Ring-bill was dropping a small object resembling a bottle cap and then making practically vertical,

slightly twisting dives of some meters and catching it. It did this three times while I watched, then on a fourth drop the object fell into the water and was lost.

25 November 1972: An adult flew about fairly high with something in its bill that looked like a small ball. Five times it dropped the object, then dived steeply and caught it from a few to a number of meters down. On a sixth drop it failed to make the catch, and abandoned it.

28 December 1972: An adult on the water lunged and got a sodden leaf, flew up in the air a meter or so with it, dropped it into the water (where it sank) flew down and got it again, and again flew up and dropped it. Over and over it did this. Shortly several more Ring-bills, including at least one subadult, began doing the same thing, and some others tried to snatch the leaves from the holders. From the size, shape and color I could tell with certainty that it was always the same leaf that a bird retrieved and dropped again. These activities went on for at least 5 min after I first noticed them. Ficken (op. cit., 577, 578) cites a report of contagious play by captive Common Ravens (*Corvus corax*).—HERVEY BRACKBILL, 2620 Poplar Drive, Baltimore, MD 21207. Received 10 February 1978, accepted 19 May 1978.

**Temperature and Snowfall Effects on Feeding Station Activity.**—The feeding activity of birds often appears to increase on colder days or days with snowfall. A quantitative study of such weather effects became possible by using an automatic feeder designed to count total feeding visits per day. The feeding station was a simple tray mounted five feet above the ground in a small yard in Princeton, Mercer County, New Jersey. Tray walls allowed access only to one side of the feeder, and a photocell projected across this side. Any bird visiting the station interrupted the light beam, causing a relay to record a count. The actual circuit details of the automatic recorder have been reported earlier (Leck and Leck, *Amer. Birds*, **30**: 140–141, 1976).

The feeder was operated from 11 November 1973 to 27 February 1974, and 10,629 visits were recorded. Field data included total number of visits and maximum–minimum temperatures each day (excluding a December vacation period). The most common bird was the locally abundant House Finch (*Carpodacus mexicanus*), but Blue Jays (*Cyanocitta cristata*), Starlings (*Sturnus vulgaris*), House Sparrows (*Passer domesticus*), and other species also frequented this urban feeder. (A feeding station on the ground might well have added additional birds, but the results would probably parallel those from this elevated station.) The seed was a commercial mixture of cracked corn, millet, and sunflower. Changes in feeding station activity with different weather conditions were dramatic. The results are best presented in a chronological review:

#### NOVEMBER (315 visits)

11–30 November. Mild with daily maxima usually between 40° and 60°F. Several record warm days (max. 75°F). Average 16 visits per day with few or none on warmer days. Most active day (50 visits) was also the coldest.

#### DECEMBER (1,803 visits)

1–15 December. Mostly mild, including several warm days (max. 62°). Daily maxima usually between 30° and 54°F. Average 35 visits per day.

16–19 December. SNOWSTORM (Heaviest snowfall in two years); daily minima 10°–16°F. Activity increases with an average of 395 visits per day.

20 December. Mild; reduced activity (91 visits).

#### JANUARY (5,638 visits)

1–8 January. Seasonally cool; daily minima 20°–30°F. Average 100 visits per day.

9–11 January. HEAVY SNOWSTORM (three-inch accumulation). Rapid increase in activity with an average of 865 visits per day.

12–14 January. Cold with snow cover. Average 484 visits per day.

15–17 January. Mild weather; daily maxima 40°–50°F. Average 150 visits per day.