

- _____. 1981. On the ecology and biology of Tengmalm's Owl (*Aegolius funereus*) in Southern Ostrobothnia and Suomenselkä, western Finland. *Acta Univ. Ouluensis ser. A. Sci. rerum not.* 118 Biol. 13:1-84.
- _____. 1984a. Population dynamics of birds of prey in relation to fluctuations in small mammal populations in western Finland. *Annales Zoologici Fennici* 21 (in print).
- _____. 1984b. Optimal diet of the Kestrel *Falco tinnunculus* in breeding season. *Ornis Fennica* 61 (in print).
- _____. 1984c. Prey choice strategies of the Kestrel *Falco tinnunculus* in relation to small mammal abundance and Finnish birds of prey. *Annales Zoologici Fennici* 21 (in print).
- _____, S. IKOLA, R. HAAPOJA, AND J. KIRKKOMÄKI. 1977. On the ecology of Long-eared, Short-eared and Tengmalm's Owls as well as Kestrel and Hen Harrier in Alajoki in 1977. *Suomenselän Linnut* 12:100-117.
- _____, E., S. IKOLA, R. HAAPOJA, AND O. HEMMINKI. 1979. On the occurrence and breeding of raptors in Alajoki in 1978. *Suomenselän Linnut* 14:44-51.
- MASCHER, J.W. 1963. Tornfalk (*Falco tinnunculus*) övertar byte från jorduggla (*Asio flammeus*) i flykten. *Vår Fågelvärld* 22:293-294.
- MIKKOLA, H. 1983. Owls of Europe. T & A D Poyser, Calton. 397 pp.
- NILSSON, I. 1975. Tornfalk parasiterar på jorduggla. *Anser* 14:133.
- REESE, R.A., AND E. BALFOUR. 1973. Food piracy between kestrels and short-eared owls. *British Birds* 66:227-228.
- WATSON, D. 1977. The Hen Harrier, T & A D Poyser, Calton. 307 pp.
- WOOD, C.R. 1976. Piratical short-eared owl. *British Birds* 69:272.

Kp.4, SF-62200 Kauhava, FINLAND

Received 17 December 1983; Accepted 4 June 1984

An Unusual Observation of 'Homing' To Prey By A Migrating Immature Peregrine Falcon

CARL SAFINA

On 6 October 1981 at 10:05 EST, while operating a raptor banding station on the Long Island, New York barrier beach, I saw a hatching year female Peregrine Falcon (*Falco peregrinus*) land approximately 50 m from my blind and begin eating a small passerine (probably a White-throated Sparrow (*Zonotrichia albicollis*, based on feathers). The falcon was on a low, beach heather (*Hudsonia*) covered rise in a broad, open rolling area. Soon after the falcon began eating, an Osprey (*Pandion haliaetus*), carrying a fish, came over the falcon, vocalizing. The Peregrine flew up and chased the Osprey over the bay to the north, stooping repeatedly at it before flying out of sight. Within 5 min the falcon reappeared on its kill. As the prey was small, cryptically colored, and in a broad, non-descript area of the beach, and since the Peregrine was not seen searching for it, apparently the falcon remembered exactly where it had left its kill and was able to return there from a point out of sight.

Falcons frequently return to cached prey. What is interesting about this incident is that the falcon was a first year migrant and the topography was almost certainly not familiar (the banding station had been manned daily for 3 w prior, and no Peregrines appeared to have been staying in the area). Enderson (Auk. 81:332-352, 1964) described wintering Prairie Falcons (*Falco mexicanus*) leaving their prey on the ground and driving off other rapors in a similar manner. His falcons seldom had difficulty relocating the prey, but this is not surprising because they were familiar with the area.

National Audubon Society, Scully Sanctuary, 306 South Bay Avenue, Islip, NY 11751.

Received 30 March 1983; Accepted 30 April 1984

Errata - Raptor Research 18(2)

Page 44 (Table 1 concluded), 0.420^g should appear in the column for shell thickness, 0.49^h should appear in the column for HE and 0.27^h should appear in the column for DDE; page 47, paragraph 3, line 6, > 8 ppm should appear as ≥ 8 ppm; page 61, Literature Cited, the Sawby et al. reference appeared in *Condor* 76:479-481; page 70, photographs are reversed.