

ACKNOWLEDGMENTS

I thank B. Lehman, R. Hipp, and J. Boyce for companionship and assistance in the field. T. Kaiser, B. Millsap, J. Parrish, D. Runde and C. White provided critical review of the manuscript. The U.S. Bureau of Land Management, Dr. and Mrs. D. A. Boyce, and Mr. and Mrs. R. T. Whipple provided financial support for this study. I dedicate this article to D. Britt who loved wildlife, especially birds of prey.

LITERATURE CITED

- BOYCE, D. A., JR. 1982. Prairie Falcon fledgling productivity in the Mojave Desert, California. M.S. Thesis. Humboldt State University, Arcata, CA. 97 pp.
- BOYCE, D. A., L. FISHER, W. E. LEHMAN, B. HIPP AND J. PETERSON. 1980. Prairie Falcons nest on an artificial ledge. *Raptor Res.* 14:46-50.
- , C. M. WHITE, R. ESCANO AND W. E. LEHMAN. 1982. Enhancement of cliffs for nesting Peregrine Falcons. *Wildl. Soc. Bull.* 10:380-381.
- , R. L. GARRETT AND B. J. WALTON. 1986. Distribution and density of Prairie Falcons nesting in California during the 1970s. *Raptor Res.* 20:71-74.
- BROWN, L. H. AND D. AMADON. 1968. Eagles, hawks and falcons of the world. Country Life Books, London.
- DECKER, F. R. 1931. Prairie Falcon's nest. *The Oologist* 48:96.
- DENTON, S. J. 1975. Status of Prairie Falcons breeding in Oregon. M.S. Thesis. Oregon State Univ., Corvallis. 58 pp.
- DICE, L. R. 1943. The biotic provinces of North America. Univ. Mich. Press, Ann Arbor. 76 pp.
- ENDERSON, J. H. 1964. A study of the Prairie Falcon in central Rocky Mountain region. *Auk* 81:332-352.
- JAEGER, E. C. 1957. The North American Deserts. Stanford Univ. Press, Stanford, CA. 308 pp.
- LEEDY, R. R. 1972. The status of the Prairie Falcon in western Montana: special emphasis on possible effects of chlorinated hydrocarbon insecticides. M.S. Thesis. Univ. of Montana, Missoula. 96 pp.
- MACLAREN, P. A., D. E. RUNDE AND S. H. ANDERSON. 1984. A record of tree-nesting Prairie Falcons in Wyoming. *Condor* 86:487.
- MILLSAP, B. A. 1984. Distributional status of Falconiformes in westcentral Arizona . . . with notes on ecology, reproductive success, and management. U.S. Dept. of Interior, B.L.M. Tech. Note 355.
- OGDEN, V. T. 1973. Nesting density and reproductive success of Prairie Falcons in southwestern Idaho. M.S. Thesis. Univ. of Idaho, Moscow. 39 pp.
- AND M. G. HORNOCKER. 1977. Nesting density and reproductive success of Prairie Falcons in southwestern Idaho. *J. Wildl. Manage.* 41:1-11.
- PLATT, S. W. 1974. Breeding status and distribution of the Prairie Falcon in northern New Mexico. M.S. Thesis. Oklahoma State Univ., Stillwater. 68 pp.
- PORTER, R. D. AND C. M. WHITE. 1973. The Peregrine Falcon in Utah, emphasizing ecology and competition with the Prairie Falcon. *Brigham Young Univ. Sci. Bull.* 28:1-74.
- RUNDE, D. E. AND S. H. ANDERSON. 1986. Characteristics of cliffs and nest sites used by breeding Prairie Falcons. *Raptor Res.* 20:21-28.
- TYLER, J. G. 1923. Observations on the habits of the Prairie Falcon. *Condor* 25:95-97.
- WILLIAMS, R. N. 1981. Breeding ecology of Prairie Falcons at high elevations in central Colorado. M.S. Thesis. Brigham Young Univ., Provo, UT. 44 pp.
- . 1984. Eyrie aspect as a compensator for ambient temperature fluctuations: a preliminary investigation. *Raptor Res.* 18:153-155.
- ZAR, J. H. 1984. Biostatistical Analysis. Second ed. Prentice Hall. Englewood Cliffs, NJ. 736 pp.

Department of Zoology, 574 WIDB, Brigham Young University, Provo, UT 84602.

Received 28 October 1986; Accepted 14 January 1987

J. Raptor Res. 21(1):38-39

© 1987 The Raptor Research Foundation, Inc.

ADULT PAIR OF MERLINS IN SOUTHERN UTAH IN JUNE

JAMES E. SAILER

While participating in a Utah Peregrine Falcon (*Falco peregrinus*) survey, the author and a co-observer sighted and confirmed the presence of an adult pair of Merlins (*Falco columbarius*) in southern Utah. The falcons were light in color and were thought to be *F. c. richardsonii*. Positive identification of the subspecies by a tail-band count was not possible.

Observations occurred on 1 June 1984, between 0955 and 1008 H (MDT), south of Bryce Canyon National Park (approx. 112°20'W longitude, 37°27'N latitude). Observations were made using a spotting scope (15 × 30). The female Merlin was sighted at 0955 H perched in a snag at the top of a 120 m, southeast-facing cliff, approximately 0.5 km from the observation site. At 0958 H she

made a shallow stoop to the northeast, parallel to the cliff face, and was lost from view behind a large outcropping on the cliff. Upon her disappearance, an adult male Merlin was sighted in the immediate area perched on a small rock spire of the outcropping. He remained until 1008 H and was subsequently lost from view. Neither falcon was seen again during the remaining two hr of observations. Further investigations were not possible due to the continuing Peregrine survey, and we were unable to document the Merlins' breeding status. The area of the sightings was approximately 2650 m above sea level, dropping 425 m in one km to the valley below. Flora above the cliff consisted chiefly of ponderosa pine (*Pinus ponderosa*), white fir (*Abies concolor*), and Douglas fir (*Pseudotsuga menziesii*). Habitat below the cliff varied from Pinyon pine (*Pinus edulis*) and Utah Juniper (*Juniperus osteosperma*) to grasslands as the elevation decreased (Elias 1980).

Two additional Merlin sightings were made in Canyonlands National Park on 17 and 18 June from two observation points approximately two km apart. Both Merlins sighted were adult males and were thought to be the same bird. Observations were made from the rim of the Colorado River Canyon on both occasions. Flora above the canyon was sparse; however, the canyon floor and many small side canyons in the area maintained populations of pine (*Pinus* sp.). The Merlin's breeding range extends from the taiga through the prairie-parkland biomes (Temple 1972). During the 19th century, it was found as a breeding bird in the Wasatch Mountains of northern Utah (Hayward et al. 1976), although no subsequent breeding records have been reported for this region (Oliphant 1985). Bailey and Niedrach (1965) considered this species a rare winter migrant of Utah, while Hayward et al. (1976) classified it as a sparse breeding resident. The nearest recent breeding by Merlins has been found in southern Idaho, southern Wyoming, northern Colorado and Nebraska (Lock and Craig 1975; Craig and Renn 1977; Lock 1979; Oliphant 1985; Becker pers. comm.).

Occurrence of this species outside of its documented breeding range during the summer months suggests it is not only a winter migrant within Utah as previously indicated. My observation dates fall within incubation and hatching periods of *F. c. richardsonii* in southeastern Montana (Becker and Sieg 1985) and correspond temporally with reports of Merlin breeding chronologies in south-

central Idaho (Craig and Renn 1977). Becker and Sieg (1985) reported that spring arrivals of Merlins at their study area in southeastern Montana occurred between 11 March and 23 April. Based upon this evidence, Merlins observed during my survey were not thought to be migrant birds returning north and may represent a breeding component in Utah.

ACKNOWLEDGMENTS

Ritt Enderson was co-observer and confirmed these sightings in the field. We thank the Peregrine Fund, Inc., and the National Park Service for enabling us to make these observations while participating in the 1984 Peregrine survey. Christopher Servheen, Dale Becker and Lynn Oliphant reviewed and commented on the manuscript.

LITERATURE CITED

- BAILEY, A. M. AND R. J. NIEDRACH. 1965. Birds of Colorado. Vol. I. Denver Mus. Nat. Hist., Denver, CO. Pp. 246-248.
- BECKER, D. M. AND C. H. SIEG. 1985. Breeding chronology and reproductive success of Richardson's Merlins in southeastern Montana. *Raptor Res.* 19(2/3):52-55.
- CRAIG, T. AND F. RENN. 1977. Recent nestings of the Merlin in Idaho. *Condor* 79:392.
- ELIAS, T. 1980. Trees of North America. Time Mirror Magazines, NY.
- HAYWARD, C. L., C. COTTAM, A. M. WOODBURY AND H. H. FROST. 1976. Birds of Utah. Gr. Basin Natural. Memoirs. Brigham Young Univ., Provo, UT. 229 pp.
- LOCK, R. A. 1979. A second nesting record for Merlins in Nebraska. *Nebraska Bird Rev.* 42(2):39.
- AND R. CRAIG. 1975. Merlin nest in Nebraska. *Nebraska Bird Rev.* 43:78-79.
- OLIPHANT, L. W. 1985. North American Merlin breeding survey. *Raptor Res.* 19(2/3):37-41.
- TEMPLE, S. A. 1972. Systematics and evolution of the North American Merlins. *Auk* 89:325-338.

**P.O. Box 5832, Missoula, MT 59806. Present address:
RD #1, Box 43A, Glen Gardner, NJ 08826.**

Received 1 June 1986; Accepted 16 December 1986