
# appalachia



## Robert C. Leberman

Powdermill Nature Reserve Carnegie Museum of Natural History HC 64 Box 453 Rector, Pennsylvania 15677-9605 rcleberm@charterpa.net

By Appalachian standards it was a mild winter, with a warmer-than-usual December, a few weeks of cold, old-fashioned winter in late January, and spring-like conditions by the last week of February. At Erie, Presque Isle Bay remained open until mid-January and was free of ice again by 25 February, allowing a wide variety of waterfowl to linger there. Open water at many inland rivers, lakes, and reservoirs also provided habitat for a variety of wintering waterbirds. Many raptors, including Northern Harrier, Red-shouldered Hawk, and American Kestrel, likely benefited from low snow accumulations in the north that made small-mammal prey more accessible. No doubt because of these same mild weather conditions, many backyard feeder-watchers viewed the season as "dull" or, at least, "less than spectacular."

A scattering of hummingbirds attempted to winter in the southern part of the Region. As Kevin Calhoon noted, it was undoubtedly the first time ever that three different hummingbird species wintered at a single feeding station in Tennessee—or, this editor would add, anywhere else in the Appalachians! Northern Shrikes wintered in record-high numbers in western Pennsylvania, where a survey by Paul Hess account-

ed for a minimum of 68 birds scattered over 26 counties. Only rather moderate numbers of northern finches invaded the area, but, as noted below, the flight did take several species deep into the southern mountains. During the last week of February, with warm spring-like conditions, there were strong northward flights of ducks, Turkey Vultures, American Woodcocks, Eastern Phoebes, and Fox Sparrows throughout the Region.

**Abbreviations:** M.L.W.A. (Mosquito Lake Wildlife Area, Trumbull, OH); P.N.R. (Powdermill Nature Reserve, Rector, Westmoreland, PA); P.I.S.P. (Presque Isle State Park, Erie, PA).

#### **LOONS THROUGH VULTURES**

Fourteen Red-throated Loons were listed during a systematic waterbird count at Sunset Pt., P.I.S.P., through 1 Jan (JM); the only report s. of L. Erie came from Meander Cr. Res., Mahoning, OH, 18 Dec (fide CBa). Common Loons, recorded mostly as singles, were found nearly throughout the Region where there was suitable open water, with a high count of 28 at Douglas L., TN, 12 Dec (RKn, DE). The 46 Pied-billed Grebes listed on the Garrett, MD, CBC, 20 Dec, represented a new high count for that area (FP). Six hundred Horned Grebes were counted just offshore along a 5-mi stretch of beach at P.I.S.P. 18 Dec (JM); a high count of 76 was at Douglas L., TN, 12 Dec (RKn, DE), and seven were at L. Julian, Buncombe, NC, 20 Dec (WF). The season's only Red-necked Grebe was found off P.I.S.P. 7 Dec (JM). Now found regularly in the Region, up to six Eared Grebes wintered on S. Holston L., TN (RKn, WC), and as many as four were at P.I.S.P. 11-19 Dec (JD, LR, JM). A very late American White Pelican was an unusual find at Booker T. Washington S.P., TN, 18 Dec (JCh, BJ, HW); another was observed in Mahoning, OH, 19-21 Dec (fide DJH). Ten Doublecrested Cormorants were flying down the Allegheny R. at Warren, PA, as late as 6 Jan (DW), and about 15 stayed to winter at Kingsport, TN (RKn).

An Am. Bittern was a rare winter find near Brevard, NC, 22 Dec (NS); even more surprising was one near Oakland, MD, 13 Jan (fide FP). Six Great Egrets, first seen 12 Dec at Hiwassee W.R., TN (LW, BJ, JCh), remained through the season. A Green Heron flushed from the edge of Keystone L. during the Bushy Run CBC, Westmoreland, PA, 26 Dec (fide DBy) was certainly late for

that locality, but was the bird at Herrington Manor S.P., MD, 26 Feb (FP) wintering or an exceptionally early spring migrant? The only reports of wintering Black-crowned Night-Herons were of ≤10 at Kingsport, TN (RKn). In recent years there has been a trend for Turkey Vultures to attempt wintering farther n. in the Region; one at Erie 6 Jan (BS) represented the first winter record for that area, one was in *Indiana*, PA, 27 Dec (RMH), and two were in *Trumbull*, OH, 9 Jan (DJH).

#### WATERFOWL

Reports of a Ross's Goose from three Westmoreland/Indiana, PA, sites 26 Dec-28 Feb may all have been of the same individual (fide PH), but another was at Blythe Ferry W.M.A., Meigs, TN, 6 Feb (fide KC). At Pymatuning L., PA, Canada Geese peaked at over 11,200 birds 19 Dec (RH); 3000 were at M.L.W.A., OH, 30 Jan (DJH). An unmarked Trumpeter Swan, first seen at L. Arthur, PA, in July, remained until mid-January freeze-up (PH); five wing-tagged adults were at P.I.S.P. 31 Dec-16 Jan (JM, JSt). Late southward flights of Tundra Swans included 300 on 18 Dec over Allegheny/Butler, PA, (PH) and 100 at Tidioute, PA, 31 Dec (JK); 150 wintered on Erie Bay (JM), and an early northbound flock of 30 was over Ligonier, PA, 23 Feb

Over 700 Am. Black Ducks were at M.L.W.A., 19 Dec (CBa), while high counts of Mallards included nearly 3200 at Pymatuning L. 1 Jan (RFL), and 2000 at Quemahoning L., Somerset, PA, 25 Dec (TM). Two Blue-winged Teal at P.I.S.P. 18 Dec were especially late so far north (AnT), and late N. Shovelers included five at Silver L., Dayton, VA, 26 Dec (JBa) and 19 at Pymatuning L. 11 Dec (RFL). A female King Eider 18 Dec (†JM) was one of the highlights of the P.I.S.P. waterbird count. In Tennessee an imm. male Harlequin Duck on the S. Holston R. 1 Jan-early Feb (AH, JBr, m.ob.) represented just the 3rd state record. In w. Pennsylvania, away from L. Erie (where they are regular), single Surf Scoters were on the Ohio R. at Montgomery Dam, Beaver, 18 Dec (KB, BH), Kahle L., Venango, 24 Dec (JSt), and on the Alleheny R. near Pittsburgh 31 Jan (PH, SK); two were at L. Julian, NC, 20 Dec (WF, DBI) and two at Hiwassee W.R., TN, 12 Dec (RHo, DM). Two Black Scoters were at P.I.S.P. 8 Dec, with one there on the 28th (JM); two at Hiwassee 12 Dec were a rare find (RHo, DM). Exceptional Ruddy Duck counts included 1707 at L. Arthur, PA, 18 Dec (GW) and 944 in Garrett, MD, 18 Dec (FP).



The thousands of Common Redpolls that pushed southward during fall 1999 and continued into winter 1999/2000 were accompanied by a few Hoary Redpolls, such as this female at Presque Isle State Park, Pennsylvania, 10 Feb 2000. Photograph by Jerry McWilliams.

#### **RAPTORS THROUGH TERMS**

A late Osprey was at Cherokee L., TN, 12 Dec (RKn, DE), but one along the Allegheny R., at Warren, PA, 18 Feb (TG) was likely an exceptionally early spring migrant. Wintering N. Goshawks apparently were mostly confined to nw. Pennsylvania, but one was seen as far s. as Green Ridge State Forest, MD, 20 Feb (JBC). Rough-legged Hawks invaded the Region in the best numbers in several years: over 80 were reported, including 11 in nw. Warren, PA, 19 Dec (DW, TG), ten at Belleville, Mifflin, PA, 16 Jan (GG), and six in Allegany, MD, in mid-winter (GB, Rki). In Tennessee, reports came from Washington 5 Feb, Johnson 8-26 Feb (fide RKn), and Bledsoe 12 Dec and 10 Jan (fide KC). Wintering Golden Eagles included one at Acme, Westmoreland, PA, 6 Feb (JG), and another in Cedar Cr. Valley, Shenandoah, VA, 26 Dec (DDv). At least ten Merlins were reported from w. PA, with scattered birds seen elsewhere in the Region. The raptor of the season was a white-morph Gyrfalcon at the Stone Mt. hawkwatch, Huntingdon, PA, 7 Dec (†DBr). Single Peregrine Falcons were at P.I.S.P. 3 Jan (LM), Parkersburg, WV, in mid-Jan (fide JE), and Watauga L., TN, 9 Jan (HPL). Up to five wintered in Pittsburgh (PH) and a pair had returned to their nest site at Chickamauga Dam, TN, by 10 Jan (HS).

Early in the season, large rafts of Am. Coots settled on Regional lakes, including 3500 on Conneaut L., PA, 18 Dec (RCL, RFL), 2191 on L. Arthur 19 Dec (GW), and 475 at L. Moomaw, *Bath*, VA, 1 Jan (YL). On Erie Bay many coots did not survive to the end of February, when about three

dozen were found dead on the ice surrounding open patches of water (JM). A record count of 10,891 Sandhill Cranes was made at Hiwassee W.R., TN, 12 Jan (fide KC); one in Mercer, PA, 1 Jan was unexpectedly late so far north (NT). A record-late Lesser Yellowlegs was at Pymatuning L., Ashtabula, OH, 19 Dec (RCL, RFL); an early northbound migrant had reached Brainerd Levee, TN, 25 Feb (JHe). Among other interesting winter shorebirds were 31 Least Sandpipers and a Dunlin at Candies Cr. W.M.A., Bradley, TN, 10 Jan (KC). The only Purple Sandpiper was at P.I.S.P. 4–11 Dec (JHo, LC).

Single imm. dark-morph Pomarine Jaegers were spotted off P.I.S.P. 2 & 28 Dec and 2 Jan, with a probable fourth bird 30 Dec; a probable Parasitic Jaeger was there 30 Dec, all exciting results of McWilliams' systematic waterbird counts. An imm. Little Gull at Pymatuning L., 19 Dec represented the first Crawford, PA, record for the species (RCL, RFL), and one was at P.I.S.P. 8 Jan (JM). High counts of Bonaparte's Gulls included 608 at M.L.W.A 19 Dec (CBa), and 750 at Douglas L., TN, 12 Dec (RKn, DE). A Herring Gull near Brevard, NC, 27 Feb was just the 3rd for Transylvania (NS). The only Iceland Gull report was from P.I.S.P. 11 Feb (JM). A Lesser Black-backed Gull along the Monongahela and Ohio Rivers, Allegheny, PA, 28 Dec was an exciting find (ph. WDS), as were two at P.I.S.P. 30 Jan (JM). Five Glaucous Gulls were listed at Erie 31 Dec to 4 Feb (JM, DS). Single Great Black-backed Gulls at Pittsburgh 1 Jan (PH) and 6 Feb (JHo) were rare finds for that area, as was an imm. Black-legged Kittiwake along the Ohio R. at Montgomery Dam 14 Dec (SK). A Com. Tern was still at P.I.S.P. 10 Dec (JM), and a remarkably late Forster's Tern was at Nickajack L., Marion, TN, 27 Dec (JHe).

#### DOVES THROUGH HUMMINGBIRDS

The Eur. Collared-Dove population continued to increase in Hamilton, TN, with a high count of 61 on 30 Jan (DG); five were at Pikeville, a new location in Bledsoe, TN, 5 Feb (DV). Barn Owls, seldom reported in recent years, were located in at least three sites in ne. Tennessee (RKn). The lone report of Snowy Owl came from P.I.S.P. 29-30 Dec (JM). Long-eared Owls were more widely reported than usual in w. Pennsylvania, led by a total of 23 at three separate roosts in Juniata, PA (AdT), 9 birds in Somerset, PA (SB, DDa), and 4 in Venango (JSt); one or two in Shady Valley, TN, 22-26 Jan, represented the 3rd local record (RKn). Short-eared Owls were in good numbers in, ne. Ohio, Pennsylvania, and w. Maryland, with more than 70 reported; ≤3 were regular at Hooper Lane, Henderson, NC, after 26 Dec (WF) Northerly Reports of N. Saw-whet Owls came from 13 sites in w. Pennsylvania, with ≤7 at Moraine, S.P. (DDa). Southward, ten were found in Big S. Fork National Recreation Area, TN/KY (SJS); singles were at Bristol 26 Dec, and Roan Mt., TN, 11 Feb (fide RKn); and ≤3 were near Clayton, Rabun, GA, 8 Jan–3 Mar (JSe).

Incredibly, three species of hummingbirds wintered at a feeding station operated by Gil Brewster at East Brainerd, Hamilton, TN, and all were captured and banded (CSI, PM). An imm. male Ruby-throated Hummingbird trapped there 12 Dec was seen occasionally thereafter through the end of the period; an ad. female Rufous Hummingbird, also captured 12 Dec, was not seen subsequently; and an imm. male **Black-chinned Hummingbird** banded 4 Dec remained through Feb (fide KC). Elsewhere, an imm. male Rufous Hummingbird banded 4 Dec (CS, PM) at Signal Mt., TN, stayed throughout the period; a banded female was in Sullivan, TN, into Feb (fide WC). At Brevard, NC, two Selasphorus hummingbirds frequented a feeder mid-Oct-3 Jan, when one disappeared; the second bird was banded as a Rufous 5 Jan (SC) and remained through Feb (fide NS).

## SHRIKES THROUGH FINCHES

The only reports of Loggerhead Shrike were of a few scattered birds wintering in ne Tennessee (RKn). Northern Shrikes, however, staged their greatest invasion on record. In ne. Ohio, five were seen in Trumbull (DJH) and one in Ashtabula (CBa), while in w. Pennsylvania a fantastic 68 shrikes were reported from 26 counties (fide PH, RCL). At least two Northerns crossed the Mason-Dixon line into Maryland, with one seen during the Garrett CBC (FP, CSk) and another at Dan's Mt, Allegany, 31 Dec-4 Jan (RKi). Near Clintonville, PA, Jerry Stanley witnessed a protracted chase and the eventual capture of a bat by a Northern Shrike on 24 Feb!

Blue-headed Vireos wintered in good numbers in the s. third of the Region, and one was listed as far north as Pipestem, WV, 16 Jan (JPh). Two early Tree Swallows had reached Staunton, VA, by 29 Feb (YL). In mid-Dec two Black-capped Chickadees were banded in *Sullivan*, TN, where they are rare, with one remaining into March (RL) Following last fall's heavy southward chickadee migration, significant northward movement could be detected at the P.N.R

178 NORTH AMERICAN BIRDS

banding station by early Feb (RCL, RM). Two Sedge Wrens and a Marsh Wren were good finds on the Hiwassee W.R. CBC 1 Jan, and a Marsh Wren was at Brainerd Levee, TN, 18 Dec (KC). A Townsend's Solitaire was briefly observed but welldescribed near Slickville, Westmoreland, 15 Jan—a first w. Pennsylvania winter report (†MM). American Robins were unusually abundant in the north; high counts included 17,500 at a Washington, PA, state game lands 29 Jan (KB, BH), over 10,000 at a roost in Fairview, PA, in early Feb (JM), and 5863 at Raccoon Cr. S.P., PA, 28 Dec (fide PH) An unusually high Am. Pipit count was of 374 birds on the Henderson, NC, CBC 19 Dec (WF). Orange-crowned Warblers, four in all, were found at three sites in Hamilton and Marion, TN, between 18 Dec and 31 Jan (fide KC). Several Pine Warblers were seen in Transylvania, NC, in February (NS), and one was in Carter, TN, 20 Jan (FA). In e. Tennessee, ten Palm Warblers were listed between 18 Dec-1 Jan (RKn, KC), and one was near Pisgah Forest, NC, 15 Jan (NS). A mid-winter report of Com Yellowthroat came from Carter, TN, 9 Jan (fide RKn).

A Spotted Towhee found at Joe Corsales' bird feeder at Export, Westmoreland, PA, during the Bushy Run CBC 19 Dec, represented just the 4th Pennsylvania record; it remained through the end of the period, allowing many birders to add this rarity to their lists (DBy, m.ob.). Eastern Towhees wintered in above-average numbers nearly throughout the Region, A Savannah Sparrow at Coneautville, PA, 22-26 Jan was a rare midwinter find so far north (fide JM) and 18 at Swoope, Augusta, VA, 13 Feb represented a new high winter count there (YL). A late Nelson's Sharp-tailed Sparrow was a rare find in Hawkins, TN, 18 Dec (RKn). Usually scarce in Pennsylvania, White-crowned Sparrows wintered in record numbers, with 91 on the Bushy Run CBC (DBy); a Gambel's White-crowned was at a feeder near Shelocta, Indiana, PA, 6-27 Dec (RMH). Snow Buntings were fairly common in the north, with ≤200 in late Jan at Neshannock Falls, Lawrence, PA (BD); the most southerly report was of two in Garrett, MD, 5 Feb (JBC). A male Rose-breasted Grosbeak visited a feeder at Little River, NC, 14 Feb (fide NS). A Brewer's Blackbird appeared at Amnicola Marsh, Hamilton, TN, 14 Jan (CBl); three were at Brainerd Levee 10 Feb (KC), with another on the 19th (JHe). In Pennsylvania, Baltimore Orioles briefly appeared at feeders in Harborcreek 15 Jan (SP), and Sayre 17 Jan (RS).

After a heavy flight of Purple Finches through Pennsylvania during the fall, they were unaccountably scarce at P.N.R. by early winter (RM, RCL), but they were numerous in Transylvania, NC (NS) and in "substantial numbers" in Pipestem, WV (JPh). Small numbers of Red Crossbills invaded the Region, reaching as far s. as Cades Cove, TN (SJS), and Henderson, NC (fide WF). Whitewinged Crossbills were only slightly more numerous, with a high count of ten in Franklin, PA (ISt) and a few wandering s. to Roan Mt. and Signal Mt., TN (fide KC, RKn), Flocks of 100 or more Com. Redpolls were occasionally observed, especially in Erie, Crawford, and Lawrence, PA, and single birds reached as far s. as Shady Valley, TN, 25 Jan-23 Feb (JSh) and Augusta, VA, 8-19 Feb (YL). A female **Hoary Redpoll** (ph. JM) was a rare find at P.I.S.P. 10 Feb. Following a fairly heavy autumn flight, Pine Siskins were uncommon or absent over much of sw. Pennsylvania, including P.N.R., by mid-winter (PH, RCL), but were plentiful in Transylvania, NC (NS), Augusta, VA (YL), and, locally, in se. Tennessee (KC). Evening Grosbeaks were rather widespread in the Region but nowhere common—the highest count was of just over 125 at Roan Mt. 15 Jan (fide RKn).

Contributors and cited observers: Fred Alson. Carole Babyak (CBa), Scott Bastian, John Bazuin (JBa), Ken Behrens, Ralph Bell, D Blee, Clyde Blum (CBI), Gwen Brewer, James Brooks (JBr), Nancy Brundage, Don Bryant (DBr), Dick Byers (DBy), Kevin Calhoon, Susan Campbell, Janice Chadwell (JCh), Louise Chambers, John B. Churchill (JBC), Wallace Coffey, Joe Corsales (JCo), Dave Darney (DDa), David Davis (DDv), Barbara Dean, Jon Dunn, Dean Edwards, Jeanette Esker, Wayne Forsythe, Danny Gaddy, Kathie Goodblood, Ted Grisez, Grea Grove, Jim Gruber, Ron Harrell (RH), John Henderson (JHe), Paul Hess, Roger and Margaret Higbee (RMH), Bill Hintze, Dave and Judy Hochadel (DJH), Audrey Hoff, Ron Hoff (RHo), Joyce Hoffmann (JHo), Bonnie Johnson, Ray Kiddy (RKi), Jim King, Scott Kinzey, Rick Knight (RKn), YuLee Larner, Portia MacMillen, Mark McConaughy, Linda McWilliams, Jerry McWilliams, H. P. Langridge (HPL), Ronald F. Leberman (RFL), Richard Lewis (RL), Tony Marich, Robert Mulvihill, Dollyann Myers, Jeff and Retta Payne. Jim Phillips (JPh), Fran Pope, Shirley Post, Larry Rosche, Jeff Sewell (JSe), Walt and Dana Shaffer (WDS), Harold Sharp, Bob Shaw (BSh), Richard Shelling, John Shumate (JSh), Norma Siebenheller, Connie Skipper (CSk), Chris Sloan (CSI), Michael Smith, Don Snyder, Jerry Stanley (JSt), Stephen J. Stedman, Bob Sunday, Aden Troyer (AdT), Andy Troyer (AnT), Neil Troyer, David Vogt, Don Watts, Gene Wilhelm, Havden Wilson, Libby Wolfe.

# BirdBase Software for a BirdArea

Nearly all the ABA birders reporting life lists over 6000 who use computers use our software. It is the software most widely used by all those over the 1400 bird threshold. Why? Because it lets them easily keep track of birds seen and also helps them see new ones with its ability to list the birds of any major world area, labeling all endemics and birds previously seen in the area, outside it, or both. View an on-screen demonstration at http://members.aol.com/sbsp

BirdBase can switch between displaying all the birds in James Clements' world list or any selection of them, when recording sightings. It accepts 5000-word sighting notes. And it instantly updates any number of annual/life lists (e.g., yard, county, state, nation, ABA region/area, world). These can give details of all sightings or of only each bird's first sighting, or just name birds seen. BirdBase can list all sightings of a bird, list in date-order all trips or first sightings of all birds, tabulate Christmas counts, etc. It handles all changes in the world list, and puts them in sightings already recorded.

**BirdArea** can list in detail the range of any bird. And it can produce check lists of the birds of all major world areas (i.e., U.S. states, Canadian provinces, nations, ABA regions/areas, important islands) with endemics labeled. Printouts allow multiple check marks and notes. Shawneen Finnegan's annually-updated ranges come from more than 800 publications in 10 languages and from many experts birding the areas.

If BirdBase is used BirdArea can label birds already seen on each check list, make lists of birds not already seen, and find any recorded sighting in which a bird is outside its known range. If BirdArea is used BirdBase can switch to displaying only birds whose ranges include any of the world areas when recording sightings of a trip to that area. This makes recording very much easier and calls to attention out-of-range sightings.

BirdBase is \$59.95 + \$4.00 shipping; so is BirdArea. Buy both for \$99.95 + \$6.00. Or start with BirdBase and N. American birds for \$39.95 + \$3.00. Add 8% for CA orders. Double overseas shipping. Visa, MasterCard OK. For Windows 3.X, 9X, NT.

#### Santa Barbara Software Products

1400 Dover, S.B. CA 93103 USA; Phone/fax: 805 963 4886; E-mail: sbsp@aol.com