

Individual variation in behavior among male Blue Grouse

Martin K. McNicholl

Ornithologists, and banders in particular, studying populations of birds in the wild are frequently impressed by individual variations in behavior, with certain individuals consistently acting in a manner different from other conspecifics of the same sex and age. As noted by Thomson (1964), such variability might be expected to be even greater than morphological variation, as behavior may be determined by genetics, experience, or both. Indeed, variation to an individual level is documented well for some behavioral components, such as call notes and song (e.g. Falls 1969; Beer 1970; Falls and McNicholl in press). Yet, few behavior studies stress such variability, except in relation to such practical problems as trapability (e.g. Doan 1976; Hamerstrom and Hamerstrom 1977), band removal affecting population estimates, longevity data and other information based on banding returns (Wiseman 1977), and persistent difficulty of feeding some individuals in captivity (Berry 1975). Studies, such as those by Kennard (1894), Lockley (1940), and Partridge (1976), emphasizing individual variation in behavior, are relatively few.

The current development of fast and efficient methods of analyzing large volumes of numerical data make the quantification of behavior both valuable and tempting. Yet behavior patterns are rarely so stereotyped as to allow ready tabulations without loss of qualitative information. Knowledge of individual variation, however, does allow quantification of behavior with considerable confidence. For example, 13 male Blue Grouse (*Dendragapus obscurus*) on Vancouver Island, British Columbia showed distinct differences in responses to playbacks of taped hooting (the song of Blue Grouse), at the boundaries of their territories in 1973 (Tables 1 and 2). Nevertheless, by combining several behavioral parameters, J.B. Falls and I were able to demonstrate that each individual showed less response to the songs of neighbors from the correct direction than to songs of other individuals (Falls and McNicholl in press). All but one of these birds were color-banded, enabling us to confirm individual identity.



Several recent investigators have attempted to test for genetic differences in Blue Grouse at a population level by comparing behavior (e.g. Mossop 1971; Willie 1971; Hemus 1972; Donaldson 1973; Low 1974; Bergerud and Hemus 1975; Cooper 1977). With the exception of Cooper's study, these investigations were based on such behavioral parameters as reaction to an "arena" consisting of three mirrors, a female squat dummy and female calls, and reactions to observers. Although these researchers did indeed find differences in these parameters, indicating possible differences in populations, their results are inconclusive without background data on individual variation within the population, and variability in behavior of each individual. If individual behavior is marked and consistent within each bird, their results likely reflect genetic differences, but if behavioral variation varies markedly within each individual, their results may reflect only chance reactions of each bird on a particular encounter. During detailed studies of color-banded males from 1971 to 1974, I did find marked individual differences in reactions to me (McNicholl 1978a), "tameness" (McNicholl 1978a: table 3), individual hooting patterns as to number of syllables per song and frequency of

Table 1. Postural responses of male Blue Grouse to taped hooting at edge of territory.

Experimental subject	Responses to two experiments, each including 3 playbacks ¹
1	Remained in neutral posture throughout first experiment; in 2nd, no display on 1st N, partial display on both S.
2	Remained in neutral posture throughout 1st experiment; in 2nd, showed aggressive reactions on both S, no display on N.
3	In 1st experiment, white around lateral apertia on S and 1st N, no display on 2nd N; aggressive reactions to both S and no display to N of 2nd experiment.
4	No display on either N of 1st experiment; full display on S. In 2nd experiment, no display on N; full display on 1st S; aggressive on 2nd S.
5	In 1st experiment, no display on either N, full display on S; in 2nd experiment, no display on N, full display on 1st S, aggressive on 2nd S.
6	In 1st experiment, no display on either N, partial display on S; in 2nd experiment, no display on N, full display on 1st S, aggressive on 2nd S.
7	In 1st experiment, no display on either N, partial display on S; in 2nd experiment, no display on N, full display on both S.
8	In 1st experiment, no display on either N; on S, partial display, in 2nd experiment, no display on N, full display on both S playbacks.
9	No display on 1st N in 1st experiment, full display on S, and full display to J.B.F. on 2nd N; in 2nd experiment, no display on N, full display on both S, with aggressive reaction on both S.
10	No display on either N in 1st experiment, partial display on S. In 2nd experiment, full display on both S, no display on N.
11	Full display on both N of 1st experiment, aggressive reaction on S; in 2nd experiment, full display on 1st S, no display on N, partial display on 2nd S.
12	In 1st experiment, no display on either N, partial display on S; in 2nd experiment, no display on N, partial display on 1st S, full display on 2nd S.
13	In 1st experiment, no display on either N, full display on S; in 2nd experiment, no display on N, full display on both playbacks.

¹1st experiment: playback of correct neighbor (N), then stranger (S), then same N; 2nd experiment, playback of different S, then N, then S. Order of presentation and other details in Falls and McNicholl, in press.

singing (McNicholl 1978b: table 8), and individual tendency to sing (McNicholl 1978b: table 7). Although most birds showed occasional atypical responses, behavioral differences were generally consistent within individuals. Such results suggest that differences found by the several workers in the studies cited above reflected true behavioral differences among birds of different populations; a remote chance still remains that the researchers encountered atypical or individualistic responses in the birds studied. This possibility is also in-

Table 2. Movements¹ of male Blue Grouse in response to taped hooting at edge of territory.

Experimental subject	N-S-N ²			S-N-S ²		
	N	S	N	S	N	S
1	0	1/119	0	0	0	0/(119)
2	0	0	0/(66)	125/140	0	15/15
3	4/55	8/51	0	64/64	0	0
4	0.3/115	4/115	-4/115 (moved back)	115/115 (but circled to 15 away)	0	15/15
5	0	2.5/82	0/79.5	62/82	0	20/20(flew)
6	0	12/50	0/38	52/91	0	37/39+5 (went behind speaker)
7	0	4/56	-4/56 (moved back)	12/52	0	26/40
8	0	59/117	0/58	30/64	0	27/34
9	0/33	46/33 (behind speaker)	0/13	59/59	0	0, but circled speaker
10	0	6/85	0/79	32/93 (flew)	0	53/61
11	55/55 (flew)	0 (circled speaker)	0 (circled speaker)	15/270	0	45/255
12	0	2/40	0/38	8/35	0	25/27
13	0	57/60	0/3	23/50	0	20/28

¹Distance moved in M.K.M. paces/remaining distance between bird and speaker in M.K.M. paces. All movements on foot unless otherwise noted.

²N-S-N and S-N-S experiments as per Table 1 footnote.

dicated by the work of Cooper (1977), who found marked individual variation in behavior of particular grouse in cages.

Thus, behavior can be compared at a population level, but the results are difficult to interpret without both knowledge of the amount of behavioral variation within each population, and the consistency of behavior of the individual birds studied. Banders and others studying wild populations of birds can provide welcome additional information on this topic through careful and detailed observations of individually marked birds.

Acknowledgements

I am grateful to John H. Alway, Fred C. Zwickel and an anonymous referee for comments on an earlier version of the manuscript. These studies were supported by the National Research Council of Canada, an Isaac Killam Fellowship, the British Columbia Fish and Wildlife Branch, and the University of Alberta. Research was conducted on lands of Crown Zellerbach Canada Ltd., Courtenay Division.

Literature cited

- Beer, C.G. 1970. Individual recognition by voice in the social behavior of birds. *Adv. Study Behav.* 3:27-74.
- Bergerud, A.T. and D.H. Hemus. 1975. An experimental study of the behavior of Blue Grouse (*Dendragapus obscurus*). I. Differences between the founders from three populations. *Can. J. Zool.* 53:1222-1237.
- Berry, H.H. 1975. Hand-rearing Lesser Flamingos. pp. 109-116 in Kear, J. and N. Duplaix-Hall (Editors). *Flamingos. T. & A.D. Poyse, Berkhamsted.*
- Cooper, C.R. 1977. Differences in behavior between populations of captive Blue Grouse. M. Sc. thesis, Univ. Toronto, Toronto.
- Doan, L.A. 1976. Behavior of Black Terns on the nesting grounds. *Inland Bird Banding News* 48:49-54.
- Donaldson, J.L. 1973. A behavioral and demographic study of an insular population of Blue Grouse. B. Sc.thesis, Univ. Victoria, Victoria.
- Falls, J.B. 1969. Functions of territorial song in the White-throated Sparrow. pp. 207-232 in Hinde, R.A. (Editor). *Bird vocalizations.* Cambridge Univ. Press, Cambridge.
- Falls, J.B. and M.K. McNicholl. In press. Neighbor-stranger discrimination by song in male Blue Grouse. *Can J. Zool.*
- Hamerstrom, F. and F. Hamerstrom. 1977. On the art of Harris' Hawk trapping. *Inland Bird Banding News* 49:4-8.
- Hemus, H.D. 1972. A behavioral and demographic comparison of three Blue Grouse populations released on Moresby Island. B.C. B.Sc. (Hons) thesis, Univ. Victoria, Victoria.
- Kennard, F.H. 1894. The habits and individualities of the Red-shouldered Hawk (*Buteo lineatus*) in the vicinity of Brookline, Mass. *Auk* 9:197-210.
- Lockley, R.M. 1940. The seabird as an individual. Results of ringing experiments. Smithsonian rept. for 1939:341-352.
- Low, D.J. 1974. Qualities of individuals in a replacement stock of Blue Grouse. M. Sc.thesis, Univ. British Columbia, Vancouver.
- McNicholl, M.K. 1978a. Reactions of male Blue Grouse to intrusions by observers. Appendix 2 in McNicholl, M.K. *Behavior and social organization in a population of Blue Grouse on Vancouver Island.* PhD. thesis, Univ. Alberta, Edmonton.
- _____ 1978b. Vocal communication among male Blue Grouse. Paper 3 in McNicholl, M.K. *Behavior and social organization in a population of Blue Grouse on Vancouver Island.* PhD. thesis, Univ. Alberta, Edmonton.
- Mossop, D.H. 1971. A relation between aggressive behavior and population dynamics in Blue Grouse. M. Sc. thesis, Univ. British Columbia, Vancouver.
- Partridge, L. 1976. Individual differences in feeding efficiencies and feeding preferences of captive Great Tits. *Animal Behav.* 24:230-240.
- Thomson, A.L. 1964. *A new dictionary of birds.* McGraw-Hill, New York.
- Willie, F.B. 1971. Behavior as a possible regulatory mechanism in three Blue Grouse populations on lower Vancouver Island. B. Sc. thesis, Univ. Victoria, Victoria.
- Wiseman, A.J. 1977. Band removal by Cardinals. *Inland Bird Banding News* 49:33-37. ♠

Department of Zoology, University of Alberta, Edmonton, Alberta, T6G 2E9 Canada. Present address: Beak Consultants Ltd., 3530 11 A St. N.E., Calgary T2E 6M7, Alberta, Canada.



H. Cox Jones

Corrections to MTAB 36

The BBL has informed us of two errors in the recent MTAB:

5. "Page 26" should read "Page 2—7."
Delete new band sizes for Broad-winged Hawk. 5—6 are the correct sizes.