

# **ORNITOLOGIA NEOTROPICAL**

EDITOR

RAYMOND McNEIL

## **INDEX TO VOLUME 16, 2005**

(Authors, key words, families, common names, latin names)

An International Journal of Neotropical Ornithology

published by

**THE NEOTROPICAL ORNITHOLOGICAL SOCIETY**

- Abourachid, A., 471–479  
 Abundance, 219–230  
*Aburria aburri*, 205–217  
*Accipiter striatus*, 85–99, 109–115, 219–230  
*Adelomyia melanogenys*, 297–319  
 Advertisement flights, 197–204  
*Aegialaius thilius*, 551–555  
 Aging, 481–491  
 Agreda, A., 297–319  
 Ahumada, G. A., 53–63  
*Aimophila mystacalis*, 101–104  
 Alava, J. J., 175–185  
*Alcedo atthis*, 141  
 Almonte, J., 219–230  
*Alopochelidon fucata*, 85–99  
 Altitudinal ranges, 435–436  
 Alvarado Orellana, S., 271–275  
 Alvarez, M. E., 231–251  
*Amazilia amazilia*, 297–319  
*Amazilia chionogaster*, 231–251  
*Amazilia fimbriata*, 1–14, 85–99  
*Amazilia franciae*, 129–136  
*Amazilia tobaci*, 1–14  
*Amazilia tzacatl*, 129–136, 297–319  
*Amazona aestiva*, 231–251, 573–578  
*Amazona amazonica*, 85–99  
*Amazona ochrocephala*, 1–14  
*Amazona tucumana*, 43–52  
*Amazona ventralis*, 405–417  
*Amazona vittata*, 405–417  
 Amazonia, 289–296  
*Amblycercus holosericeus*, 297–319  
*Ammodramus humeralis*, 85–99  
*Anabacerthia variegaticeps*, 297–319  
*Anairetes fernandezianus*, 371–385  
 Andean premontane forest, 231–251  
 Andes, 53–63, 137–140, 205–217, 231–251, 283–286, 427–432, 439–440, 511–518, 567–571  
*Anhinga anhinga*, 85–99  
*Anthracothorax dominicus*, 219–230  
*Anthracothorax*, 1–14  
*Apbrastura masafuerae*, 371–385  
*Ara ambiguus*, 297–319  
*Ara auricollis*, 85–99  
*Ara chloroptera*, 85–99  
*Ara manilata*, 289–296  
 Aragon Tapia, A., 279–281  
 Aramburu, R., 573–578  
*Aramides cajanea*, 85–99  
*Aramus guarauna*, 85–99  
*Aratinga erythrogenys*, 297–319  
*Aratinga leucophthalma*, 43–52, 231–251  
*Aratinga mitrata*, 43–52  
*Aratinga wagleri*, 1–14  
*Ardea cocoi*, 85–99  
 Argentina, 31–42, 43–52, 117–121, 145–161, 231–251, 267–270, 435–436, 503–510, 519–527, 529–537, 539–546, 563–566  
 Arid zones, 101–104  
*Arremon abeillei*, 297–319  
*Arremon aurantiirostris*, 129–136, 297–319  
 Arthropods, 529–537  
*Asio flammeus*, 277–278, 371–385  
*Asturina nitida*, 231–251  
*Athene cunicularia*, 529–537  
 Atlantic forest, 65–83, 197–204, 547–549  
*Atlapetes tricolor*, 129–136  
*Attila spadiceus*, 1–14  
*Attila torridus*, 297–319  
*Aulacorhynchus haematopygus*, 129–136, 297–319  
 Austral summer grounds, 267–270  
*Automolus leucophthalmus*, 65–83  
*Automolus ochrolaemus*, 297–319  
 Avifauna, 1–14, 85–99, 219–230  
 Azpiroz, A. B., 105–108  
 Barro Colorado Island, 187–195  
*Baryphthengus martii*, 129–136  
*Basileuterus culicivorus*, 85–99  
*Basileuterus flaveolus*, 65–83  
*Basileuterus fraseri*, 297–319  
*Basileuterus hypoleucus*, 65–83  
*Basileuterus tristriatus*, 129–136  
*Batara cinerea*, 231–251  
 Bechard, M. J., 267–270  
 Becker, C. D., 297–319  
 Behavior, 105–108, 137–140, 197–204, 271–275  
 Behavioral evolution, 15–29  
 Berkunsky, I., 573–578  
 Biogeography, 371–385  
 Biondi, L. M., 31–42  
 Bird Bandung, 145–161  
 Bird communities, 337–346  
 Bird nest predation, 187–195  
 Birds of prey, 197–204  
 Blendinger, P. G., 231–251  
 Bó, M. S., 31–42

- Bolivia, 85–99, 109–115, 337–346  
*Branco, L. C.*, 529–537  
*Brazil*, 65–83, 547–549  
 Breeding association, 117–121  
 Breeding organization, 15–29  
 Breeding period, 31–42  
 Breeding, 117–121, 123–126  
 Brood parasitism, 435–436  
*Brooks, D. M.*, 85–99., 289–296  
*Brotogeris chiriri*, 85–99  
*Brotogeris versicolorus*, 289–296  
*Buarremon brunneinucha*, 297–319  
*Bubo magellanicus*, 539–546  
*Bubulus ibis*, 85–99, 129–136  
*Bücker, A. D. L.*, 283–286  
*Buteo albigena*, 271–275  
*Buteo brachyurus*, 129–136  
*Buteo jamaicensis*, 85–99, 219–230  
*Buteo magnirostris*, 1–14 , 231–251  
*Buteo nitidus*, 297–319  
*Buteo plathypterus*, 109–115  
*Buteo polyosoma*, 271–275, 371–385  
*Buteo swainsoni*, 109–115, 267–270  
*Buteogallus meridionalis*, 109–115  
*Buteogallus urubitinga*, 85–99, 109–115, 297–319  
*Cabanne, G. S.*, 197–204, 547–549  
 Cacao plantation, 1–14  
*Cacicus cela*, 1–14, 289–296  
*Calyptophilus tertius*, 219–230  
*Campephilus gayaquilensis*, 129–136, 297–319  
*Campephilus leucopogon*, 231–251  
*Campephilus melanoleucus*, 231–251  
*Camptostoma obsoletum*, 85–99, 129–136, 297–319  
*Campylorhynchus trochilirostris*, 297–319  
*Campylorhynchus turdinus*, 85–99  
*Caplonch, P.*, 145–161  
*Caprimulgus anthonyi*, 297–319  
*Capsiempis flaveola*, 297–319  
 Captive-reared, 405–417  
*Cariama cristata*, 85–99, 471–479  
*Cariamidae*, 471–479  
 Caribbean, 405–417  
*Casiornis rufa*, 85–99, 231–251  
*Casmerodium alba*, 85–99  
*Castillo Meza, A. L.*, 101–104  
*Cathartes aura*, 85–99, 109–115, 129–136, 231–251  
*Cathartes burrovianus*, 85–99  
*Catharus bicknelli*, 219–230  
*Catharus dryas*, 231–251, 297–319  
*Catharus ustulatus*, 231–251, 297–319  
*Celeus lugubris*, 85–99  
 Census methods, 337–346  
 Census, 175–185  
*Cercococcyx tyrannina*, 129–136  
*Cerrado*, 85–99  
*Certhiaxis cinnamomea*, 85–99  
*Ceryle torquata*, 85–99  
*Chaco*, 145–161  
*Chaetocercus bombus*, 297–319  
*Chaetura cinereiventris*, 297–319  
*Chaetura meridionalis*, 231–251  
*Chaetura ocyptates*, 297–319  
*Chalbura buffoni*, 1–14  
*Charadrius collaris*, 117–121  
 Chewing lice, 573–578  
*Chile*, 271–275, 371–385  
*Chiroxiphia lanceolata*, 1–14  
*Chituitano*, 85–99  
*Chlorestes notatus*, 1–14  
*Chloroceryle amazona*, 85–99, 289–296  
*Chloroceryle americana*, 1–14, 85–99  
*Chloroceryle inda*, 85–99  
*Chlorospingus canigularis*, 129–136, 297–319  
*Chlorospingus parvirostris*, 437–438  
*Chlorostilbon aureoventris*, 85–99, 231–251  
*Chlorostilbon melanorhynchus*, 129–136, 297–319  
*Chlorostilbon swainsonii*, 219–230  
*Chondrohierax uncinatus*, 109–115, 231–251, 297–319  
*Chrysuronia oenone*, 1–14  
*Chunga burmeisteri*, 471–479  
*Cinclodes oustaleti*, 371–385  
*Cinclus leucocephalus*, 129–136  
*Circus buffoni*, 109–115  
*Cistothorus palustris*, 277–278  
 Cloud forest, 137–140, 283–286, 439–440  
*Cnemotriccus fuscatus*, 1–14, 85–99, 231–251  
*Coccyzus americanus*, 85–99  
*Coccyzus melacoryphus*, 231–251  
*Coereba flaveola*, 1–14, 85–99, 219–230, 297–319  
 Cold fronts, 109–115  
*Collazo, J. A.*, 405–417  
*Colombia*, 53–63, 205–217  
*Columba fasciata*, 129–136  
*Columba livia*, 371–385

- Columba picazuru*, 85–99  
*Columbina picui*, 85–99  
*Columbina squammata*, 85–99  
*Columbina talpacoti*, 85–99  
 Community, 1–14, 231–251, 337–346, 371–385, 503–510  
*Conirostrum speciosum*, 85–99  
*Conopophaga lineata*, 65–83  
 Conservation, 1–14, 43–52, 85–99, 175–185, 183–174, 219–230, 297–319, 337–346, 371–385, 419–426, 503–510  
*Contopus cooperi*, 297–319  
*Contopus fumigatus*, 297–319  
*Contopus punensis*, 297–319  
 Convergence, 387–396  
 Cooperative searching, 271–275  
 Corado, R., 277–278  
*Coragyps atratus*, 85–99, 109–115, 129–136, 231–251, 297–319  
*Coryphospingus cucullatus*, 85–99  
*Cotinga maynana*  
 Cowbird host, 435–436  
 Cracids, 205–217  
*Cranioleuca erythrops*, 297–319  
*Cranioleuca pyrrhophia*, 231–251  
*Crax fasciolata*, 85–99  
*Crotophaga ani*, 85–99  
*Crypturellus obsoletus*, 471–479  
*Crypturellus parvirostris*, 85–99  
*Crypturellus sonori*, 129–136, 297–319  
*Crypturellus tataupa*, 231–251  
*Crypturellus undulatus*, 85–99  
 Cuba, 141  
*Cyanerpes caeruleus*, 129–136  
*Cyanocompsa cyanoides*, 1–14, 297–319  
*Cyanocorax chrysops*, 85–99, 231–251  
*Cyanocorax cyanomelas*, 85–99  
*Cyanocorax yncas*, 1–14  
*Cyclarhis gujanensis*, 231–251, 297–319  
*Cyphorhinus phaeocephalus*, 297–319  
*Dacnis egregia*, 129–136  
 Daily variation, 337–346  
*Damophila julie*, 297–319  
 Decades, 321–336  
*Dendrocincla fuliginosa*, 1–14, 129–136, 297–319  
*Dendrocolaptes certhia*, 297–319  
*Dendrocolaptes picumnus*, 231–251  
*Dendroica caerulea*, 219–230  
*Dendroica tigrina*, 219–230  
*Dendrocygna leucophrys*, 557–561  
 Density, 205–217  
 Detectability, 337–346  
 Diet, 31–42, 65–83, 519–527  
 Diet overlap, 539–546  
 Display, 15–29  
 Distribution, 85–99, 175–185, 361–370, 371–385, 529–537  
 Diversity, 129–136, 219–230  
*Dives warzenszczii*, 129–136  
 Dobbs, R. C., 427–432, 511–518  
 Dominican Republic, 405–417  
*Donacobius atricapillus*, 85–99  
 Doucet, S. M., 433–434  
*Dryocopus lineatus*, 1–14, 297–319  
 Duetting, 397–404  
 Duråes, R., 65–83, 481–491  
*Dysithamnus mentalis*, 65–83, 297–319  
 Ecology, 183–174  
 Ecotone, 539–546  
 Ectoparasites, 573–578  
 Ecuador, 129–136, 137–140, 175–185, 297–319, 321–336, 427–432, 437–438, 511–518, 567–571  
 Ecuadorian authors, 321–336  
 Ecuadorian ornithology, 321–336  
 Eggs, 101–104, 137–140, 427–432, 437–438, 493–501, 557–561, 567–571  
*Egretta thula*, 289–296  
 Eitniear, J. C., 279–281  
 El Salvador, 557–561  
*Elaenia albiceps*, 145–161  
*Elaenia fallax*, 219–230  
*Elaenia flavogaster*, 85–99  
*Elaenia obscura*, 231–251  
*Elaenia parvirostris*, 145–161  
*Elaenia strepera*, 145–161  
*Elanoides forficatus*, 109–115, 231–251, 297–319  
*Elanus leucurus*, 109–115  
 Elevational gradient, 43–52  
*Empidonax virescens*, 297–319  
*Empidonax varius*, 85–99, 231–251  
 Endemic birds, 219–230, 297–319  
 Endemism, 371–385  
 Environmental stability, 289–296  
 Ericson, P. G. P., 347–359  
*Eubucco bourcierii*, 297–319  
*Eucometis penicillata*, 1–14

- Euphonia chlorotica*, 85–99  
*Euphonia laniirostris*, 1–14, 297–319  
*Euphonia musica*, 219–230  
*Euphonia xanthogaster*, 1–14, 129–136, 297–319  
*Euryptila helias*, 85–99  
*Euscarthmus meloryphus*, 297–319  
*Eutoxeres aquila*, 129–136  
 Evolution, 347–359  
*Falco peregrinus*, 85–99, 109–115  
*Falco rufipectoralis*, 85–99, 297–319  
*Falco sparverius*, 109–115, 219–230, 371–385  
 Falconiformes, 109–115  
 Favero, M., 31–42  
 Fedy, B. C., 397–404  
 Feeding guilds, 231–251  
 Feeding rates, 283–286  
 Fernández, E. M., 219–230  
 Figueroa Rojas, R. A., 271–275  
 Fjeldså, J., 347–359  
 Flock size, 43–52  
*Florisuga mellivora*, 1–14  
*Fluvicola nengeta*, 129–136  
 Food abundance, 529–537  
 Foraging ecology, 427–432, 563–566  
 Foraging, 397–404  
 Forest conservation, 297–319  
 Forestry concession, 337–346  
*Formicarius nigricapillus*, 297–319  
*Formicivora rufa*, 85–99  
 Formoso, A., 573–578  
 Fraga, R. M., 15–29., 435–436  
 Fragmentation, 53–63  
 Freile, J. F., 321–336  
 Frere, E., 519–527  
 Furnariidae, 347–359  
*Furnarius cinnamomeus*, 129–136  
*Furnarius rufus*, 85–99  
*Galbalcyrhynchus leucotis*, 289–296  
*Gallbulia ruficauda*, 1–14, 85–99  
*Gampsophylax swainsonii*, 85–99  
 Gandini, P., 519–527  
 Garcia, G. O., 563–566  
 García-Chávez, J., 101–104  
 Garrido, H., 449–470  
 Garrido, O. H., 141  
 Garúa forest, 297–319  
 Gatto, A., 123–126  
 Gelis, R. A., 427–432, 511–518  
 Generalist, 31–42  
*Geothlypis auricularis*, 297–319  
*Geothlypis trichas*, 219–230  
*Geotrygon montana*, 1–14, 129–136, 297–319  
*Geranospiza caerulescens*, 297–319  
 Geupel, G. R., 183–174  
 Girón, L. E., 557–561  
*Glaucidium brasiliense*, 85–99  
*Glaucidium peruanum*, 297–319  
*Glaucis hirsuta*, 1–14  
*Glyptosternon spirurus*, 129–136, 297–319  
 Gopar, A., 529–537  
*Grallaria albicollis*, 231–251  
*Grallaria guatimalensis*, 297–319  
*Grallaria quitensis*, 567–571  
*Grallaria watkinsi*, 297–319  
*Grallaricula nana*, 137–140  
 Grassland, 503–510  
 Greeney, H. F., 137–140, 283–286, 427–432,  
     437–438, 439–440, 511–518, 567–571  
 Guatemala, 419–426  
 Guayaquil Gulf, 175–185  
 Guerrero, S. A., 405–417  
 Guevara, E., 129–136  
 Guevara, J., 129–136  
 Guild, 129–136  
*Guira guira*, 85–99  
*Gymnopithys leucaspis*, 297–319  
 Habitat association, 43–52  
 Habitat fragmentation, 129–136  
 Habitat patch size, 529–537  
 Habitat use, 371–385, 405–417, 419–426  
 Habitat, 129–136, 205–217, 405–417  
 Hahn, I., 371–385  
 Haiti, 219–230  
 Halupka, K., 427–432  
*Haplospiza rustica*, 493–501  
 Harbers, N., 283–286  
*Harpagus bidentatus*, 297–319  
*Harpagus diodon*, 197–204, 547–549  
 Hayes, F. E., 263–266  
*Heliodoxa jacula*, 297–319  
*Heliomaster furcifer*, 231–251  
*Helornis fulica*, 85–99  
*Heliothryx barroti*, 129–136  
*Helminthorhynchus vermivorus*, 219–230  
*Hemitraupis guira*, 1–14, 85–99, 129–136  
*Hemitriccus margaritaceiventer*, 85–99, 231–251  
*Henicorhina leucophrys*, 129–136, 297–319  
*Herpetotheres cachinnans*, 297–319

- Herpsilochmus atricapillus*, 85–99 231–251  
*Herrera*, M., 85–99  
*Heterospizias meridionalis*, 85–99  
*Hirundinea ferruginea*, 85–99  
*Hirundo rustica*, 85–99  
 Hispaniolan endemic bird, 219–230  
 Historical review, 321–336  
 Höfling, E., 471–479  
 Home range, 405–417  
 Homogenization, 503–510  
 Honig, R. A., 85–99  
 Hurricane, 405–417  
*Hydropsalis brasiliiana*, 85–99  
*Hylocharis chrysura*, 85–99  
*Hylocryptus erythrocephalus*, 297–319  
*Hylophilus decurtatus*, 297–319  
*Hylophilus flavipes*, 1–14  
*Hylophylax*, 187–195  
*Hyptilus ruficollis*, 1–14  
*Icterus bonana*, 449–470  
*Icterus cayanensis*, 85–99  
*Icterus dominicensis*, 449–470  
*Icterus icterus*, 85–99  
*Icterus laudabilis*, 449–470  
*Icterus melanopsis*, 449–470  
*Icterus northropi*, 449–470  
*Icterus oberi*, 449–470  
*Icterus portoricensis*, 449–470  
*Icterus praecox*, 449–470  
*Icterus prosthemelas*, 449–470  
*Ictinia mississippiensis*, 109–115  
*Ictinia plumbea*, 109–115, 197–204, 231–251, 297–319  
*Ilicura militaris*, 65–83  
 Incubation, 137–140  
 Independent evolution, 347–359  
 Indicador species, 297–319  
 Introductions, 371–385  
 Irestedt, M., 347–359  
 Itatiaia National Park, 547–549  
*Jacana jacana*, 85–99  
 Juan Fernández Island, 371–385  
 Kinematics, 471–479  
 Kirkconnell, A., 141, 449–470  
*Klais guimeti*, 1–14  
*Knipolegus aterrimus*, 231–251  
*Knipolegus signatus*, 231–251  
 Komar, O., 557–561  
 Kun, M., 539–546  
 Labels, 277–278  
 Landscape structure, 529–537  
*Larus atlanticus*, 123–126  
*Lathrotriccus euleri*, 65–83, 231–251  
*Lathrotriccus griseipectus*, 297–319  
 Latin America, 183–174  
 Latta, S. C., 183–174  
*Legatus leucophaius*, 85–99  
*Lepidocolaptes souleyetii*, 1–14, 129–136, 297–319  
*Lepidocolaptes angustirostris*, 85–99  
*Lepidotrix coronata*, 481–491  
*Leptodon cayanensis*, 197–204, 297–319  
*Leptopogon amaurocephalus*, 65–83, 231–251  
*Leptotila megalura*, 231–251  
*Leptotila ochraceiventris*, 297–319  
*Leptotila pallida*, 297–319  
*Leptotila verreauxi*, 1–14, 85–99, 129–136, 231–251, 297–319  
*Leucopernis albicollis*, 85–99  
*Leucopernis occidentalis*, 297–319  
 Leveau, C. M., 503–510  
 Leveau, L. M., 503–510  
*Limnoctites*, 347–359  
*Limnornis*, 347–359  
*Limnothlypis swainsonii*, 219–230  
 Lobo, R., 145–161  
 Loetti, V., 551–555  
 Londoño, G. A., 205–217  
 López de Aquino, S., 557–561  
*Lophotriccus pileatus*, 129–136, 297–319  
 Lower Paraná, 117–121  
*Loxia megaplaga*, 219–230  
*Loxigilla violacea*, 219–230  
 Macaya Biosphere Reserve, 219–230  
 Machalilla National Park, 297–319  
 Machicote, M., 529–537  
*Malacoptila panamensis*, 129–136  
 Malizia, L. R., 231–251  
*Manacus manacus*, 297–319  
 Manakins, 481–491  
 Mar Chiquita Reserve, 563–566  
 Mariano-Jelicich, R., 563–566  
 Marini, M. A., 65–83  
 Martin, P. R., 427–432, 567–571  
 Martinez, J. J., 529–537  
 Massoni, V., 551–555  
 Mature forest, 297–319  
 Maugeri, F. G., 117–121

- Mecocerculus bellmayri*, 231–251  
*Mecocerculus leucophrys*, 231–251  
*Megarynchus pitangua*, 85–99, 129–136, 297–319  
*Melanerpes candidus*, 85–99  
*Melanerpes pucherani*, 297–319  
*Melanerpes rubricapillus*, 1–14  
*Melanerpes striatus*, 219–230  
*Melanopareia elegans*, 297–319  
 Mennill, D. J., 433–434  
*Mesembrinibis cayanaensis*, 85–99  
 Methodology, 65–83  
 México, 101–104, 279–281  
*Micrastur ruficollis*, 231–251  
*Micrastur semitorquatus*, 297–319  
*Microbates cinereiventris*, 129–136  
*Microcerulus marginatus*, 129–136, 297–319  
 Migration, 109–115, 547–549  
 Migratory birds, 183–174  
 Migratory routes, 145–161  
 Millones, A., 519–527  
*Milvago chamao*, 31–42  
*Milvago chimachima*, 85–99  
 Mimicry, 387–396  
*Mimus dorsalis*, 435–436  
*Mimus gilvus*, 263–266  
*Mionectes oleagineus*, 129–136, 297–319  
*Mionectes olivaceus*, 129–136, 297–319  
*Mionectes rufiventris*, 65–83  
*Miozetetes similis*, 129–136  
 Mites, 573–578  
*Mniotilla varia*, 219–230  
 Molecular phylogeny, 15–29  
 Molecular sexing, 481–491  
 Molecular systematics, 347–359  
*Molothrus bonariensis*, 85–99, 129–136, 435–436  
 Molt, 267–270, 481–491  
*Momotus momota*, 85–99, 231–251  
*Monasa nigrifrons*, 85–99  
 Monitoring, 183–174  
 Montane cloud forest, 419–426  
 Montane forest, 43–52, 53–63, 53–63, 231–251  
 Morton, E. S., 397–404  
 Multilayered eggshell, 263–266  
 Muñoz, M. C., 205–217  
*Myadestes genibarbis*, 219–230  
*Myadestes ralloides*, 129–136  
*Mycteria americana*, 85–99  
*Myiarchus swainsoni*, 85–99, 231–251  
*Myiarchus tuberculifer*, 129–136, 231–251, 297–319  
*Myiarchus tyrannulus*, 85–99  
*Myiarchus venezuelensis*, 1–14  
*Myiobius sulphureipygius*, 129–136, 297–319  
*Myiobius villosus*, 297–319  
*Myioborus miniatus*, 129–136, 297–319  
*Myiodinastes maculatus*, 1–14, 85–99, 231–251, 297–319  
*Myiopagis gaimardi*, 1–14  
*Myiopagis subplacens*, 297–319  
*Myiopagis viridicata*, 85–99, 231–251  
*Myiophobus fasciatus*, 129–136, 297–319  
*Myiotriccus ornatus*, 129–136  
*Myiozetetes similis*, 289–296  
*Myiozetetes cayanensis*, 85–99  
*Myrmeciza exsul*, 129–136, 397–404  
*Myrmeciza immaculata*, 297–319  
*Myrmeciza longipes*, 397–404  
*Myrmeciza*, 187–195  
*Myrmecophilus strigilatus*, 85–99, 231–251  
*Myrmotherula pacifica*, 129–136  
*Myrmotherula schisticolor*, 297–319  
 Natural history, 105–108, 283–286, 427–432, 437–438, 439–440  
*Nemosia pileata*, 85–99  
 Neognathous birds, 471–479  
*Nesocites micromegas*, 219–230  
 Nest predation, 53–63, 253–262  
 Nest, 101–104, 137–140, 279–281, 283–286, 427–432, 433–434, 437–438, 493–501, 511–518, 557–561, 567–571  
 Nesting, 427–432  
 Nestlings, 283–286, 437–438, 439–440, 511–518, 573–578  
 Nidification, 347–359  
 Nocolossi, G., 231–251  
 Nocturnal activity, 123–126  
*Nothura boraquira*, 85–99  
*Notiochelidon cyanoleuca*, 85–99, 129–136, 297–319  
*Nyctibeus griseus*, 85–99, 297–319  
*Nycticorax nycticorax*, 85–99  
*Nyctidromus albicollis*, 297–319  
*Nystalus chacuru*, 85–99  
*Nystalus maculatus*, 85–99  
*Nystalus radiatus*, 297–319  
 O’Shea, B. J., 361–370  
 Ocmin-Petit, A., 289–296

- Odontophorus erythrops*, 129–136, 297–319  
*Odontophorus gujanensis*, 557–561  
 Off-territory forays, 397–404  
 Olivo, C., 109–115  
 Olson, S. L., 347–359  
*Oporornis philadelphus*, 1–14  
 Opportunist, 31–42  
 Orcés, G., 321–336  
*Ornithonyx bursa*, 573–578  
*Ortalis erythroptera*, 297–319  
*Ortalis ruficauda*, 1–14  
 Ortiz, F., 321–336  
*Oryzoborus angolensis*, 85–99  
 Otongachi, 129–136  
*Otus roboratus*, 297–319  
*Otus watsonii*, 85–99  
 Ovenbird, 347–359  
*Pachyramphus cinnamomeus*, 129–136  
*Pachyramphus polychopterus*, 85–99, 231–251  
*Pachyramphus validus*, 85–99  
*Pachyramphus viridis*, 85–99, 231–251  
 Palacios-Silva, R., 101–104  
 Paleognathous birds, 471–479  
 Pampa, 503–510  
 Panama, 187–195  
*Pandion haliaetus*, 109–115  
 Pando-Vasquez, L., 289–296  
*Parabuteo unicinctus*, 109–115, 297–319  
*Paragoniocotes heterogenitalis*, 573–578  
*Paragoniocotes semicingulatus*, 573–578  
 Paramo, 567–571  
 Parasite removal, 137–140  
 Parental care, 511–518, 551–555  
*Paroaria capitata*, 85–99  
 Parrots, 43–52  
*Parula pitiayumi*, 85–99, 129–136, 297–319  
*Passer domesticus*, 371–385  
*Patagioenas cayennensis*, 231–251  
*Patagioenas flavirostris*, 279–281  
*Patagioen's inornata*, 219–230  
*Patagioen's squamosa*, 219–230  
 Patagonia, 123–126, 539–546  
*Penelope obscura*, 231–251  
*Penelope purpurascens*, 297–319  
*Penelope superciliaris*, 85–99  
*Penelopina nigra*, 419–426  
 Persistente, 253–262  
 Peru, 387–396  
*Phacellodomus rufifrons*, 85–99  
*Phaenicophilus poliocephalus*, 219–230  
*Phaethornis anthophilus*, 1–14  
*Phaethornis baroni*, 297–319  
*Phaethornis pretrei*, 231–251  
*Phaethornis striigularis*, 1–14  
*Phaetornis yaruqui*, 129–136  
*Phalacrocorax gaimardi*, 519–527  
*Pharomachrus auriceps*, 129–136  
*Pheucticus chrysogaster*, 297–319  
*Phyllomyias griseiceps*, 297–319  
*Phylloscartes ventralis*, 231–251  
*Piaya cayana*, 85–99, 129–136, 231–251, 297–319  
*Piculus rubiginosus*, 129–136, 231–251, 297–319  
*Picumnus albosquamatus*, 85–99  
*Picumnus cirratus*, 231–251  
*Picumnus squamulatus*, 1–14  
*Pilherodius pileatus*, 85–99  
*Pionus chalcopterus*, 297–319  
*Pionus maximiliani*, 43–52, 85–99, 231–251  
*Pionus menstruus*, 129–136, 289–296  
*Pionus sordidus*, 1–14  
*Pipile cumanensis*, 85–99  
*Pipra chloromeros*, 433–434  
*Pipra erythrocephala*, 1–14  
*Pipra filicauda*, 1–14, 481–491  
*Pipra pipra*, 481–491  
*Pipridae*, 433–434, 481–491  
*Piranga flava*, 85–99  
*Piranga lutea*, 297–319  
*Piranga rubra*, 297–319  
*Pitangus lictor*, 85–99  
*Pitangus sulphuratus*, 1–14, 85–99, 231–251  
*Platalea ajaja*, 175–185  
*Platycichla flavipes*, 1–14  
*Platyparis homochrous*, 297–319  
*Platyrinchus mystaceus*, 65–83, 297–319  
*Podager naevia*, 85–99  
*Poecilotriccus plumbeiceps*, 231–251  
*Poecilotriccus ruficeps*, 427–432  
*Pogonotriccus ophthalmicus*, 129–136  
*Polioptila dumicola*, 85–99  
*Polioptila plumbea*, 129–136, 297–319  
 Politi, N., 43–52, 231–251  
*Polyborus plancus*, 85–99, 109–115  
 Population change, 289–296  
 Population, 175–185, 205–217  
 Porzecanski, A. L., 85–99  
 Predators, 187–195

- Prey size, 539–546  
*Priotelus roseigaster*, 219–230  
*Prugne chalybea*, 85–99  
*Propyrrhura auricollis*, 43–52, 231–251  
 Protective nesting, 117–121  
*Psarocolius decumanus*, 1–14, 289–296  
*Psarocolius decumanus*, 85–99  
*Pseudotriccus pelzelni*, 511–518  
*Pseutes*, 187–195  
*Pteroglossus castanotis*, 85–99  
*Pteroglossus erythropygius*, 129–136, 297–319  
 Puebla, 101–104  
 Puerto Rico, 405–417  
*Pulsatrix perspicillata*, 297–319  
*Pyriglenia leuconota*, 297–319  
*Pyrocephalus rubinus*, 85–99  
*Pyrrhura molinae*, 43–52, 85–99, 231–251  
 Quintana, F., 123–126  
 Radiotelemetry, 397–404  
 Ralph, C. J., 183–174  
*Ramphastos brevis*, 297–319  
*Ramphastos toco*, 85–99, 231–251  
*Ramphastos tucanus*, 289–296  
*Ramphocelus carbo*, 1–14  
*Ramphocelus icteronotus*, 129–136, 297–319  
 Raptors, 109–115, 197–204, 547–549  
 Rarity, 205–217  
 Reboreda, J. C., 551–555  
 Renesting interval, 253–262  
 Renner, S. C., 419–426  
 Renous, S., 471–479  
 Reproduction, 101–104  
 Reproductive success, 253–262  
 Research, 321–336  
 Restoration, 405–417  
*Rhynchoscyllus pacificus*, 297–319  
*Rhynchotus rufescens*, 85–99  
*Rhynchotus rufescens*, 471–479  
 Ría Deseado, 519–527  
 Richness, 503–510  
 Kimmer, C. C., 219–230  
 Ríos, M. M., 205–217  
 Rivera, L. O., 43–52, 231–251  
 Robinson Crusoe Islands, 371–385  
 Robinson, T. R., 187–195  
 Robinson, W. D., 187–195  
 Rodríguez, Y., 141  
 Rodríguez-Ferraro, A., 105–108  
 Rombough, J. F., 439–440  
 Römer, U., 371–385  
 Rompré, G., 187–195  
 Roper, J. J., 253–262  
*Rostrhamus sociabilis*, 85–99, 109–115  
*Rupicola peruviana*, 105–108  
 Rural environment, 503–510  
 Ryder, T. B., 481–491  
 Saavedra, A. M., 85–99  
 Sahores, M., 539–546  
*Saltator atripennis*, 129–136  
*Saltator grossus*, 297–319  
*Saltator maximus*, 129–136, 297–319  
*Saltator striaticeps*, 1–14, 297–319  
 Sánchez, C., 493–501  
*Sappho sparganura*, 231–251  
*Sarcoramphus papa*, 85–99, 109–115, 297–319  
*Sauvagea longirostris*, 219–230  
*Sayornis nigricans*, 129–136  
*Schiffornis turdinus*, 297–319  
 Schlatтер, R. P., 371–385  
*Sclerurus guatemalensis*, 297–319  
*Sclerurus mexicanus*, 297–319  
*Scytalopus latrans*, 283–286  
*Scytalopus parkeri*, 439–440  
 Seasonal variation, 231–251  
 Secondary forest, 297–319  
 Seijas, S., 539–546  
 Seipke S. H., 547–549  
*Seiurus aurocapillus*, 219–230  
*Seiurus noveboracensis*, 1–14  
*Sephanoides fernandensis*, 371–385  
*Sephanoides sephanioides*, 371–385  
*Serpophaga cinerea*, 129–136  
*Serpophaga subcristata*, 85–99, 231–251  
 Serra da Mantiqueira, 547–549  
*Setophaga ruticilla*, 1–14, 219–230  
 Sexual size dimorphism, 551–555  
 Shrubs, 529–537  
 Sierra Yalijux, 419–426  
 Sipaliwini Savanna, 361–370  
*Sittasomus griseicapillus*, 1–14, 85–99, 231–251, 297–319  
 Snakes, 187–195  
 Social foraging, 271–275  
 Solórzano, A., 1–14  
 Song, 397–404, 493–501  
 Sornosa, F., 137–140  
 Species abundance, 85–99  
 Species composition, 231–251

- Species richness, 85–99  
 Species turnover, 231–251  
*Spindalis dominicensis*, 219–230  
*Spizaetus tyrannus*, 297–319  
*Spizaetus ornatus*, 297–319  
*Sporophila corvina*, 297–319  
*Sporophila intermedia*, 1–14  
*Sporophila lineola*, 289–296  
*Sporophila luctuosa*, 387–396  
*Sporophila nigricollis*, 1–14  
 Stability, 289–296  
*Stelgidopteryx ruficollis*, 1–14, 85–99, 129–136  
*Sterna superciliaris*, 117–121  
*Sterna trudeani*, 563–566  
*Sternoclyta cyanopectus*, 1–14  
 Stomach contents, 277–278  
*Streptoprocne zonaris*, 129–136, 231–251, 297–319  
*Strix nigrolineata*, 297–319  
*Strix virgata*, 297–319  
 Study skins, 277–278  
 Stutchbury, B. J. M., 397–404  
 Suárez, N., 123–126  
*Sublegatus modestus*, 85–99  
 Suriname, 361–370  
 Survey design, 337–346  
 Survey, 205–217  
*Synallaxis albescens*, 85–99  
*Synallaxis azarae*, 231–251  
*Synallaxis brachyura*, 129–136, 297–319  
*Synallaxis frontalis*, 231–251  
*Synallaxis gujanensis*, 85–99  
*Synallaxis scutata*, 231–251  
*Syndactyla rufosuperciliata*, 231–251  
 Syntopic breeding, 387–396  
*Tachycineta euchrysea*, 219–230  
*Tachyphonus luctuosus*, 129–136  
*Tachyphonus rufus*, 1–14, 129–136  
*Tanager conothraupis*, 387–396  
*Tangara cyanicollis*, 129–136, 297–319  
*Tangara gyrola*, 129–136, 297–319  
*Tangara icterocephala*, 129–136, 297–319  
*Tapera naevia*, 231–251  
*Taraba major*, 85–99, 231–251, 297–319  
 Tartar emetic, 65–83  
 Taxonomy, 449–470  
 Tehuacán-Cuicatlán, 101–104  
 Terrestrial locomotion, 471–479  
 Territorial behavior, 397–404  
 Territory defense, 397–404  
*Thalurania fannyi*, 129–136, 297–319  
*Thamnophilus atrinucha*, 253–262  
*Thamnophilus caerulescens*, 65–83, 85–99, 231–251  
*Thamnophilus doliatus*, 1–14, 85–99  
*Theristicus caudatus*, 231–251  
*Thlypopsis sordida*, 85–99  
*Thraupis episcopus*, 1–14, 129–136, 297–319  
*Thraupis palmarum*, 1–14, 129–136, 297–319  
*Thraupis sayaca*, 85–99  
*Thryothorus guarayanus*, 85–99  
*Thryothorus mystacalis*, 297–319  
*Thryothorus rufalbus*, 1–14  
*Thryothorus sclateri*, 297–319  
*Tiaris bicolor*, 219–230  
*Tiaris fuliginosa*, 1–14  
*Tiaris olivaceus*, 219–230  
*Tigrisoma lineatum*, 85–99  
 Tinamidae, 471–479  
*Tinamus solitarius*, 471–479  
*Tityra cayana*, 85–99  
*Tityra inquisitor*, 85–99  
*Tityra semifasciata*, 129–136, 297–319  
*Todirostrum cinereum*, 297–319  
*Todus angustirostris*, 219–230  
*Tolmomyias flavotectus*, 297–319  
*Tolmomyias sulphurescens*, 1–14, 231–251, 297–319  
 Townsend, A. K., 219–230  
 Townsend, J. M., 219–230  
 Trejo, A., 539–546  
*Trichothraupis melanops*, 65–83  
 Trijillo, G., 53–63  
*Tringa solitaria*, 85–99  
 Trinidad, 263–266  
*Troglodytes aedon*, 1–14, 85–99, 129–136  
*Trogon caligatus*, 297–319  
*Trogon collaris*, 1–14, 297–319  
*Trogon curucui*, 85–99, 231–251  
*Trogon mesurus*, 297–319  
 Trophic guilds, 65–83  
 Tropical forest, 289–296  
 Tropical montane forest, 53–63  
 Tropical mountain cloud forest, 419–426  
*Turdus albicollis*, 65–83  
*Turdus daguae*, 297–319  
*Turdus falcklandii*, 371–385  
*Turdus fumigatus*, 1–14

- Turdus hauxwelli*, 85–99  
*Turdus maculirostris*, 129–136, 297–319  
*Turdus nudigenis*, 1–14  
*Turdus plumbeus*, 219–230  
*Turdus reevei*, 297–319  
 Tyrannidae, 145–161  
*Tyrannus melancholicus*, 85–99, 129–136  
*Tyrannus savana*, 85–99  
*Tyto alba*, 539–546  
 Understory, 1–14  
 Urban environment, 503–510  
 Urbanization, 503–510  
 Venezuela, 1–14, 105–108  
*Veniliornis callonotus*, 297–319  
*Veniliornis frontalis*, 231–251  
*Veniliornis kirkii*, 129–136, 297–319  
 Verea, C., 1–14  
 Videography, 187–195  
 Vilella, F. J., 405–417  
 Villarreal, D., 529–537  
*Vireo olivaceus*, 1–14, 85–99, 231–251, 297–319  
 Vocalizations, 15–29  
*Volatinia jacarina*, 85–99  
 Weidensaul, C. S., 267–270  
 Welcker, J. J., 85–99  
 West Indies, 449–470  
 White, T. H. Jr., 405–417  
 Wiley, J. W., 141, 449–470  
 Wintering, 145–161, 183–174  
 Witt, C. C., 387–396  
 Wolfenden, B. E., 397–404  
 Woltmann, S., 337–346  
*Xanthopsar flavus*, 15–29  
*Xenoligea montana*, 219–230  
*Xenops minutus*, 1–14, 297–319  
*Xenops rutilans*, 85–99, 231–251, 297–319  
*Xiphocolaptes major*, 85–99, 231–251  
*Xiphocolaptes promeropirhynchus*, 129–136  
*Xiphorhynchus erythropygius*, 297–319  
*Xiphorhynchus picus*, 85–99  
*Xiphorhynchus susurans*, 1–14  
 Yorio, P., 123–126  
 Yungas, 145–161  
 Zaldaña Fonseca, C. M., 557–561  
*Zenaida auriculata*, 85–99  
*Zenaida aurita*, 219–230  
*Zenaida macroura*, 219–230  
*Zonotrichia capensis*, 85–99

## REVIEWERS FOR ORNITOLOGÍA NEOTROPICAL, VOLUME 16

Each manuscript submitted for publication in *Ornitología Neotropical* is generally reviewed by two specialists, often including a member of the editorial board, and frequently by a third one. The following individuals kindly reviewed one or more (\*) manuscripts for Volume 16: Jorge L. B. Albuquerque, José Luis Alcantara, Marina Aanciäes, Adrián Azpiroz\*, John M. Bates, Marc Béchard, A. R. Biknevicius, Keith L. Bildstein, John Blake\*, Pedro Blendinger, Elisa Bonaccorso\*, Sérgio H. Borges, D. Brightsmith\*, Daniel M. Brooks, D. E. Burhans, Daniel Cadena\*, Patricia Caplonch, J. Carroll, R. J. Casaux, J. F. Chace, M. Cohn-Haft, Jennifer O. Coulson, Andres M. Cuervo, Liliana M. Davalos, Andrew Dawson, R. D. Dawson, R. C. Dobbs, David Duffy, Jessica Eberhard, Christian Érard\*, Carla S. Fontana, Rosendo Fraga\*, Juan Freile\*, Adrián Di Giacomo, Julio C. Gallardo del Angel, Rudolphe A. Gelis, Stephen S. Germaine, Michael Gochfeld, Patricia Gonzalez, F. González-Garcia, Jennifer Gervais, Harold F. Greeney\*, Floyd E. Hayes\*, Fritz Hertel, Juan Pablo Isaach, Frankie D. Jackson, Fabián Jaksic\*, Jaime Jiménez\*, A. R. Keith, Oliver Komar, Niels Krabbe, T. A. Langen, Steve Latta, Catherine Lindell\*, Bette A. Loiselle, I. J. Lovette, Russell Main, Tiego Marques, Raymond McNeil\*,

Felix M. Medina, Myrian E. Mermoz, J. Michael Meyers, M. C. Meyers, Javier Millan, Diego Montalti, Luis Gonzalo Morales, Sergio G. Mosa, Luis Germán Naranjo, Manuel Nores\*, Christian Olivo, K. E. Omland, Vincenzo Penteriani, P. J. Pietz, Augusto Piratelli, Pablo Porras-Peñaanda, Brigitte Poulin\*, E. R. Price, Richard O. Prum, Peter Pyle, Flavio Quintana, J. van Remsen, Nate Rice, Gary Ritchison, W. Douglas Robinson\*, G. H. Rosenberg, Brett K. Sandercock, J. A. Sánchez-Zapata, Virginia Sanz, Jean-Pierre Savard, Roberto Schlatter, Karl-L. Schuchmann, Tom S. Schulenberg\*, Cagan Sekercioğlu, Thomas Sherry, Rodney Siegel, Neil G. Smith, Roger Smith, J. K. Schmutz, N. S. Sodhi\*, F. Gary Stiles, José Tello, Marc Théry\*, Jean-Marc Thiollay\*, R. Thorstrom, J. C. Torres-Mura, Olivier Tostain, Alejandro Travaini, Daniel Varland, David Varrichio, Carlos Verea, Pascal Villard, François Vuilleumier\*, P. J. Weatherhead, M. S. Webster, Thomas H. White, Jim Wiley\*, Edwin O. Willis\*, L. Wolf, Pablo Yorio, Kristof Zyskowski\*.

The abstracts were kindly corrected by Cristina Yumi Miyaki for the Portuguese, and by Elisa Bonaccorso for the Spanish. Page-setting was done by Barbara Durán, Elisa Muñoz and María Rojas G.