

T. V.	1893.	GENERAL NOTES.
Portland, Conn.	common	
Belcherstown, Mass.		seen 5-10-'83
Phillips, Me.	common	

A bird of the lowest branches and underbrush, often feeding upon the ground. At once recognized by the white spot at the base of the wing quills. Prof. Kumlein finds it a common migrant in Wisconsin: "apparently less common than 20 or 30 years ago."

Dendroica coronata (Linn.) YELLOW-RUMPED WARBLER.

T. V.	S. R.	1893.	GENERAL NOTES.
Starke, Fla.	tol. com.		
Coffeyville, Kan.	common		
Garden City, Kan.	common	4-21	
Jackson Co., Mo.	common		
Grinnell, Iowa.	abundant		one pair known to breed
St. Paul, Minn.	abundant		
Winnebago Co., Ill.	abundant	4-12	L. 5-12-'93
Delavan, Wis.	common		
Milton, Wis.	common	4-19	
Meridian, Wis.	abundant		
Palmer, Mich.	common	4-30	9-17 to 10-17-'93
Hamilton Co., O.	abundant		4-29-'91 10-3 to 10-25-'91
Lorain Co., O.	common	5-10	4-12-'94 L. 5-12-'94
Berwyn, Pa.	common	4-12	9-1 to 10-30-'92
Cortland Co., N. Y.	abundant	5.5	{ '92 record: males 5-2; L. males; F. females 5-18; L. female 5-24.
Oswego City, N. Y.	abundant		
Portland, Conn.	common		
Wethersfield, Conn.	common		
Duxbury, Mass.	common		} remained all winter and fed on grass seeds
Belcherstown, Mass.	common	rare	one pair bred
Bangor, Me.	common	common	
Phillips, Me.	common	common	
Toronto, Can.		not com.	

Usually very common in Lorain Co., O.; I did not see a dozen birds during the spring of 1893. This bird is worthy of special study. He is the earliest of the warblers.