

campus of the Agricultural College. On this latter date their similarity to Cowbirds was quite marked, and suggested a very probable reason for their being overlooked in past years. Their habits are certainly very dissimilar to those of the closely related Rusty Blackbird (*Euphagus carolinus*), and this fact should be borne in mind in watching for these birds in migration. Lack of suspicion is another trait characteristic of the small flocks seen in western North Carolina and Georgia, so positive identification is never a difficult matter.—THOS. D. BURLEIGH, *Bureau of Biological Survey, New Orleans, La.*

A Record of the Trumpeter Swan from the Late Pleistocene of Illinois.

—In material secured at Aurora, Illinois, by Professor Clarence R. Smith of Aurora College, forwarded to me for examination by Dr. L. A. Adams of the University of Illinois, I have identified humeri, a broken sternum, and part of the scapula of the Trumpeter Swan (*Cygnus buccinator*). According to Professor Smith¹ these specimens were found in a marl deposit underlying a peat bog above which was a layer of muck forming the bottom of a swamp in Phillip's Park in the southwest part of Aurora at a point a mile and a half east of the Fox River. They were obtained by C. W. A. workers during excavation of the swamp to make an artificial lake.

The swan remains were associated with bones of mastodon (*Mastodon americanus*) and giant beaver (*Castoroides ohioensis*) and are believed to have been deposited in the bed of a post-glacial lake. According to the findings of Dr. O. P. Hay² this would seem to place them in what Dr. Hay has termed the Wabash beds in late Pleistocene. Dr. Adams writes me that mollusks of Pleistocene species were taken from the pneumatic foramina of the swan humeri.

The bones are distinguished at a glance as those of the Trumpeter Swan. The sternum, while in fragments, shows the characteristic bulbous swelling projecting into the body cavity at the anterior end in addition to the swollen channel in which the trachea is folded. The humeri exhibit two sizes, possibly indicative of sexual difference in wing measurement, and are greater in size than the largest Whistling Swans. All of the specimens, which bear numbers 111, 734, B, C, D, F, H, J, and K, are fresh and clean in appearance, and are in good state of preservation.

Previously this swan has been reported from Pleistocene deposits in Oregon and Florida, the present being the first occurrence of it in the central portion of our country. The find is one of definite importance in view of the few reports of birds that have come from Pleistocene beds of the area in question.

The specimens have been returned to Professor Smith, to whose kindness and that of Dr. Adams I am indebted for permission to place this occurrence on record, and are preserved by the city of Aurora in a museum at Phillip's Park.—ALEXANDER WETMORE, *U. S. National Museum, Washington, D. C.*

Notes on the Nesting of Captive Mute Swans.—In the latter part of June, 1930, four Mute Swans (*Cygnus olor*) about a year old, and still in the gray juvenile plumage, were placed upon eighty-seven acre Fowler Lake, in the city of Oconomowoc, Waukesha County, Wisconsin. These birds were a single brood from a captive pair. Two additional birds that had been hatched the previous year, but from another pair, were placed on the lake later in

¹Science, Vol. 81, April 19, 1935, p. 380.

²Carnegie Inst. Washington, Publ. 322, 1923, p. 13.

the summer. The four original birds stayed by themselves on the lake, and would not allow the other two to come near them, but pursued them whenever they could get near them. All birds were wing-clipped at the time, but were pinioned before being taken to winter quarters.

The following spring two of the original four birds mated and started building a nest during the latter part of April. The site chosen was in a cattail marsh in the Oconomowoc River, where the stream flows into Fowler Lake. It could easily be seen from a bridge, and as a number of people came to observe the building operations, the pair abandoned this site and moved to a more secluded spot farther up stream. On June 8, 1931, the pair was seen on the lake with five newly hatched young, which must have been less than twenty-four hours old.

The Mute Swan is ordinarily supposed not to mate before three years of age, but this pair was approximately two years of age at the time of producing young.—WILLIAM JOHNSTON HOWARD, *Petoskey, Mich.*

The 1934 Fall Migration at Cleveland's Public Square.—Migrating birds were much less common at the Square this season than most years, according to my observations. The only reason that I can propose is that, with the unusual amount of clear weather which occurred during the fall months, migrants were flying higher over Lake Erie on their southward journey, and passed over the city before descending to rest in more attractive surroundings beyond the congested area.

My records for the ten species which I observed this season are as follows:

Species	First Record	Last Record	No. Days Seen	Largest No. in One Day
Canada Goose	Nov. 1	—	1	8
Sparrow Hawk	Sept. 13	Dec. 20	4	1
Herring Gull	Sept. 17	Dec. 17	11	9
Mourning Dove	Aug. 23	—	1	1
Northern Yellow-throat	Sept. 19	Oct. 22	2	1
White-crowned Sparrow	Sept. 19	Oct. 27	2	1
White-throated Sparrow	Oct. 2	Nov. 7	19	3
Song Sparrow	Oct. 10	Oct. 15	2	1
Swamp Sparrow	Oct. 8	—	1	1
Lincoln Sparrow	Sept. 18	Sept. 19	2	1

In addition, evening newspapers carried a story on October 2 that an American Bittern (identified by a qualified ornithologist) had been captured alive in the Square that morning. The geese and the Mourning Dove, the only new additions to my list, were observed flying over at a low altitude.

Starlings were as numerous as ever. This year they took over the terminal tower as one of their roosting spots, many of them finding crannies five or six hundred feet above the street, where a strong wind is blowing much of the time. This must have been a miserable roost in severe weather, but it seemed to make little difference.—WILLIAM H. WATTERSON, *Cleveland, Ohio.*

Some Corrections.—I should like to make the following corrections to my article on Walter John Hoxie in the WILSON BULLETIN, XLVI, September, 1934: On page 173, in the map of the Sea Islands, "Hunters Island" should read Hunting Island. On page 174, last line, the date should read July 2, 1906. On page 176, line 17, it was a mistake to say that "Later observations have confirmed Hoxie's records" of the Buff-breasted Sandpiper. Hoxie's notes indicate one was taken on St. Helena Island, South Carolina, on May 5, 1886, but thus far the whereabouts of the specimen is unknown. Among Hoxie's