

Eastern Lark Sparrow, *Chondestes grammacus grammacus*. On June 22, 1944, I discovered an adult, followed by two recently fledged young, on a dry limestone-studded slope west of the Potomac River and six miles north of Shepherdstown, Jefferson County. Breeding records of this species east of the mountains are rather rare.—LAWRENCE E. HICKS, *Ohio State University, Columbus, Ohio*.

**Bird remains from an Indian village site in Ohio.**—In 1917 Mills ("Certain Mounds and Village Sites in Ohio," Vol. 3, Pt. 1, p. 46) reported six species of birds from the Feurt Village Site which is located about five miles north of Portsmouth in Clay Township, Scioto County, Ohio, just east of the Scioto River. From 1937 to 1942, excavations for archaeological specimens were carried on at this site by Dr. Stanley Copeland, H. R. McPherson, and Philip Kientz, of Columbus, Ohio, who generously allowed me to study the animal remains that they recovered there. The material is presumed to date from the fifteenth or the sixteenth century.

From among the remains 646 bird bones were identified; 439 are from the Turkey, but altogether, at least 30 species of birds are represented. Eight of these have been reported by Alexander Wetmore (1943, *Wils. Bull.*, 55:55 and 127) to whom I am indebted for assistance in identifying most of the bones. The complete list of birds now known from the site is as follows:

Common Loon <i>Gavia immer</i>	Ruffed Grouse <i>Bonasa umbellus</i>
Pied-billed Grebe <i>Podilymbus podiceps</i>	Prairie Chicken <i>Tympanuchus cupido</i>
Great Blue Heron <i>Ardea herodias</i>	Bob-white <i>Colinus virginianus</i>
Trumpeter Swan <i>Cygnus buccinator</i>	Turkey <i>Meleagris gallopavo</i>
Canada Goose <i>Branta canadensis</i>	Little Brown Crane <i>Grus canadensis canadensis</i>
Blue or Snow Goose <i>Chen</i> sp.	Sandhill Crane <i>Grus canadensis tabida</i>
Mallard <i>Anas platyrhynchos</i>	Woodcock <i>Philohela minor</i>
Black Duck <i>Anas rubripes</i>	Passenger Pigeon <i>Ectopistes migratorius</i>
Blue-winged Teal <i>Anas discors</i>	Great Horned Owl <i>Bubo virginianus</i>
Wood Duck <i>Aix sponsa</i>	Barred Owl <i>Strix varia</i>
Ring-necked Duck <i>Nyroca collaris</i>	Flicker <i>Colaptes auratus</i>
Lesser Scaup Duck <i>Nyroca affinis</i>	Pileated Woodpecker <i>Ceophloeus pileatus</i>
Red-tailed Hawk <i>Buteo jamaicensis</i>	Ivory-billed Woodpecker <i>Campephilus principalis</i>
Red-shouldered Hawk <i>Buteo lineatus</i>	Raven <i>Corvus corax</i>
Bald Eagle <i>Haliaeetus leucocephalus</i>	Crow <i>Corvus brachyrhynchos</i>
Marsh Hawk <i>Circus cyaneus hudsonius</i>	

—ROBERT GOSLIN, *Department of Physiology, Ohio State University, Columbus, Ohio*.