

Commentary

A CRITIQUE OF WANG YONG AND FINCH'S FIELD-IDENTIFICATIONS OF WILLOW FLYCATCHER SUBSPECIES IN NEW MEXICO

John P. Hubbard¹

In a recent paper in the *Wilson Bulletin*, Wang Yong and Finch (1997; henceforth Y&F) reported that they subspecifically identified 83 of 84 Willow Flycatchers (*Empidonax traillii*) captured, banded, and released in central New Mexico in spring and autumn 1994 and 1995. Given the nature of these subspecies and the means by which Y&F apparently identified them, I am extremely doubtful about the reliability of their determinations and thus the validity of these as scientific data. The fact is that identifying these taxa is quite difficult, even for trained taxonomists working in the laboratory under the best protocols and conditions. This difficulty stems from a number of factors, the major one being the pervasive subtlety of the plumage-color characters by which these subspecies mainly differ. Not surprisingly, these differences are difficult to describe in words, which is exacerbated by the fact that none of the available classification systems accurately portrays the range of plumage coloration observed in this flycatcher (e.g., Browning 1993). This means that this species' plumage-color characters are best observed in specimens (i.e., study or flat skins), which also provide the best avenue for identifying subspecies. To do this, one must first assemble series of skins representing all relevant taxa, as well as such important subcategories as age classes (e.g., adult vs immature) and seasonal groupings (e.g., spring vs autumn). Then one sorts "unknowns" (which could include live birds) into subcategories and compares them to the taxa therein, which should produce at least tentative subspecific identifications. In fact, this is the standard laboratory approach for identifying color-based subspecies, and it is the only means proven

reliable for this purpose in the Willow Flycatcher.

As my earlier comments suggest, I do not believe Y&F used the approach described above in their attempts to identify subspecies in the Willow Flycatcher. In other words, they did not take synoptic series of study skins into the field, against which the birds they captured were compared to determine subspecific identities. However, I cannot be 100% certain about this because the methods section in their paper is so incomplete and otherwise deficient one can only guess at many aspects of their approach. Nonetheless, it seems logical that if they had used skins as the basis for their identifications, they would have said so. Given this assumption, if they did not use skins, how did they go about identifying their birds to subspecies? On this matter Y&F are at best vague, providing a few clues but no definitive explanations of their identification methodology. For example, we are told that they "... adopted the four-subspecies classification system of Hubbard (1987) and Unitt (1987)," in which "subspecies identity ... is based [in part] on ... coloration of the head [= crown] and neck [= forenape] and its contrast with the back, and the contrast between the breastband and the throat (see Phillips 1948, Hubbard 1987, Unitt 1987, Browning 1993)." Based on this, I assume that Y&F chose literature descriptions (as opposed to specimen comparisons) as the basis for their identification of Willow Flycatcher subspecies. In addition, I also suspect they converted these descriptions into the color values of Smithe (1975), as this is the system they used to classify coloration in birds captured in the field. Beyond this, one could also speculate on such matters as (a) how converted values were actually used to identify birds, e.g., whether in a dichotomous key, probability table, or other

¹ Route 5, Box 431, Española, New Mexico 87532.

framework; or (b) what Y&F's perceptions were of color characters in various races, given that no such descriptions were offered by them. However, I see no purpose in further speculation concerning these or other aspects of their methodology. This is because if they did base their identifications on the literature rather than specimens, I believe the process became so flawed that the details are irrelevant—like rearranging deck chairs on the sinking Titanic!

The message here is that the literature is no substitute for specimen comparisons for anyone attempting to identify Willow Flycatcher subspecies, at least if attaining the most reliable scientific data is the goal. Furthermore, given logistical and other problems, I doubt even specimen comparisons would consistently yield reliable identifications of live birds under field conditions. Not only would it be unwieldy to take and use museum skins in the field, but setting up and maintaining constant conditions (e.g., lighting) would also be difficult. In addition, except for recaptures, only one opportunity would be available to identify each live bird in the field. This means that one could not reassess identifications at a later time, which is both frequent and necessary when studying specimens in the laboratory. In this regard, photographs and certainly color readings (e.g., from Smithe 1975) would not be adequate for such reexaminations because these do not exactly duplicate colors observed in the birds or specimens themselves. Given these considerations, I believe that identifying subspecies in the Willow Flycatcher is best done in the laboratory, using study skins examined under proper protocols and procedures by people trained in the process. In other words, this is a task that should be left to an alpha-taxonomic approach, which is appropriate when one considers that subspecies arose and largely remain as products of that realm.

Even when approached as outlined above, the reality is that not every specimen or even population of this flycatcher can be reliably assigned to subspecies. Intergradation and overlap occur in all characters that distinguish these taxa, so birds exhibiting such characteristics may be un- or misidentified as a result. In addition, characteristics in some populations remain poorly known, mainly because of the paucity of specimens from these areas. For

example, in the latest revision of the species, Browning (1993) could only assemble 270 specimens of breeding season adults—including fewer than 20 of the endangered subspecies *E. t. extimus* of the Southwest. As a consequence, it is not surprising that he questioned boundaries between four of the five subspecies recognized in his paper. Even when populational characteristics are better known, opinions may differ as regards their taxonomic treatment. Thus, Browning (1993) recognized two subspecies (i.e., *E. t. traillii* and *E. t. campestris*) as breeding in the region east of the Rocky Mountains, whereas Unitt (1987) merged the latter with the nominate form. Differences in opinion also exist on a broader scale, such as concerning the overall number of subspecies recognizable in the Willow Flycatcher. For example, some taxonomists maintain that none should be recognized (e.g., Mayr and Short 1970, Traylor 1979), while others accept four to six as valid (e.g., Phillips 1948, Aldrich 1951, Wetmore 1972, Oberholser 1974, Unitt 1987, Browning 1993). Thus, although specimen comparisons provide our only reliable means for identifying subspecies in this flycatcher, this approach must be used with the clear recognition that it is just the first step in this very difficult endeavor.

Incidentally, the above differences in taxonomic opinion present a problem for those that rely largely or entirely on the literature for their knowledge of geographic variation in this species. That is, how does one choose which authorities to follow and thus which viewpoints to accept on this subject? Among others, one way around this would be to adhere strictly to a single point of view, such as the recent revision of this flycatcher by Browning (1993). However, Y&F chose not to do this, instead electing to cobble their concept of variation from a variety of sources (e.g., Phillips 1948, Hubbard 1987, Unitt 1987, Browning 1993). Given the lack of consensus among these sources, this was a questionable decision. In fact, it would be a challenge even for people with firsthand experience with geographic variation in this species, as seen from the variety of opinions cited above. As a consequence, it is not surprising that I would quibble with Y&F's choices, including that of which authorities to follow.

For example, as indicated earlier, they cited my unpublished paper (Hubbard 1987) as a basis for the "four-subspecies classification system" adopted in their study. However, that so-called system was actually a cobbling job itself, my aim being to summarize color characters of various subspecies from the treatments of Phillips (1948), Aldrich (1951), Wetmore (1972), and Oberholser (1974). As such, it was not meant either to provide definitive descriptions of these subspecies or to recommend which should be recognized as valid. For it to have been otherwise used by Y&F may seem flattering, but it certainly was not a sound decision from a taxonomic viewpoint.

Given the flawed nature of their approach, it is no surprise that Y&F's findings on Willow Flycatcher subspecies would also be open to question. For example, when compared with what is known from specimens (e.g., Hubbard 1987), significant differences emerge on the New Mexico status of three of the four taxa recognized in that study. (In light of the relative scientific standing of the two sources, I would obviously accept the specimen version over that of Y&F in every case.) The most significant difference occurs in the subspecies *E. t. brewsteri* (sensu stricto), which breeds along the Pacific slope of North America. Although occurring regularly in migration eastward to Arizona (Monson and Phillips 1981), this form has rarely been collected east and north of that state, e.g., in Utah (Behle 1985), Colorado (Bailey and Niedrach 1965), Oklahoma (Sutton 1967), and Texas (Oberholser 1967). Hard data from New Mexico clearly conform to this pattern, with only two (4.7%) of the 43 specimens so attributed in Hubbard (1987) and even these were somewhat equivocal. By contrast, Y&F identified 33 (39.8%) of their 83 birds as *E. t. brewsteri*, which is about 8.5 times more frequent than reported by Hubbard. Another notable departure involves the subspecies *E. t. traillii* (in which Y&F include *E. t. campestris*), which breeds from the Great Plains to the northeastern Atlantic Coast. In the Southwest, *E. t. traillii/campestris* occurs regularly in the plains of eastern Colorado (Bailey and Niedrach 1965) and New Mexico (Hubbard 1987), but it has not been collected as far west as Arizona (Monson and Phillips 1981). Yet Y&F reported that 8.4% of their birds were

this form, even though the the middle Rio Grande Valley lies some 200 miles west of the nearest specimen localities in New Mexico. Finally is the race *E. t. adastus*, which breeds widely in the interior U.S. north of the southwestern states, through which it passes in both spring and autumn. In New Mexico, it comprised 25.6% of the specimens reported by Hubbard (1987), compared to 10.8% in Y&F's sample.

As for the fourth subspecies (*E. t. extimus*), Y&F identified 34 (41.1%) of their birds as this form, compared to the 48.8% from throughout New Mexico by Hubbard (1987). Thus, on the face of it, their findings would seem not to differ significantly from what is known from specimens of this taxon. However, the number of questionable literature records of this subspecies suggests it may be more subject to misidentification than certain other forms, such *E. t. brewsteri* and *E. t. traillii* (both sensu lato). Birds that might be mistaken for *E. t. extimus* could include sun-bleached or worn individuals of other races, as well as pale variants of *E. t. adastus*, intergrades between the latter and *E. t. extimus*, and carelessly-examined *E. t. campestris*. If so misidentified, such instances could help explain records of *E. t. extimus* from areas outside its known breeding range, such as the northern two-thirds of Colorado (Bailey and Niedrach 1965) and Texas east of the Trans-Pecos region (Oberholser 1974). As for New Mexico, I am dubious of *E. t. extimus* records from the eastern plains, such as two specimens reported in Hubbard (1987) from Roosevelt County. In addition, I have definitely reidentified two of the purported *E. t. extimus* from that report, one from San Juan County (= *E. t. adastus* > *extimus*) and another from Socorro County (= *E. t. extimus* > *adastus*). Of course, as mentioned earlier, we do not have the luxury of reexamining *E. t. extimus* (or other subspecies) reported by Y&F, so their identifications cannot be reassessed in light of potential sources of misidentification. Given this and their flawed methodology, I see no reason to regard their findings on this form as any more acceptable than those on the other races reported in their paper. As a final point, Y&F make no mention of the differences between their findings on the various subspecies and the specimen record as discussed above.

While the need for this would not have been obvious as regards *E. t. extimus* and perhaps even *E. t. adastus*, this could hardly have been the case with *E. t. traillii* and especially *E. t. brewsteri*.

To summarize, geographic variation in the Willow Flycatcher mainly involves subtle differences in plumage coloration, concerning which taxonomists disagree in terms of the number of subspecies that should be recognized. Anyone contemplating identifying these subspecies should do so with these caveats in mind, as well as by approaching the process through the use of specimen comparisons—preferably in the laboratory under controlled conditions and with proper training in alpha-taxonomic procedures. Given that Y&F's approach appears to have been otherwise, I submit that their field identification of these subspecies cannot be regarded as a bona fide assessment of this parameter in the birds they processed in New Mexico in 1994 and 1995. Furthermore, for those that would use their subspecific findings, I urge them to do so with extreme caution to say the least. Beyond this, I would like to state that as an alpha-taxonomist, I am dismayed that a study with such a flawed approach to subspecies identification could make its way into print in a major ornithological journal. To wit, ornithology has come to rely almost entirely on non-specimen data for monitoring the distribution and status of birds on this planet. While not necessarily a bad thing, sometimes we may fail to recognize the very real limitations of such data. No better example of this exists than as regards the identification of difficult taxa, of which subspecies in *Empidonax traillii* provide a perfect case in point.

LITERATURE CITED

- ALDRICH, J. W. 1951. A review of the races of Traill's Flycatcher. *Wilson Bull.* 63:192–197.
- BAILEY, A. M. AND R. J. NIEDRACH. 1965. *Birds of Colorado*, vol. 2. Denver Museum of Natural History, Denver, Colorado.
- BEHLE, W. C. 1985. Utah birds: geographic variation and systematics. *Utah Mus. Nat. Hist. Occ. Pub.* 5:1–147.
- BROWNING, M. R. 1993. Comments on the taxonomy of *Empidonax traillii* West. *Birds* 24:241–257.
- HUBBARD, J. P. 1987. The status of the Willow Flycatcher in New Mexico. Unpublished report, New Mexico Dept. of Game and Fish, Santa Fe.
- MAYR, E. AND L. L. SHORT, JR. 1970. *Species taxa of North American birds*. Nuttall Ornithological Club, Cambridge, Massachusetts.
- MONSON, G. AND A. R. PHILLIPS. 1981. Annotated checklist of the birds of Arizona. Univ. of Arizona Press, Tucson.
- OBERHOLSER, H. C. 1974. *The birdlife of Texas*, vol. 2. Univ. of Texas Press, Austin.
- PHILLIPS, A. R. 1948. Geographic variation in *Empidonax traillii*. *Auk* 65:507–514.
- PHILLIPS, A. R., J. MARSHALL, AND G. MONSON. 1964. *The birds of Arizona*. Univ. Arizona Press, Tucson.
- SMITHE, F. B. 1975. *Naturalist's color guide*. American Museum of Natural History, New York.
- SUTTON, G. M. 1967. *Oklahoma birds*. Univ. of Oklahoma Press, Norman.
- TRAYLOR, M. A., JR. 1979. Check-list of birds of the world, vol. 8. Harvard Univ., Cambridge, Massachusetts.
- UNITT, P. 1987. *Empidonax traillii extimus*: an endangered subspecies. *West. Birds* 18:137–162.
- YONG, W. AND D. M. FINCH. 1997. Migration of the Willow Flycatcher along the Middle Rio Grande. *Wilson Bull.* 109:253–268.
- WETMORE, A. 1972. The birds of the Republic of Panama, part 3. *Smithsonian Misc. Coll.* 150(3):1–631.